Published online 2017 January 24.

Letter

Stainless Steel Crowns in Pediatric Restorative Dentistry: Applications and Advantages

Nahid Ramazani^{1,*}

¹Children and Adolescents Health Research Center, Oral and Dental Disease Research Center, Pediatric Dentistry Department, Zahedan University of Medical Sciences, Zahedan, IR Iran

^{*} *Corresponding author*: Nahid Ramazani, Children and Adolescents Health Research Center, Oral and Dental Disease Research Center, Pediatric Dentistry Department, Zahedan University of Medical Sciences, Zahedan, IR Iran. Tel: +98-9155009085; +98-5433414005, E-mail: ramazani_nahid@yahoo.com

Received 2016 October 06; Revised 2016 November 23; Accepted 2017 January 10.

Keywords: Crown, Pediatric Dentistry, Tooth

Dear Editor,

Many children suffer from dental caries as the most common infectious disease (1-3). Consequently, proper dental restoration is one of the most critical issues in pediatric dentistry. Stainless steel crowns (SSCs) are characterized by full crown coverage and increased caries prevention through protecting the entire tooth; they are also more durable than multi-surface amalgam restorations (4). SSCs are prefabricated shells in different sizes, composed of metals such as iron, nickel, and chromium.

Despite major advancements in the development of modalities in pediatric dentistry (such as laser technology) (5), there are certain circumstances which necessitate the application of SSCs. In general, crowns are routinely used for restoring pulp-treated teeth. The concept of full coverage is to eliminate the risk of breaking tooth to buccal and lingual halves due to reduced dentin strength.

Another indication of SSCs is dental treatment under general anesthesia. In dental rehabilitation under general anesthesia, the approach is to restore teeth more with SSCs and less with other dental materials (4). Children with hypophosphatemia, who develop spontaneous dental abscesses, are also among cases requiring SSCs (6). Other indications include heritable dental defects, such as amelogenesis imperfecta, dentinogenesis imperfecta, and environmentally-induced defects such as enamel hypoplasia.

In pediatric dentistry, the strategy is to deliver efficient treatment and simultaneously reduce the time of operation (7). Application of SSCs reduces the time of operation and promotes the child's cooperation. In fact, compared to other restoration methods, SSCs placement is less disturbing for the patient.

Despite various advantages of crowns, microleakage occurs in the marginal area due to the prefabricated entity. In in vitro studies, microleakage has been reported with respect to the extent of trace material (eg, dye) penetration in the interface between the tooth and the material (8). However, this drawback is not critical, compared to various advantages of crowns. In general, restoration cannot jeopardize the treatment and guarantees the life span of primary teeth.

Parental satisfaction plays a fundamental role in dental health care. Esthetically, parents may not consent to a crown-included treatment plan; therefore, dentists can alleviate their concern by explaining the benefits of SSCs. Children also agree with their parents and accept the plan by receiving proper and age-appropriate information from the parents. To mask the metallic appearance of crowns, a modification, called pre-veneered SSC, is available. These crowns have a composite-bonded veneer to produce a more pleasant appearance, although they require more preparation and are more expensive.

The high cost of SSCs is another reason for rejecting this restoration method; in this regard, cost-effectiveness should be taken into account. Accordingly, dentists have to explain that restorations other than SSCs may result in good outcomes for only one-surface or two-surface restorations. Parents should be also informed that although SSCs are more expensive compared to other restorations, treatment is certainly worth the cost.

Studies have measured post-treatment pain following dental treatments (9). In fact, immediate post-treatment pain and discomfort are inevitable consequences of SSCs. It has been revealed that SSCs cause significantly more discomfort than extraction in patients (10). Nevertheless, post-operative pain can be well controlled by pain killers.

References

- Ramazani N. Child dental neglect: a short review. Int J High Risk Behav Addict. 2014;3(4):21861. doi: 10.5812/ijhrba.21861. [PubMed: 25741483].
- 2. Ramazani N, Sadeghi P. Bacterial Leakage of Mineral Trioxide Aggregate, Calcium-Enriched Mixture and Biodentine as Furcation Perfora-

Copyright © 2017, Growth & Development Research Center. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribute the material just in noncommercial usages, provided the superior original work is properly cited. tion Repair Materials in Primary Molars. *Iran Endod J.* 2016;**11**(3):214–8. doi: 10.7508/iej.2016.03.013. [PubMed: 27471534].

- 3. Ramazani N, Rezaei S. Evaluation of the prevalence of clinical consequences of untreated dental caries using PUFA/pufa index in a group of Iranian children. *Iran J Pediatr.* 2016;**27**(1) doi: 10.5812/ijp.5016.
- Ramazani N. Different Aspects of General Anesthesia in Pediatric Dentistry: A Review. *Iran J Pediatr.* 2016;26(2):2613. doi: 10.5812/ijp.2613. [PubMed: 27307962].
- 5. Ramazani N, Ahmadi R, Daryaeian M. Oral and dental laser treatments for children: applications, advantages and considerations. *J Lasers Med Sci.* 2012;**3**(1):44–9.
- 6. Ramazani N. Hypophosphatemic rickets and its dental significance. *Iran J Pediatr.* 2013;**23**(6):714. [PubMed: 24910759].
- 7. Ramazani N, Mohammadi A, Amirabadi F, Ramazani M, Ehsani F. In vitro investigation of the cleaning efficacy, shaping ability, prepara-

tion time and file deformation of continuous rotary, reciprocating rotary and manual instrumentations in primary molars. *J Dent Res Dent Clin Dent Prospects*. 2016;**10**(1):49–56. doi: 10.15171/joddd.2016.008. [PubMed: 27092215].

- Zarenejad N, Asgary S, Ramazani N. Coronal microleakage of three different dental biomaterials as intra-orifice barrier during nonvital bleaching. *Dent Res J (Isfahan)*. 2015;12(6):581–8. doi: 10.4103/1735-3327.170582.
- Ramazani M, Hamidi MR, Moghaddamnia AA, Ramazani N, Zarenejad N. The Prophylactic Effects of Zintoma and Ibuprofen on Postendodontic Pain of Molars with Irreversible Pulpitis: A Randomized Clinical Trial. *Iran Endod J.* 2013;8(3):129–34. [PubMed: 23922575].
- Staman NM, Townsend JA, Hagan JL. Observational study: discomfort following dental procedures for children. *Pediatr Dent.* 2013;35(1):52– 4. [PubMed: 23635899].

Iran J Pediatr. 2017; 27(3);e9272.