



The Relationship Between Early Weight Z-Score Changes and Nutritional Differences in Very Low Birth Weight Infants During Hospital Stay

Chu-yi Guo,¹ Mu-xue Yu,^{1*} Lin-xia Deng,¹ Wang-kai Liu,¹ and Si-qi Zhuang¹

¹Department of Pediatrics, The First Affiliated Hospital, Sun Yat-sen University, Guangzhou, China

*Corresponding author: Mu-xue Yu, Department of Pediatrics, The First Affiliated Hospital, Sun Yat-sen University, Guangzhou, Guangdong Province 510080, China. E-mail: yumuxue@163.com

Received 2017 October 06; Revised 2018 January 19; Accepted 2018 March 11.

Abstract

Background: Advances in neonatal care have resulted in improved survival for very low birth weight infants (VLBWIs). However, the main factors that lead to early weight Z-score [standard deviation score (SDS)] changes during hospital stay in VLBWIs still need to be studied more.

Objectives: In this study we aimed to investigate the differences in specific factors that relate to VLBWIs' early weight Z-score changes as the scientific basis for clinical practice of VLBWIs during the entire hospital stay.

Methods: We retrospectively reviewed baseline characteristics, nutrition management, clinical outcomes, and complications between non-catch-down (NCD) (Δ weight SDS ≥ -0.67) and catch-down (CD) groups (Δ weight SDS < -0.67) in VLBWIs during hospital stay.

Results: A total of 227 infants (NCD group = 117, CD group = 110) met the inclusion criteria. No significant differences were found in baseline characteristics between the two groups. The NCD group received higher parenteral lipids (g/kg/d) from day 5 to day 7 and higher enteral feeding volume (mL/kg/d) from day 5 to week 5. The NCD group had higher protein:energy ratios at day 2 to day 4 and day 7 and higher total energy (kcal/kg/d) during the first six weeks. The NCD group had lower incidence of bronchopulmonary dysplasia (BPD) and parenteral nutrition-associated cholestasis (PNAC).

Conclusions: This study showed that adequate parenteral lipid and energy at the first week of life and earlier enhanced enteral feeding volume might significantly improve the VLBWIs' weight Z-scores during hospital stay. The VLBWIs with catch-down growth might be more prone to developing BPD and PNAC.

Keywords: Parenteral Nutrition, Enteral Nutrition, Very Low Birth Weight Infant

1. Background

The preterm infants with birth weight less than 1500 g were defined as very low birth weight preterm infants (VLBWIs). VLBWIs may have suck-swallow un-coordination, and deeply immature gastric and intestinal functions (1). Therefore, full enteral nutrition (EN) cannot be established immediately. Parenteral nutrition (PN) is a 'bridge' that establishes enteral nutrient intakes and improves the VLBWIs' growth (2). Previous studies suggested that most VLBWIs compared with the median birth weight of a reference fetus with the same postmenstrual age did not receive catch-up growth in neonatal intensive care unit (NICU) during hospitalization (3, 4). This deficiency manifested as a growth curve falling and a falling Z-score [standard deviation score (SDS)] during the early postnatal period. Z-

score (SDS) is used to describe the size (weight, length, and head circumference) of an infant compared with other infants of the same age and sex (5). For example, positive SDS is greater than the 50th percentile, 0 SDS is equivalent to the 50th percentile and negative SDS is less than the 50th percentile on a growth curve. A Z-score change of zero represents unchanged or stable growth status. The Z-score was considered to be an indicator to evaluate the individual or groups' growth. Ong et al. (6) found that 0.67 SDS represents the width of each percentile band on standard growth charts, i.e., 2nd to 9th percentile, 9th to 25th, 25th to 50th, etc. It could be a dynamic index to evaluate the growth status of preterm infants.

In today's NICUs, especially in developing countries such as China, the problem of postnatal growth failure is still serious in VLBWIs (7). In this retrospective study, we

used Δ weight SDS between birth and hospital discharge to investigate the differences in specific factors that relate to VLBWIs' weight Z-score changes during the entire hospital stay.

2. Methods

2.1. Study Population

Following approval of the institutional review board, the medical record was searched to identify patients evaluated at the First Affiliated Hospital of Sun Yat-sen University with a diagnosis code. This retrospective study was performed at a single university-based NICU (the First Affiliated Hospital of Sun Yat-sen University). The nutritional protocol and recordkeeping were provided as standard care in our department. The university hospital perinatal database was queried to identify and obtain the demographics of VLBWIs (birth weight < 1500 g, gestational age < 37 weeks, and hospitalized within 24 h of birth) admitted from 1 June 2010 to 30 March 2017. Infants who had cardiac anomalies, congenital gastrointestinal anomalies, or hereditary metabolic diseases; were transferred after 24 postnatal hours, died in the hospital, had a hospital stay of < 15 days, or had unstable vital signs upon hospital discharge were excluded.

Infant demographics, including Z-score (SDS) for weight, length, and head circumference between birth and discharge using the Fenton 2013 growth chart (8), as well as age upon initiation, were obtained. All infants were divided into two groups (non-catch-down [NCD] group: Δ weight SDS \geq -0.67; catch-down [CD] group: Δ weight SDS < -0.67). All data were compared between the two groups for further evaluation.

2.2. Perinatal Characteristics and Complications

The data of maternal characteristics and complications that affected the wellbeing of an infant, such as gemellary pregnancy, pregnancy-induced hypertension, steroid use, delivery route, premature rupture of membrane, fetal distress, in vitro fertilization and embryo transfer, intracytoplasmic sperm injection, twin-twin transfusion syndrome and multifetal reduction were analyzed. Additionally, the data of infants' characteristics and complications such as small for gestational age (SGA), bronchopulmonary dysplasia (BPD), sepsis, necrotizing enterocolitis (NEC), intraventricular hemorrhage (ICH), retinopathy of prematurity (ROP), extrauterine growth restriction (EUGR), and parenteral nutrition associated cholestasis (PNAC) for the two groups were also recorded. Total days of return to birth weight and hospital stay were calculated and analyzed.

2.3. Anthropometric Parameters at Birth and Discharge

The infants' weight, length, and head circumference and Z-score (SDS) for weight, length, and head circumference at birth and discharge were also obtained and studied.

2.4. Nutrition Management

Daily intakes of all nutrients including parenteral nutrition (PN) and enteral nutrition (EN) recorded in patient charts according to NICU protocol were assessed. After birth, PN was initiated within 24 hours and contained glucose, amino acids, and lipids in our NICU. Energy was increased daily based on metabolic tolerance. The protein content and energy density of enteral nutritional intake were calculated according to the guideline (2). Formula milk composition was calculated by using the manufacturers' datasheets. Total days of PN and laboratory biochemical indicators (blood urea nitrogen [BUN], creatinine [Cr], aspartate transaminase [AST], and alanine transaminase [ALT]) during hospitalization were also obtained.

2.5. Statistical Analyses

Baseline characteristics, daily changes in protein and energy intake with PN and EN, clinical outcomes, and complications during the NICU stay were compared between the two groups. Continuous variables are presented as mean \pm standard deviation or median or quartile range (P25, P75). Student's t-test was used to compare quantitative data with a normal distribution. Parameters without a normal distribution were analyzed by the Mann-Whitney U test. Categorical variables such as maternal and clinical outcomes of the two groups are presented as a percentage that was analyzed by chi-square test and Fisher's exact test. Calculations were performed using SPSS software version 17.0 (SPSS Inc, Chicago, IL, USA). The level of significance was set at a $P < 0.05$.

3. Results

3.1. Perinatal Characteristics and Complications

A total of 227 VLBWIs (NCD group = 117, CD group = 110) met the inclusion criteria. Perinatal characteristics and complications are listed in Table 1. No significant differences in maternal characteristics and complications were found between the two groups. The incidence of most common complications (except for BPD and PNAC) related to prematurity and mortality was not significantly different between the two groups. The incidence of BPD (0.348 vs 0.518, respectively; $P < 0.05$) and PNAC (0.196 vs 0.341, respectively; $P < 0.05$) were significantly lower in the NCD group than in the CD group. The mean PN duration, length

of hospital stay, and mean days of return to BW were significantly shorter in the NCD group ($P < 0.05$).

3.2. Anthropometric Parameters at Birth and Discharge

The groups were compared for anthropometric parameters and Z-score (SDS) at birth and discharge (Table 2). No significant differences were found between the two groups at birth. The anthropometric parameters at discharge were significantly higher in the NCD group ($P < 0.05$). Additionally, the SDS for weight and length and head circumference at discharge was significantly higher in the NCD group ($P < 0.05$).

3.3. PN and EN Intake

Daily changes in protein and energy intake from birth to the sixth week of life are presented in Figures 1 and 2. During the first week of life, the daily average parenteral amino acid (g/kg/d) and glucose (g/kg/d) administered to the two groups were not significantly different, while parenteral lipids (g/kg/d) administered to the NCD group were significantly higher than those of the CD group from day 5 to day 7 ($P < 0.05$) (Figure 1A-1C). After the first week of life, the NCD group received significantly lower parenteral glucose (g/kg/d) from week 2 to week 3 and lower parenteral amino acid (g/kg/d) at week 3 ($P < 0.05$) (Figure 1A and 1B). The NCD group received higher mean daily enteral feeding volume (mL/kg/d) than the CD group from day 5 to week 5 ($P < 0.05$) (Figure 1D).

The NCD group had a greater mean daily EN protein intake (mL/kg/d) from day 5 to week 5 ($P < 0.05$) (Figure 2A). The NCD group had higher parenteral energy (kcal/kg/d) than that of the CD group from day 5 to day 7 ($P < 0.05$) (Figure 2B). Moreover, the NCD group had a significantly increased energy intake (kcal/kg/d) during the first six weeks of life ($P < 0.05$), except for the first four days (Figure 2B). The P:E ratio was higher at the first week and significantly higher at day 2 to day 4 and day 7 in NCD group ($P < 0.05$) (Figure 2C). However, from week 2 to week 3, the P:E ratio was significantly higher in the CD group ($P < 0.05$) (Figure 2C).

3.4. Main Laboratory Test

No differences were found between the two groups regarding mean values of the main laboratory tests from the first to the sixth week of life (Table 3).

4. Discussion

Our study demonstrated that the NCD group was given more parenteral lipid and total energy at the first week, and received earlier enhanced enteral feeding volume

than the CD group during hospital stay. This study revealed that parenteral and enteral nutrition might play the most important role in the postnatal growth of VLBWIs during hospitalization. The improved weight Z-score of VLBWIs during hospital stay might associate with the earlier adequate parenteral lipid and energy intake at the first week of life and enhanced enteral feeding volume during hospital stay. The infants with significant decline in weight Z-scores might be more prone to developing BPD and PNAC.

Parenteral lipids are one of the most important energy providers (9), that can provide calories required for the process of protein synthesis. Increased parenteral lipids significantly improve nitrogen utilization and retention, resulting in positive nitrogen balance in preterm VLBWIs (10). In our study, we found that parenteral lipids and total parenteral energy administered to the NCD group were significantly higher than those administered to the CD group from day 5 to week 1. Similar to our findings, researches on parenteral lipids had shown that higher dosages of intravenous lipids helped to decrease postnatal weight loss and have an earlier birth weight regain (11, 12). Dit Trolli et al. (13) demonstrated that early lipid supply (lipids were initiated at the first three days after birth and increased 0.5 to 1 g/kg/d to reach 4 g/kg/d) during the first two weeks of life could improve the weight gain from birth to day 28 of life and ameliorate the neurological development at a corrected age of one year for preterm VLBWIs. Moreover, recent randomized controlled trials (RCTs) have indicated that increasing parenteral energy intakes during the first few days of life significantly improve the preterm infants' head growth (14, 15). Thus, our results add to the evidence that earlier administration of adequate parenteral lipids and energy may help to meet the numerous nutritional needs of VLBWIs.

In our study, the mean daily enteral feeding volume administered to the NCD group was higher than the CD group from day 5 to week 5. This finding indicated that an earlier and enhanced enteral feeding plays an important role in the growth for VLBWIs. Early EN is known to decrease gut atrophy and intestinal permeability (16) and has been associated with improved postnatal growth (17). A recent RCT (18) demonstrated a significantly better weight gain during hospitalization and adequate growth of head circumference in the enhanced feeding group. Enhanced enteral feeding might also improve weight gain during hospitalization (19). Morgan et al. (20) suggested that advancing enteral feed volumes (30 to 40 mL/kg/d) compared to slow enteral feed volumes (15 to 24 mL/kg) does not increase the incidence of NEC or death in VLBWIs. Additionally, they found that the slow enteral feed volumes might delay in establishing full enteral feeds and increase the risk of infection in VLBWIs. Guidelines for feeding VLBWIs of McMas-

Table 1. Main Perinatal Complications and Complications^a

	NCD Group (n = 117)	CD Group (n = 110)	t/z/ χ^2	P Values
Maternal				
Gemellary pregnancy	58 (49.57)	60 (54.55)	0.562	0.454
Pregnancy-induced hypertension	29 (24.79)	30 (27.27)	0.182	0.669
Prenatal steroids	62 (52.99)	49 (44.55)	1.619	0.203
Cesarean delivery	21 (17.95)	23 (20.09)	0.318	0.573
PROM	41 (35.04)	35 (31.82)	0.265	0.607
Fetal distress	18 (15.38)	20 (18.18)	0.318	0.573
Meconium stained amniotic fluid	12 (10.26)	14 (12.73)	0.341	0.559
Placenta praevia	11 (9.40)	10 (9.09)	0.007	0.936
IVF-ET	25 (21.37)	26 (23.64)	0.168	0.682
ICSI	4 (3.42)	3 (2.73)	0.091	0.763
TTTS	10 (8.55)	15 (13.64)	1.498	0.221
Multifetal reduction	9 (7.69)	10 (9.09)	0.145	0.704
Infant				
GA, w	30.2 (29.0, 32.5)	30.4 (28.6, 32.3)	1.109	0.282
SGA	42 (35.90)	44 (40.00)	0.406	0.524
RDS	104 (88.89)	100 (90.91)	0.254	0.614
Apnea	27 (23.08)	29 (26.36)	0.330	0.566
Pulmonary infection	45 (38.46)	56 (50.91)	3.557	0.059
BPD	49 (41.88)	65 (59.09)	6.717	0.010
NEC	23 (19.65)	22 (20.00)	0.004	0.949
Sepsis	36 (30.77)	31 (28.18)	0.344	0.558
ICH	36 (30.77)	29 (26.36)	0.539	0.463
ROP	38 (32.48)	34 (30.91)	0.064	0.800
PNAC	39 (33.33)	55 (50.00)	0.210	0.646
EUGR	28 (28.3)	54 (60.0)	15.553	0.000
Duration of PN, d	31 (24.41)	40 (29.52)	3.487	0.000
Return to BW, d	6 (1.8)	8 (1.11)	2.256	0.028
Hospital stay, d	39 (30.52)	49 (38.62)	3.259	0.001

Abbreviations: BL, birth length; BW, birth weight; BPD, bronchopulmonary dysplasia; EUGR, extrauterine growth restriction; GA, gestational age; HC, head circumference; ICH: intracranial hemorrhage; ICSI, Intracytoplasmic sperm injection; IVF-ET, in vitro fertilization and embryo transfer; NEC, necrotizing enterocolitis; PN, parenteral nutrition; PNAC, parenteral nutrition associated cholestasis; PROM, premature rupture of membrane; RDS, respiratory distress syndrome; ROP, retinopathy of prematurity; SGA, small for gestational age; TTTS, twin-twin transfusion syndrome.

^aValues are expressed as No. (%) or median (P25, P75).

ter University (21) suggested that infants weighing < 1000 g at birth reach full enteral feeding (~150 to 180 mL/kg/d) by about two weeks in and by about one week in infants weighing 1000 to 1500 g by using evidence-based feeding protocols. Furthermore, we found that the NCD group received lower parenteral protein and energy intake after two weeks. We inferred that the NCD group had received higher EN during the first week, then gradually reduced PN and transitioned to full enteral feeding. Meanwhile, this study

also suggested that enhanced enteral feeding volume during the first week might help to shorten the time of PN duration and achieve full enteral feeding quickly.

We noticed that the NCD group was administered a higher total protein intake from day 5 to week 5 after birth and had a higher P:E ratio during the first week of life. Protein is most important nutrient for the growth of preterm VLBWIs as the main functional and structural component of the body cells and structure (2). It has been estimated

Table 2. Anthropometric Parameters at Birth and Discharge^a

	NCD Group (n = 117)	CD Group (n = 110)	t/z	P Value
Weight at birth, kg	1.22 ± 0.23	1.19 ± 0.21	1.024	0.153
Weight SDS at birth	-0.64 ± 0.51	-0.56 ± 0.45	1.255	0.105
Length at birth, cm	27.6 ± 1.6	27.5 ± 1.7	0.457	0.324
Length SDS at birth	-0.64 ± 0.37	-0.62 ± 0.34	0.423	0.336
HC at birth, cm	37.6 ± 3.4	37.9 ± 3.5	0.655	0.257
HC SDS at birth	-0.37 ± 0.29	-0.34 ± 0.25	0.832	0.203
Weight at discharge, kg	2.48 ± 0.41	2.22 ± 0.81	3.022	0.003
Weight SDS at discharge	-0.75 ± 0.35	-0.97 ± 0.87	2.471	0.015
Length at discharge, cm	45.8 ± 2.8	43.4 ± 3.6	4.256	0.000
Length SDS at discharge	-0.41 ± 0.65	-1.29 ± 0.48	4.707	0.000
HC at discharge, cm	31.4 ± 2.5	30.4 ± 1.7	3.542	0.000
HC SDS at discharge	-0.81 ± 0.76	-1.12 ± 0.64	3.314	0.001

Abbreviation: HC, head circumference.


^aValues are expressed as means ± SD.

Figure 1. Mean daily PN, EN intakes and protein energy ratio from birth to 6th week from birth to the sixth week. During the first week of life, the daily average parenteral amino acid (g/kg/d) and glucose (g/kg/d) administered to the 2 groups were not significantly different, while parenteral lipids (g/kg/d) administered to the NCD group were significantly higher than those of the CD group from day 5 to day 7 (A-C). From week 2 to week 3, the NCD group was administered significantly lower parenteral amino acid than the CD group (A). At week 3, the NCD group was administered significantly lower parenteral amino acid (g/kg/d) than the CD group (B). The NCD group was administered a higher mean daily enteral feeding volume (mL/kg/d) than the CD group from day 5 to week 5 (D). Error bars represent standard deviations; *P < 0.05; **P < 0.01.

that early high protein intake could improve weight gain and the other growth outcomes at discharge (22). The European society for paediatric gastroenterology hepatology and nutrition (ESPGHAN) recommended that the infants

up to 1000 g should receive 4.0 to 4.5 g/kg/day protein, and the infants from 1000 to 1800 g should receive 3.5 to 4.0 g/kg/day protein that will meet the needs of most preterm infants (23). The types of EN such as human milk and for-




Figure 2. Total protein intake and total energy intake via PN and EN from birth to the sixth week. A, the NCD group had a significantly increased total protein intake (g/kg/d) during day 5 to day 6 and week 4 to week 5. The NCD group had a greater mean daily enteral protein intake from day 5 to week 5. The CD group had increased parenteral protein intake week 2 to week 5 and significantly higher than the NCD group during week 3; B, the NCD group had a significantly increased energy intake (kcal/kg/d) during the first 6 weeks of life, except for the first four days. The NCD group had higher parenteral energy than those of the CD group from day 5 to day 7. The CD group had increased parenteral energy intake week 2 to week 5 and significantly higher than the NCD group from week 2 to week 4. The P:E ratio was higher at the first week and significantly higher at day 2 to day 4 and day 7 in the NCD group (2C). During week 2 to week 3, the P:E ratio was significantly lower in the NCD group (C). Error bars represent standard deviations; * $P < 0.05$, ** $P < 0.01$ for EN; $\Delta P < 0.05$ for PN; # $P < 0.05$ for total protein/energy intake; & $P < 0.05$, && $P < 0.01$ for the P:E ratio.

mula milk can meet the VLBWIs' protein and energy needs (2). Moreover, early and reasonable energy intake could improve the preterm infants' growth and development (24). Stephens et al. (25) found that increasing first-week protein and energy intake resulted in higher mental develop-

mental index scores of extremely low birth weight infants (ELBWIs) at 18 months. Additionally, we found that the NCD group received a lower P:E ratio at 2 to 3 weeks. We speculated that the reduction of P:E ratio was due to the circumstance that NCD group had received higher total protein

Table 3. Mean Laboratory Values from the First Week to the Sixth Week of Life^a

	NCD Group (n = 117)	CD Group (n = 110)	t	P Value
Week 1				
BUN, mmol/L	4.14 ± 1.45	4.09 ± 1.60	2.247	0.805
Cr, umol/L	73.23 ± 8.22	74.66 ± 8.75	1.267	0.103
AST, U/L	30.86 ± 6.45	30.32 ± 7.10	0.599	0.275
ALT, U/L	13.42 ± 2.18	13.48 ± 1.91	0.221	0.413
Week 2				
	(n = 117)	(n = 110)		
BUN, mmol/L	4.23 ± 2.04	4.12 ± 1.94	0.416	0.678
Cr, umol/L	55.47 ± 5.12	54.69 ± 5.67	1.086	0.139
AST, U/L	27.37 ± 3.28	26.92 ± 7.63	0.571	0.284
ALT, U/L	8.59 ± 3.64	8.33 ± 3.81	0.525	0.300
Week 3				
	(n = 106)	(n = 102)		
BUN, mmol/L	3.24 ± 1.17	3.32 ± 1.02	0.531	0.596
Cr, umol/L	34.12 ± 4.34	35.46 ± 4.68	0.543	0.294
AST, U/L	25.10 ± 10.52	26.44 ± 9.73	0.954	0.171
ALT, U/L	9.05 ± 4.67	9.58 ± 5.56	0.752	0.453
Week 4				
	(n = 69)	(n = 75)		
BUN, mmol/L	3.04 ± 1.28	3.26 ± 1.01	1.138	0.257
Cr, umol/L	36.53 ± 6.21	37.45 ± 6.70	0.855	0.197
AST, U/L	25.24 ± 5.53	25.59 ± 5.56	0.378	0.353
ALT, U/L	9.12 ± 3.48	9.14 ± 4.25	0.031	0.488
Week 5				
	(n = 40)	(n = 53)		
BUN, mmol/L	3.68 ± 1.59	3.54 ± 1.39	0.452	0.652
Cr, umol/L	31.32 ± 5.95	31.78 ± 6.01	0.368	0.357
AST, U/L	24.74 ± 3.24	25.50 ± 3.15	1.133	0.131
ALT, U/L	16.39 ± 4.68	16.68 ± 3.85	0.319	0.375
Week 6				
	(n = 30)	(n = 39)		
BUN, mmol/L	3.20 ± 1.96	3.33 ± 1.55	0.3.8	0.759
Cr, umol/L	31.58 ± 3.29	30.71 ± 4.77	0.895	0.187
AST, U/L	27.24 ± 3.17	27.58 ± 3.63	0.415	0.340
ALT, U/L	18.69 ± 6.84	19.74 ± 6.40	0.650	0.259

Abbreviations: ALT, alanine transaminase; AST, aspartate transaminase; BUN, blood urea nitrogen; Cr, creatinine.

^aValues are expressed as means ± SD.

intake with higher P:E ratio at the first week, then it gradually reduced with lower proportion of protein in PN.

In this our study, the NCD group had lower incidence of BPD suggesting that BPD was closely related to greater nutritional intake. The infant with BPD has higher oxygen consumption and energy expenditure (26). Poor nutrient conditions might damage the process of immature lung tissue remodeling and develop severe chronic lung disease (26). Uberos et al. (27) suggested that lower enteral intake

of energy and total lipids during the first 14 days of life increase the risk of BPD in ELBWIs. Gianni et al. (28) showed that energy intake and severity of BPD were the major factors influencing the weight gain of preterm VLBWIs. Additionally, several studies have demonstrated that the infants who have a slower weight gain may be more prone to developing infections and BPD (29, 30).

In the meantime, we found that NCD group had a lower incidence of PNAC. Prematurity, low birth weight and long

PN duration in VLBWIs are often described as risk factors for developing PNAC (31, 32). In VLBWIs, the rate of PNAC has been reported to be as high as 50% (33). Longer PN duration is an independent risk factor for developing PNAC (34). Repa et al. (35) found that “aggressive” nutrition (amino acids and lipids concentrations of 2.0 g/kg/d and 1.0 g/kg/d initiated at the first day of life and increased stepwise by 0.5 g/kg/d to reach 4 g/kg/d; the increasing rate of EN was up to 20 mL/kg/d) was associated with a significant lower incidence of PNAC and improved postnatal growth of ELBWIs significantly. Previous studies provided adequate nutritional support through a combination of early PN and EN, followed by a progressive PN reduction, as enteral feeding volumes were steadily advanced to full EN (36, 37). Hamilton et al. (38) found that initiating enteral feeding within 24 hours of birth was associated with a significantly lower number of PN days. Our study suggested that in VLBWIs early administration of an enhanced EN nutritional regimen might reduce the incidence of intrahepatic cholestasis because of a shorter PN duration during hospitalization.

This study may help to add new idea to the management of VLBWIs in NICU. The VLBWIs’ adequate growth during hospitalization might associate with adequate parenteral lipids, energy, and enhanced enteral feeding volume. However, the benefits of this nutritional regimen regarding long-term, especially neurodevelopmental and metabolic outcomes, remain to be proven in adequately-powered RCT. Practicing early PN and higher EN nutrient supplementation requires further study especially in developing countries.

4.1. Conclusions

In conclusion, this study shows that the VLBWIs’ Z-score changes might significantly associate with parenteral and enteral nutrition during hospitalization. The VLBWIs’ weight Z-score might be significantly improved by the earlier adequate parenteral lipid, energy intake, and earlier enhanced enteral feeding volume. Additionally, the infants with significant decline in weight Z-scores might be more prone to developing BPD and PNAC.

Acknowledgments

This study was financially supported by science and technology planning project of Guangdong province, China (No. 2012B061700069) and natural science foundation of Guangdong province, China (No. 2015A030313148).

References

1. Fanaro S. Feeding intolerance in the preterm infant. *Early Hum Dev.* 2013;**89** Suppl 2:S13–20. doi: [10.1016/j.earlhumdev.2013.07.013](https://doi.org/10.1016/j.earlhumdev.2013.07.013). [PubMed: [23962482](https://pubmed.ncbi.nlm.nih.gov/23962482/)].
2. Koletzko B, Uauy R. *Nutritional care of preterm infants*. Switzerland: Karger; 2014.
3. Ehrenkranz RA, Younes N, Lemons JA, Fanaroff AA, Donovan EF, Wright LL, et al. Longitudinal growth of hospitalized very low birth weight infants. *Pediatrics.* 1999;**104**(2 Pt 1):280–9. [PubMed: [10429008](https://pubmed.ncbi.nlm.nih.gov/10429008/)].
4. Horbar JD, Ehrenkranz RA, Badger GJ, Edwards EM, Morrow KA, Soll RF, et al. Weight Growth Velocity and Postnatal Growth Failure in Infants 501 to 1500 Grams: 2000–2013. *Pediatrics.* 2015;**136**(1):e84–92. doi: [10.1542/peds.2015-0129](https://doi.org/10.1542/peds.2015-0129). [PubMed: [26101360](https://pubmed.ncbi.nlm.nih.gov/26101360/)].
5. Olsen IE, Harris CL, Lawson ML, Berseth CL. Higher protein intake improves length, not weight, z scores in preterm infants. *J Pediatr Gastroenterol Nutr.* 2014;**58**(4):409–16. doi: [10.1097/MPG.0000000000000237](https://doi.org/10.1097/MPG.0000000000000237). [PubMed: [24231639](https://pubmed.ncbi.nlm.nih.gov/24231639/)].
6. Ong KK, Ahmed ML, Emmett PM, Preece MA, Dunger DB. Association between postnatal catch-up growth and obesity in childhood: prospective cohort study. *BMJ.* 2000;**320**(7240):967–71. [PubMed: [10753147](https://pubmed.ncbi.nlm.nih.gov/10753147/)]. [PubMed Central: [PMC27335](https://pubmed.ncbi.nlm.nih.gov/PMC27335/)].
7. Collaborative Group for the Nutritional G, Developmental Study on Very Low Birth Weight I. Postnatal growth of very low birth weight infants during hospitalization. *Zhonghua Er Ke Za Zhi.* 2013;**51**(1):4–11. [PubMed: [23527924](https://pubmed.ncbi.nlm.nih.gov/23527924/)].
8. Fenton TR, Kim JH. A systematic review and meta-analysis to revise the Fenton growth chart for preterm infants. *BMC Pediatr.* 2013;**13**:59. doi: [10.1186/1471-2431-13-59](https://doi.org/10.1186/1471-2431-13-59). [PubMed: [23601190](https://pubmed.ncbi.nlm.nih.gov/23601190/)]. [PubMed Central: [PMC3637477](https://pubmed.ncbi.nlm.nih.gov/PMC3637477/)].
9. Lapillonne A. Enteral and parenteral lipid requirements of preterm infants. *World Rev Nutr Diet.* 2014;**110**:82–98. doi: [10.1159/000358460](https://doi.org/10.1159/000358460). [PubMed: [24751623](https://pubmed.ncbi.nlm.nih.gov/24751623/)].
10. Salama GS, Kaabneh MA, Almasaeed MN, Alquran M. Intravenous lipids for preterm infants: a review. *Clin Med Insights Pediatr.* 2015;**9**:25–36. doi: [10.4137/CMPed.S21161](https://doi.org/10.4137/CMPed.S21161). [PubMed: [25698888](https://pubmed.ncbi.nlm.nih.gov/25698888/)]. [PubMed Central: [PMC4325703](https://pubmed.ncbi.nlm.nih.gov/PMC4325703/)].
11. Drenckpohl D, McConnell C, Gaffney S, Niehaus M, Macwan KS. Randomized trial of very low birth weight infants receiving higher rates of infusion of intravenous fat emulsions during the first week of life. *Pediatrics.* 2008;**122**(4):743–51. doi: [10.1542/peds.2007-2282](https://doi.org/10.1542/peds.2007-2282). [PubMed: [18829797](https://pubmed.ncbi.nlm.nih.gov/18829797/)].
12. Vlaardingerbroek H, Vermeulen MJ, Carnielli VP, Vaz FM, van den Akker CH, van Goudoever JB. Growth and fatty acid profiles of VLBW infants receiving a multicomponent lipid emulsion from birth. *J Pediatr Gastroenterol Nutr.* 2014;**58**(4):417–27. doi: [10.1097/MPG.0000000000000280](https://doi.org/10.1097/MPG.0000000000000280). [PubMed: [24667866](https://pubmed.ncbi.nlm.nih.gov/24667866/)].
13. di Trolli SE, Kermorvant-Duchemin E, Huon C, Bremond-Gignac D, Lapillonne A. Early lipid supply and neurological development at one year in very low birth weight (VLBW) preterm infants. *Early Hum Dev.* 2012;**88** Suppl 1:S25–9. doi: [10.1016/j.earlhumdev.2011.12.024](https://doi.org/10.1016/j.earlhumdev.2011.12.024). [PubMed: [22264437](https://pubmed.ncbi.nlm.nih.gov/22264437/)].
14. Morgan C, McGowan P, Herwitker S, Hart AE, Turner MA. Postnatal head growth in preterm infants: a randomized controlled parenteral nutrition study. *Pediatrics.* 2014;**133**(1):e120–8. doi: [10.1542/peds.2013-2207](https://doi.org/10.1542/peds.2013-2207). [PubMed: [24379229](https://pubmed.ncbi.nlm.nih.gov/24379229/)].
15. Herrmann KR, Herrmann KR. Early parenteral nutrition and successful postnatal growth of premature infants. *Nutr Clin Pract.* 2010;**25**(1):69–75. doi: [10.1177/0885433609359001](https://doi.org/10.1177/0885433609359001). [PubMed: [20130159](https://pubmed.ncbi.nlm.nih.gov/20130159/)].
16. Rouwet EV, Heineman E, Buurman WA, ter Riet G, Ramsay G, Blanco CE. Intestinal permeability and carrier-mediated monosaccharide absorption in preterm neonates during the early postnatal period. *Pediatr Res.* 2002;**51**(1):64–70. doi: [10.1203/00006450-200201000-00012](https://doi.org/10.1203/00006450-200201000-00012). [PubMed: [11756641](https://pubmed.ncbi.nlm.nih.gov/11756641/)].

17. Flidel-Rimon O, Friedman S, Lev E, Juster-Reicher A, Amitay M, Shinwell ES. Early enteral feeding and nosocomial sepsis in very low birth-weight infants. *Arch Dis Child Fetal Neonatal Ed.* 2004;**89**(4):F289-92. doi: [10.1136/adc.2002.021923](https://doi.org/10.1136/adc.2002.021923). [PubMed: [15210657](https://pubmed.ncbi.nlm.nih.gov/15210657/)]. [PubMed Central: [PMC1721698](https://pubmed.ncbi.nlm.nih.gov/PMC1721698/)].
18. Moltu SJ, Blakstad EW, Strommen K, Almaas AN, Nakstad B, Ronnestad A, et al. Enhanced feeding and diminished postnatal growth failure in very-low-birth-weight infants. *J Pediatr Gastroenterol Nutr.* 2014;**58**(3):344-51. doi: [10.1097/MPG.0000000000000220](https://doi.org/10.1097/MPG.0000000000000220). [PubMed: [24556755](https://pubmed.ncbi.nlm.nih.gov/24556755/)]. [PubMed Central: [PMC3940525](https://pubmed.ncbi.nlm.nih.gov/PMC3940525/)].
19. Leaf A, Dorling J, Kempley S, McCormick K, Mannix P, Linsell L, et al. Early or delayed enteral feeding for preterm growth-restricted infants: a randomized trial. *Pediatrics.* 2012;**129**(5):e1260-8. doi: [10.1542/peds.2011-2379](https://doi.org/10.1542/peds.2011-2379). [PubMed: [22492770](https://pubmed.ncbi.nlm.nih.gov/22492770/)].
20. Morgan J, Young L, McGuire W. Slow advancement of enteral feed volumes to prevent necrotising enterocolitis in very low birth weight infants. *Cochrane Database Syst Rev.* 2015;(10). CD001241. doi: [10.1002/14651858.CD001241.pub6](https://doi.org/10.1002/14651858.CD001241.pub6). [PubMed: [26469124](https://pubmed.ncbi.nlm.nih.gov/26469124/)].
21. Dutta S, Singh B, Chessell L, Wilson J, Janes M, McDonald K, et al. Guidelines for feeding very low birth weight infants. *Nutrients.* 2015;**7**(1):423-42. doi: [10.3390/nu7010423](https://doi.org/10.3390/nu7010423). [PubMed: [25580815](https://pubmed.ncbi.nlm.nih.gov/25580815/)]. [PubMed Central: [PMC4303848](https://pubmed.ncbi.nlm.nih.gov/PMC4303848/)].
22. Maggio L, Cota F, Gallini F, Lauriola V, Zecca C, Romagnoli C. Effects of high versus standard early protein intake on growth of extremely low birth weight infants. *J Pediatr Gastroenterol Nutr.* 2007;**44**(1):124-9. doi: [10.1097/01.mpg.0000237927.00105.f7](https://doi.org/10.1097/01.mpg.0000237927.00105.f7). [PubMed: [17204965](https://pubmed.ncbi.nlm.nih.gov/17204965/)].
23. Agostoni C, Buonocore G, Carnielli VP, De Curtis M, Darmaun D, Decsi T, et al. Enteral nutrient supply for preterm infants: commentary from the European Society of Paediatric Gastroenterology, Hepatology and Nutrition Committee on Nutrition. *J Pediatr Gastroenterol Nutr.* 2010;**50**(1):85-91. doi: [10.1097/MPG.0b013e3181adaee0](https://doi.org/10.1097/MPG.0b013e3181adaee0). [PubMed: [19881390](https://pubmed.ncbi.nlm.nih.gov/19881390/)].
24. Hay WW. Early postnatal nutritional requirements of the very preterm infant based on a presentation at the NICHD-AAP workshop on research in neonatology. *J Perinatol.* 2006;**26** Suppl 2:S13-8. doi: [10.1038/sj.jp.7211426](https://doi.org/10.1038/sj.jp.7211426). [PubMed: [16801962](https://pubmed.ncbi.nlm.nih.gov/16801962/)].
25. Stephens BE, Walden RV, Gargus RA, Tucker R, McKinley L, Mance M, et al. First-week protein and energy intakes are associated with 18-month developmental outcomes in extremely low birth weight infants. *Pediatrics.* 2009;**123**(5):1337-43. doi: [10.1542/peds.2008-0211](https://doi.org/10.1542/peds.2008-0211). [PubMed: [19403500](https://pubmed.ncbi.nlm.nih.gov/19403500/)].
26. Shlomai NO, Patole S. Nutrition in Preterm Infants with bronchopulmonary dysplasia. *Nutrition for the Preterm Neonate.* Springer; 2013.
27. Uberos J, Lardon-Fernandez M, Machado-Casas I, Molina-Oya M, Narbona-Lopez E. Nutrition in extremely low birth weight infants: impact on bronchopulmonary dysplasia. *Minerva Pediatr.* 2016;**68**(6):419-26. [PubMed: [25407225](https://pubmed.ncbi.nlm.nih.gov/25407225/)].
28. Gianni ML, Roggero P, Colnaghi MR, Piemontese P, Amato O, Orsi A, et al. The role of nutrition in promoting growth in pre-term infants with bronchopulmonary dysplasia: a prospective non-randomised interventional cohort study. *BMC Pediatr.* 2014;**14**:235. doi: [10.1186/1471-2431-14-235](https://doi.org/10.1186/1471-2431-14-235). [PubMed: [25241811](https://pubmed.ncbi.nlm.nih.gov/25241811/)]. [PubMed Central: [PMC4177697](https://pubmed.ncbi.nlm.nih.gov/PMC4177697/)].
29. Frank L, Sosenko IR. Undernutrition as a major contributing factor in the pathogenesis of bronchopulmonary dysplasia. *Am Rev Respir Dis.* 1988;**138**(3):725-9. doi: [10.1164/ajrccm/138.3.725](https://doi.org/10.1164/ajrccm/138.3.725). [PubMed: [3059876](https://pubmed.ncbi.nlm.nih.gov/3059876/)].
30. Prince A, Groh-Wargo S. Nutrition management for the promotion of growth in very low birth weight premature infants. *Nutr Clin Pract.* 2013;**28**(6):659-68. doi: [10.1177/0884533613506752](https://doi.org/10.1177/0884533613506752). [PubMed: [24163320](https://pubmed.ncbi.nlm.nih.gov/24163320/)].
31. Hsieh MH, Pai W, Tseng HI, Yang SN, Lu CC, Chen HL. Parenteral nutrition-associated cholestasis in premature babies: risk factors and predictors. *Pediatr Neonatol.* 2009;**50**(5):202-7. doi: [10.1016/S1875-9572\(09\)60064-4](https://doi.org/10.1016/S1875-9572(09)60064-4). [PubMed: [19856863](https://pubmed.ncbi.nlm.nih.gov/19856863/)].
32. Robinson DT, Ehrenkranz RA. Parenteral nutrition-associated cholestasis in small for gestational age infants. *J Pediatr.* 2008;**152**(1):59-62. doi: [10.1016/j.jpeds.2007.06.002](https://doi.org/10.1016/j.jpeds.2007.06.002). [PubMed: [18154901](https://pubmed.ncbi.nlm.nih.gov/18154901/)].
33. Burrin DG, Ng K, Stoll B, Saenz De Pipaon M. Impact of new-generation lipid emulsions on cellular mechanisms of parenteral nutrition-associated liver disease. *Adv Nutr.* 2014;**5**(1):82-91. doi: [10.3945/an.113.004796](https://doi.org/10.3945/an.113.004796). [PubMed: [24425726](https://pubmed.ncbi.nlm.nih.gov/24425726/)]. [PubMed Central: [PMC3884104](https://pubmed.ncbi.nlm.nih.gov/PMC3884104/)].
34. Yan W, Hong L, Wang Y, Feng Y, Lu L, Tao Y, et al. Retrospective Dual-Center Study of Parenteral Nutrition-Associated Cholestasis in Premature Neonates: 15 Years' Experience. *Nutr Clin Pract.* 2017;**32**(3):407-13. doi: [10.1177/0884533616687532](https://doi.org/10.1177/0884533616687532). [PubMed: [28135431](https://pubmed.ncbi.nlm.nih.gov/28135431/)].
35. Repa A, Lochmann R, Unterasinger L, Weber M, Berger A, Haiden N. Aggressive nutrition in extremely low birth weight infants: impact on parenteral nutrition associated cholestasis and growth. *PeerJ.* 2016;**4**:e2483. doi: [10.7717/peerj.2483](https://doi.org/10.7717/peerj.2483). [PubMed: [27688976](https://pubmed.ncbi.nlm.nih.gov/27688976/)]. [PubMed Central: [PMC5036079](https://pubmed.ncbi.nlm.nih.gov/PMC5036079/)].
36. Ehrenkranz RA. Early, aggressive nutritional management for very low birth weight infants: what is the evidence? *Semin Perinatol.* 2007;**31**(2):48-55. doi: [10.1053/j.semperi.2007.02.001](https://doi.org/10.1053/j.semperi.2007.02.001). [PubMed: [17462488](https://pubmed.ncbi.nlm.nih.gov/17462488/)].
37. Ehrenkranz RA. Ongoing issues in the intensive care for the periviable infant-nutritional management and prevention of bronchopulmonary dysplasia and nosocomial infections. *Semin Perinatol.* 2014;**38**(1):25-30. doi: [10.1053/j.semperi.2013.07.005](https://doi.org/10.1053/j.semperi.2013.07.005). [PubMed: [24468566](https://pubmed.ncbi.nlm.nih.gov/24468566/)].
38. Hamilton E, Massey C, Ross J, Taylor S. Early enteral feeding in very low birth weight infants. *Early Hum Dev.* 2014;**90**(5):227-30. doi: [10.1016/j.earlhumdev.2014.02.005](https://doi.org/10.1016/j.earlhumdev.2014.02.005). [PubMed: [24612934](https://pubmed.ncbi.nlm.nih.gov/24612934/)].