Preparation and quality control of ¹⁵³Sm-[*tris*(1,10 -phenanthroline) samarium (III)] complex

Z. Naseri¹, A. Hakimi², S. Shirvani-Arani³, A.R. Jalilian^{3*}, A. Bahrami-Samani³, M. Nemati Kharat¹, M. Ghannadi-Maragheh³

¹School of chemistry, University College of Science, University of Tehran, Iran

²Health Physics and Dosimetry Research Laboratory, Department of Physics, Amir kabir University of Technology, Tehran, Iran

³Radiopharmaceutical Research and Development Lab (RRDL), Nuclear Science and Technology Research Institute (NSTRI), Tehran, Iran

Background: The ¹⁵³Sm-[tris(1,10-phenanthroline)] Samarium(III)]complex (153Sm-PL₃) was prepared in view of development of targeting therapeutic compounds for malignancies, and interesting in-vitro anti-tumor activities of lanthanide phenanthroline complexes,. Materials and Methods: Sm-153 chloride was obtained by thermal neutron flux $(4 \times 10^{13} \text{ n.cm}^{-1})$ ².s⁻¹) of enriched ¹⁵²Sm₂O₃ sample, dissolved in acidic media. The labeling was performed in ethanol in 24h, controlled by ITLC (1.0mM DTPA, pH.5, as mobile phase). The partition coefficient for the labeled compound was also determined. Results: A radiochemical yield of more than 95% was obtained. Radiochemical purity of 96% was obtained using ITLC with specific activity of about 27.75 GBg/mg. The radio-labeled complex was stable in aqueous solution at least 24 hours and no significant amount of free ¹⁵³Sm was released from the complex. The partition coefficient for the labeled compound was determined (log P. 3.4). The complex was stable in final formulation for 66h. The biological evaluation of the compound is under investigation. Conclusion: The radiolabeled compound used in this study was a very inexpensive and useful agent for the use as a therapeutic compound. Iran. J. Radiat. Res., 2012; 10(1): 59-62

Keywords: Samarium-153, 1, 10-phenanthroline, radiolabeling, quality control.

INTRODUCTION

[Tris(1,10-phenanthroline)lanthanum (III)] (La-PL₃) has been prepared previously and the rigid planar 1,10-phenanthroline (PL) molecule demonstrated distinct effects on *in vitro* cultured cells. The complex also has shown to stop DNA synthesis in CCRF-CEM and Ehrlich ascites cells leading to a cell cycle arrest in $GO/G1^{(1, 2)}$ based on the metal chelating ability of 1,10-phen ^(3, 4), several metal ions including copper, ruthenium and cobalt has shown to enhance the anticancer activity of PL ⁽⁵⁾.

On the other hand, several complexes of vanadium with PL derivatives has shown to demonstrate apoptotic effect *in vivo* and *in vitro* ⁽⁶⁻⁸⁾. Recently, La-PL₃ demonstrated anticancer activity via potent induction of cell cycle arrest and/or apoptosis with promising in vivo anticancer activity against a human colon cancer xenograft, suggesting, La-PL₃ as a new anticancer metal-drug ⁽⁹⁾.

Radioisotopes with medium-energy beta emissions and half-life of a few days are attractive candidates for systemic delivery of targeted irradiation ⁽¹⁰⁾, such as ¹⁵³Sm (T_{1/2} = 46.7 h), also having medium-energy gamma photon (103 keV) which is suitable for imaging.

In this research, ¹⁵³Sm-PL₃ (figure 1) complex was prepared and the effects of various production conditions were investigated on its labeling yield.

*Corresponding author:

Dr. Amir Reza Jalilian,

Radiopharmaceutical Research and Development Lab (RRDL), Nuclear Science and Technology Research Institute (NSTRI), Tehran, Iran, Postal code: 14155-1339.

E-mail: ajalili@aeoi.org.ir



Figure 1. Possible chemical formula for ¹⁵³Sm-PL₃.

MATERIALS AND METHODS

Production of ¹⁵³Sm was performed at Tehran Research Reactor (TRR) using ¹⁵²Sm (n, g) 153 Sm reaction with 152 Sm in purity of 98.7% (ISOTEC Inc.) 1,10-phenanthroline was purchased from Aldrich Co., Germany, without further purification. Chromatography paper (Whatman No. 2) was obtained from Whatman (Maidstone, UK). Radiochromatography was performed using a bioscan AR-2000 radio TLC scanner instrument (Bioscan, Paris, France). A high purity germanium (HPGe) detector, coupled with a Canberra[™] (model GC1020-7500SL) multichannel analyzer and a dose calibrator ISOMED 1010 (Dresden, Germany), were used for counting distributed activity in rat organs. All other chemical reagents were purchased from Merck (Darmstadt, Germany). Calculations were based on the 103 keV peak for ¹⁵³Sm. All values were expressed as mean ± standard deviation (Mean± SD), and the data were compared using Student's T-test. Statistical significance was defined as P<0.05.

Synthesis of ¹⁵³Sm-PL₃ complex

The ¹⁵³Sm was produced by neutron irradiation of 100 μ g of enriched ¹⁵²Sm₂O₃ according to reported procedures ⁽¹¹⁾ at a thermal neutron flux of 4×10¹³ n.cm⁻².s⁻¹ for 5 days. Specific activity of the ¹⁵³Sm was 27.75 GBq/mg. The irradiated target was dissolved in 200 μ l of 1.0 mol/L HCl, to prepare ¹⁵³SmCl₃ and diluted to the appropriate volume with ultra pure water, to produce a stock solution. The mixture was filtered through a 0.22 μ m biological filter and sent for use in the radiolableing step. Radionuclidic purity of the solution was tested for the presence of other radionuclides using beta spectroscopy and HPGe spectroscopy to detect various interfering beta and gamma emitting radionuclides. The radiochemical purity was also checked by Whatman No.2 chromatography paper, and developed in a mixture of 1.0 mmol/L DTPA solution as mobile phase.

The acidic solution (0.2 ml) of ¹⁵³SmCl₃ (111 MBq, 3 mCi) was transferred to a 5 mlborosilicate vial and heated to dryness using a flow of N₂ gas at 50-60°C. Two hundred microlitres of 1,10-phenanthroline monohydrate (PL) in absolute ethanol (5 mg/ml) was added to the activity-containing vial and the mixture was diluted by the addition of normal saline (300 µl) followed by vortexing at 60°C for 30-60 min. The active solution was checked for radiochemical purity by ITLC. The final solution was then passed through a 0.22 mm filter and pH was adjusted to 5.5-7. From the final product, 5 µl was applied to a Whatman No.2 strip and followed by developing in 1mM DTPA (pH.5). Radioactivity was determined by a RTLC scanner.

Stability of ¹⁵³Sm-PL₃ in final formulation

Stability tests were based on previous studies performed for radiolabeled metal complexes $^{(12)}$. A sample of 153 Sm-PL₃ (100 MBq) in aqueous solution was kept at room temperature for 6 hours while checked by RTLC. Micro-samples (5 µl) taken from the shaken mixture were transferred the TLC papers and the ratio of 153 Sm cation to 153 Sm -PL₃ were checked (eluent: 1mM DTPA).

Determination of partition coefficient

Partition coefficient (log P) of ¹⁵³Sm-PL₃ complex was calculated followed by the determination of P (P= the ration of specific activities of the organic and aqueous phases) ⁽¹³⁾. A mixture of 1 ml of 2-octanol and 1 ml of radiolabeled samarium complex at 37°C was vortexes for 2 hours and left for 30 minutes in room temperature. Then the octanol and aqueous phases were sampled (5 μ l) and counted in HPGe detector for 1000 seconds.

RESULTS

The radionuclide was prepared in a research reactor according to regular methods with a range of specific activity 600 -750 mCi/mg for radiolabeling use, after counting the samples on an HPGe detector for 5 hours, very slight amount of impurities were recorded and shown to be Eu radionuclides as shown in table 1 (figure 2).

Table 1. The radionuclidic impurities and their percentages in
the final Sm-153 samples produced form enriched Sm-152
(n=5).

Radionuclides	Impurity (%)	
Eu-154	< 2.27e-4	
Eu-155	< 1.02e-4	
Eu-156	< 4.90e-4	

The radioisotope was dissolved in acidic media as a starting sample and was further diluted and evaporated for obtaining the desired pH and volume followed by sterile filtering.

To determine stability of complex in aqueous solution, a sample of ¹⁵³Smphenantroline in aqueous solution was kept at room temperature for 6 hours while checked by RTLC. Micro-samples (5 µl) taken from the shaken mixture were transferred to the TLC papers, and the ratio of free radio Samarium to ¹⁵³Sm-Phenantroline were checked (eluent: 1mM DTPA) (figure 3).

As expected, the lipophilicity of the compound is rather high. The measured water/octanol partition coefficient, P, for the ¹⁵³Sm-complex found to be 3.4 at pH,7.

DISCUSSION

The radiochemical purity of the 153 Sm solution was checked in 1mM DTPA, and free Sm³⁺ cation is complexed to more



Figure 2. gamma spectrum for Sm-153 prepared by neutron irradiation of Sm-152 sample using an HPGe detector.



Figure 3. RTLC chromatograms of ¹⁵³Sm-SmCl₃ solution (left) and final ¹⁵³Sm-PL₃ solution (right) using 1mMDTPA solution.

Iran. J. Radiat. Res., Vol. 10, No. 1, June 2012

lipophilic SmDTPA form, and migrated to higher R_f, while small radioactive fraction remained at the origin, which could be related to other Lu ionic species, not forming SmDTPA complex, such as SmCl₄, etc. and/or colloids. While in case of ¹⁵³Sm-PL₃, the complex slowly migrated to Rf. 0.1, which was easily distinguishable from the free cation. The complexation ability of the eluting DTPA should not exceed the stability of the complex in the media since the complex can be destroyed in high molar ratios of DTPA, this phenomenon had been already reported and applied in the determination of various radionuclidic samples ⁽¹⁴⁾.

The free ¹⁵³Sm/ ¹⁵³Sm-phenantroline ratio in the labeled sample remained unchanged 4:96. The complex had significant stability in the aqueous media which made it possible to formulate the compound for upcoming biological studies in experimental animals. Stability of such complexes has also been reported while the stability in presence of human serum was also enough (>97% in 10 hours), and appropriate for future works. The lanthanide-phenanthroline complexes had logK of higher than $15^{(6-8)}$.

Regarding the high lipophilicity ratio, the high lipophilicity of the complex would be helpful since the complex could easily penetrate through blood-brain barrier (BBB) as well as other biological membranes and has allowed rapid uptake though cells, as well as lipophilic nucleus membrane (4, 5).

CONCLUSION

62

The method used in this research for the production of ¹⁵³Sm-PL₃ was quite simple and cost effective. The radiochemical purity was 96% and the labeling and quality control took 24 hours. Trace amounts of ¹⁵³SmCl₃ (~4%) were detected by ITLC which showed that radiochemical purity of the ¹⁵³Sm-PL₃ was higher than 96%. (specific activity, 4.20 TBq/mmol). The radio-labeled complex was stable in aqueous solution at least 6 hours and no significant amount of free ¹⁵³Sm was released from the complex.

Our experiments on this compound have shown satisfactory quality, and stability for future therapeutic studies.

REFERENCES

- 1. Krishnamurti C, Saryan LA, Petering DH (1980) Effects of ethylenediaminetetraacetic acid and 1, 10phenanthroline on cell proliferation and DNA synthesis of Ehrlich ascites cells. Cancer Res. 40: 4092-9.
- 2. Falchuk KH and Krishan A (1977) 1, 10-Phenanthroline inhibition of lymphoblast cell cycle. Cancer Res. 37: 2050-6.
- 3. McFadyen WD, Wakelin LP, Roos IA, Leopold VA (1985) Activity of platinum(II) intercalating agents against murine leukemia L1210. J Med Chem, 28: 1113-6.
- Sammes PG and Yahioglu G (1994) 1,10-Phenanthroline: a versatile ligand. *Chem Soc Rev,* 23: 4. 327-34
- Wang ZM, Lin HK, Zhu SR, Liu TF, Zhou ZF, Chen YT 5. (2000) Synthesis, characterization and cytotoxicity of lanthanum(III) complexes with novel 1,10-phenanthroline- 2,9-bis-alpha-amino acid conjugates. Anticancer Drug Des, 15: 405–11.
- 6. Muggia FM and Fojo T (2004) Platinums: extending their therapeutic spectrum. J Chemother, 16: 77-82.
- Green DR (2005) Apoptotic pathways: ten minutes to dead. Cell, 121: 671-4.
- 8. Szakacs G, Annereau JP, Lababidi S, Shankavaram U, Arciello A, Bussey KJ et al. (2004) Predicting drug sensitivity and resistance: profiling ABC transporter genes in cancer cells. Cancer Cell, 6:129-37.
- 9. Heffeter P, Jakupec MA, Ko"rner W, Wild S, Graf von Keyserlingk N, Elbling L, Zorbas H, Korynevska A, Knasmu"ller S, Sutterluty S, Micksche M, Keppler BK, Berger W (2006) Anticancer activity of the lanthanum compound [tris(1,10-phenanthroline)lanthanum(III)] trithiocyanate (KP772; FFC24). Biochemical Pharmacology, 71: 426-440.
- 10. Anderson PM, Gregory A. Wiseman GA, Angela Dispenzieri A et al. (2002) High-Dose Samarium-153 Ethylene Diamine Tetramethylene Phosphonate: Low Toxicity of Skeletal Irradiation in Patiens with Osteosarcoma and Bone Metastases. J Cilin Oncology, 20: 189-196.
- 11. Ferro-Flores G, De Mara Ramrez F, Tendilla JI, Pimentel -Gonzlez G, Murphy CA, Melndez-Alafort L, Ascencio JA, Croft BY (1999) Preparation and Pharmacokinetics of Samarium(III)-153-Labeled DTPA-bis-Biotin. Characterization and Theoretical Studies of the Samarium(III)-152, Conjugate. Bioconjugate Chem, 10: 726.
- 12. Lewis MR, Wang M, Axworthy DB, Theodore LJ, Mallet RW, Fritzberg AR, Welch MJ, Anderson CJ (2003) In-vivo evaluation of pretargeted 64Cu for tumor imaging and therapy. J Nucl Med. 44: 1284-1292.
- 13. Jalilian AR, Amir Hakimi A, Garousi J, Bolourinovin F, Kamali-Dehghan M, Aslani G (2008) Development of [201TI](III) oxinate complex for in vitro cell labeling. Iran J Radiat Res, 6: 145-150.
- 14. Akhlaghi M, Kamalidehghan M, Jalilian AR, Shadanpoor N (2208) Determination of [201TI]TI(III) in [201TI]TICI solutions using HPLC. Appl Radiat Isot, 66: 479-81.