

# Exposure to radiofrequency radiation emitted from mobile phone jammers adversely affects the quality of human sperm

M.E. Parsanezhad<sup>1</sup>, S.M.J. Mortazavi<sup>2,3\*</sup>, T. Doohandeh<sup>1</sup>,  
B. Namavar Jahromi<sup>1</sup>, H. Mozdarani<sup>4</sup>, A. Zarei<sup>5</sup>, M. Davari<sup>5</sup>, S. Amjadi<sup>5</sup>,  
A. Soleimani<sup>6</sup>, M. Haghani<sup>2</sup>

<sup>1</sup> Department of Obstetrics and Gynecology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

<sup>2</sup> Ionizing and Non-ionizing Radiation Protection Research Center (INIRPRC), Shiraz University of Medical Sciences, Shiraz, Iran

<sup>3</sup> Department of Medical Physics and Medical Engineering, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

<sup>4</sup> Department of Medical Genetics, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

<sup>5</sup> IVF center, Shiraz University of Medical Sciences, Shiraz, Iran

<sup>6</sup> Department of Epidemiology, School of Health, Shiraz University of Medical Sciences, Shiraz, Iran

## ABSTRACT

**Background:** The health effect of rapidly increasing everyday exposure of humans to radiofrequency radiation is a major global concern. Mobile phone jammers prevent the mobile phones from receiving signals from base stations by interfering with authorized mobile carriers' services. In spite of the fact that mobile jammer use is illegal, they are occasionally used in offices, shrines, conference rooms and cinemas. *The purpose of this study was to investigate the biological effects of short term exposure of human sperm to radiofrequency radiation emitted from a commercial mobile phone jammer.*

**Materials and Methods:** Fresh semen samples were obtained by masturbation from 50 healthy donors who had referred with their wives to Infertility Treatment Center at the Mother and Child Hospital, Shiraz University of Medical Sciences. Female problem was diagnosed as the reason for infertility in these couples. The semen sample of each participant was divided into 4 aliquots. The first aliquot was subjected to *swim-up* and exposed to jammer radiation. The second aliquot was not subjected to *swim-up* but was exposed to jammer radiation. The third and fourth aliquots were not exposed to jammer radiation but only the 3<sup>rd</sup> aliquot was subjected to *swim-up*. **Results:** Semen samples exposed to radiofrequency radiation showed a significant decrease in sperm motility and increase in DNA fragmentation. **Conclusion:** Electromagnetic radiation in radiofrequency range emitted from mobile phone jammers may lead to decreased motility and increased DNA fragmentation in human semen. It can be concluded that mobile phone jamming might exert adverse reproductive health effects.

**Keywords:** Mobile phone jammers, sperm motility, DNA fragmentation, EMFs, microwave, RF, reproductive health.

## ► Original article

### \*Corresponding author:

Dr. SMJ Mortazavi,

Fax: +98 711 2349332

E-mail:

[mortazavismj@gmail.com](mailto:mortazavismj@gmail.com)

Revised: June 2015

Accepted: Feb. 2016

*Int. J. Radiat. Res.*, January 2017;  
15(1): 63-70

DOI: 10.18869/acadpub.ijrr.15.1.63

## INTRODUCTION

Infertility is a relatively common disorder<sup>(1)</sup> affecting 10-15% of American couples and 20%

of couples at reproductive age worldwide. Modern life has prompted humans to generate, transmit and use more electricity and now electricity is an essential component of the life in

developed and developing countries. The strong link between electricity and modern life has led to exponentially increasing exposure to different sources of EMFs. New findings show that male infertility that is associated with factors such as reduced sperm production and misshapen or immotile sperms may be linked with human exposure to different electromagnetic fields (EMFs). Substantial evidence now indicates that exposure to different sources of EMFs such as mobile phones <sup>(1-5)</sup>, mobile phone jammers <sup>(6)</sup>, laptops <sup>(7)</sup> or wireless internet-connected laptops <sup>(8)</sup> or extremely low frequency electromagnetic field (ELFs) <sup>(9)</sup> may decrease the quality of human sperm.

In 2009, Mailankot and colleagues reported the result of their study on sperm function in a rat model. Their study revealed that exposure to radiofrequency radiation at 0.9/1.8GHz results in decreased motility and induction of oxidative stress in exposed sperms <sup>(10)</sup>. Agarwal et al reported a human study on unprocessed ejaculated semen after exposure to cell phone electromagnetic radiation. They compared sperm motility, reactive oxidative stress (ROS), and DNA damage in specimen exposed to mobile radiations in talk mode and intact specimen. The results showed significant decrease in motility and viability, and increased ROS and DNA damage in the group of specimens exposed to mobile electromagnetic radiations <sup>(11)</sup>.

Mobile phone jammers are electronic devices that emit radiofrequency radiation with similar frequencies to that of mobile phones to block the access of the cell phone to the mobile base stations. These devices are used in places that mobile phone utilization is prohibited such as examination halls, and places where mobile phone ringing causes disturbance such as libraries. Most of mobile jammers propagate radiofrequency radiation at 800-1900MHz within a 10 meter diameter. Over the past years, our laboratories at INIRPRC have focused on studying the health effects of exposure of laboratory animals and humans to some common and/or occupational sources of electromagnetic fields such as mobile phones <sup>(12-20)</sup> and their base stations <sup>(21)</sup>, mobile phone jammers <sup>(22)</sup>, laptop computers <sup>(23)</sup>, radars <sup>(24)</sup>,

dentistry cavitrons <sup>(25)</sup> and MRI <sup>(26,27)</sup>. To the best of our knowledge there is limited number of investigations dealing with the bioeffects of exposure to microwave radiation emitted by mobile phone jammers on DNA fragmentation. The aim of present study was to investigate possible effects of radiofrequency radiation from mobile jammer devices on sperm motility and DNA fragmentation.

## MATERIALS AND METHODS

### Study population

Semen samples were obtained by masturbation from 50 healthy volunteers who had referred to Mother and Child hospital for evaluation of their infertility and female factor infertility was confirmed by the tests. All participants had no history of any medical disorder which could affect semen quality. Systemic autoimmune disorders, history of or evident cryptorchidism, UTI symptoms or history of urogenital infections, high radiofrequency occupational exposure or residential area, smoking, cystic fibrosis, history of vasectomy, history of major testicular trauma or operation, varicocele, hydrocele, testicular hypertrophy or atrophy in physical examination were exclusion criteria for sample collection.

All participants voluntarily participated in this study and informed consent was obtained from each individual. The project was approved by the ethics committee of the institutional review board for human medical research of Shiraz University of Medical Sciences.

### Experimental protocol

Semen samples were collected according to standard protocols for semen analysis after a period of sexual abstinence (2-7 days). A total number of 50 semen samples were selected after initial evaluation according to WHO guidelines <sup>(28)</sup> for semen analysis with specific attention to sperm motility and sperm count. Collected semen specimens were protected from extremes of temperature and delivered to the laboratory within 1-2 hours of collection. To avoid the problems caused by the time interval between

semen specimen collection and its delivery to the laboratory, control specimens were treated in a similar manner.

After liquefaction, semen samples from each donor were coded and divided into 2 aliquots labeled as A, B. In group A samples, swim-up technique was performed for elimination of seminal fluid and selecting the most active and motile sperms. B-labeled samples were used fresh. The swim-up technique uses sperm self-migration to obtain a sperm sample with a motility of at least 90%. In this technique, a layer of fresh media is added to the semen sample and the majority of the motile sperms will swim out of the sample and upward into the added media.

### **Exposure system**

A mobile jammer device (MB06-Mobile Blocker) which operates in four distinct frequency ranges including global system for mobile communications (GSM, 850 MHz, 900 MHz, 1800 MHz, 1900 MHz), digital cellular service (DCS), code division multiple access (CDMA), and the third-generation (3G) was used in this study. The maximum effective radius for operation of this device was reported by the manufacturer to range 10 to 40 meters. Group A and B samples from each donor were divided in 4 aliquots. Group 1, 2, and 3 were placed at distances of 1, 3, and 5 meters from the jammer device and the fourth group placed at a point at least 500 meters away from the jammer device. After 2 hours, a sample from each specimen was evaluated for motility and DNA fragmentation. Re-sampling performed after another 2 hours in all groups for same the mentioned characteristics.

### **Sperm quality assessment**

Motility of sperms was measured according to the well-defined four categories of motility:

1. Fast progressive motility: characterized by active movement, either linearly or in a large circle with a speed higher than 25  $\mu\text{m}/\text{sec}$
2. Slow progressive motility: characterized by active movement, either linearly or in a large circle with a speed less than 25  $\mu\text{m}/\text{sec}$

3. Non-progressive motility: regarding any type of motility without effective progression such as swimming in small circles, the flagellar force hardly displacing the head, or when only a flagellar beat can be observed.

4. Immotility: when no movement is seen by the sperms

Percentage of each type of mentioned categories were measured in each sampling and compared with other groups.

For measurement of DNA fragmentation, ultrathin smears of semen specimen were prepared and fixed using 30 minutes in the 1:1 solution of acetone-ethanol. After fixation, and drying in room temperature, staining with toluidine blue 0.05% in phosphate citrate buffer for 10 minute was performed. In toluidine blue staining, those nuclei with fragmented DNA and damaged nucleic content are stained dark blue or violet and those with intact DNA materials stained light blue color. By counting sperms with each nucleus color in multiple high power fields and reporting the percentage of each type, measurement of DNA fragmentation was performed.

Inspection and evaluation of semen specimen for motility and DNA fragmentation was performed at 100X magnification by, Leica DM2700 light microscope (Leica Microsystems, Wetzlar, Germany).

### **Statistical analysis method**

Data regarding motility and DNA fragmentation were recorded. Comparison of variables among studied groups at different distances (1, 3 and 5 meters) as well as different exposure times (2 and 4 hours) were performed using repeated measure analysis of variance. SPSS 18 software (IBM corporations, Armonk, New York, U.S) was used for data analysis. P-value < 0.05 was considered as significant.

## **RESULTS**

Demographic data of the men participated in this study are summarized in table 1. The mean ( $\pm$ SD) age of the participants was  $35.49 \pm 7.00$

*Int. J. Radiat. Res., Vol. 15 No. 1, January 2017*

years (ranged 24-50 years). Effect of distance from jammer device on sperm motility and DNA fragmentation in washed specimen after 2 hours is summarized in table 2. In the exposed sperm samples, 2 hours of exposure to electromagnetic fields at a distance of 1 meter from the Wi-Fi router produced a significant decrease in the proportion of fast progressive sperms ( $p < 0.001$ ). Furthermore, exposure to the Wi-Fi electromagnetic fields produced a significant increase in DNA fragmentation ( $p < 0.01$ ).

On the other hand, sperm samples exposed for 2 hours at a distance of 3 meters from the Wi-Fi router produced a significant decrease in the proportion of fast progressive sperms ( $p < 0.001$ ). However, exposure to the Wi-Fi electromagnetic fields could not produce a significant increase in DNA fragmentation ( $p = 1$ ). Furthermore, sperm samples exposed for 2 hours at a distance of 5 meters from the Wi-Fi router produced a significant decrease in the proportion of fast progressive sperms ( $p < 0.001$ ). However, exposure to the Wi-Fi electromagnetic fields could not produce a significant increase in DNA fragmentation

( $p = 0.29$ ).

Table 3 summarizes the effect of distance from jammer device on sperm motility and DNA fragmentation in washed specimen after 4 hours. In the exposed sperm samples, 4 hours of exposure to electromagnetic fields at a distance of 1 meter from the Wi-Fi router produced a significant decrease in the proportion of fast progressive sperms ( $p < 0.001$ ). Furthermore, exposure to the Wi-Fi electromagnetic fields produced a significant increase in DNA fragmentation ( $p < 0.001$ ). On the other hand, sperm samples exposed for 4 hours at a distance of 3 meters from the Wi-Fi router produced a significant decrease in the proportion of fast progressive sperms ( $p < 0.001$ ). However, exposure to the Wi-Fi electromagnetic fields could not produce a significant increase in DNA fragmentation ( $p = 0.095$ ). Furthermore, sperm samples exposed for 4 hours at a distance of 5 meters from the Wi-Fi router produced a significant decrease in the proportion of fast progressive sperms ( $p < 0.001$ ). However, exposure to the Wi-Fi electromagnetic fields produced a significant increase in DNA fragmentation ( $p = 0.04$ ).

Table 1. Characteristics of the men participated in this study.

Characters	Frequency (Percentage)
<b>Age at the Time of Exam</b>	
Mean ± SD (Range)	35.49 ± 7.00 (24 - 50)
<b>Distribution</b>	
24-30 years	33 (33.0%)
31-35 years	23 (23.0%)
36-40 years	22 (22.0%)
41-45 years	11 (11.0%)
≥46 years	11 (11.0%)
<b>Education</b>	
High School Diploma or Less	42 (42.0%)
College Degree	14 (14.0%)
B.Sc or Master	43 (43.0%)
G.P (Medical Doctor)	1 (1.0%)
<b>Job</b>	
Jobless	3 (3.0%)
Private Business	43 (43.0%)
Teacher	17 (17.0%)
Worker	6 (6.0%)
Government Employee	26 (26.0%)
University Lecturer	3 (3.0%)
Other Jobs	2 (2.0%)

**Table 2.** Effect of distance from jammer device on sperm motility and DNA fragmentation in washed specimen after 2 hours.

Group A	Fast progressive % (Mean± SD)	Slow progressive % (Mean± SD)	Non-progressive % (Mean± SD)	Immotile % (Mean± SD)	DNA fragmentation % (Mean± SD)
Washed sample control	24.2± 13.4	40.1±10.2	12.9±6.4	22.6±9.9	14.3±8.1
1 meter distance	13.0± 10.7	36.3±9.6	25.2±8.9	25.2±10.0	15.7±8.5
3 meters distance	14.6±10.9	36.0±11.2	24.3±7.8	25.0±8.5	14.5±9.3
5 meters distance	15.0± 10.2	38.0±9.7	22.6±6.7	24.1±9.5	14.9±8.1
P-value	0.0001	0.008	0.0001	0.0001	0.1

**Table 3.** Effect of distance from jammer device on sperm motility and DNA fragmentation in washed specimen after 4 hours.

Group A	Fast progressive % (Mean± SD)	Slow progressive % (Mean± SD)	Non-progressive % (Mean± SD)	Immotile % (Mean± SD)	DNA fragmentation % (Mean± SD)
Washed sample control	22.5±11.8	41.0±8.9	14.0±6.6	22.8±11.1	15.0±7.2
1 meter distance	12.2±9.2	35.5±9.8	27.7±8.3	24.8±10.5	15.9±7.9
3 meters distance	14.0±9.7	35.4±9.1	25.7±8.0	25.2±10.7	14.5±6.3
5 meters distance	14.7±9.7	36.5±9.9	23.4±6.7	24.4±11.1	14.6±8.1
P-value	<0.0001	<0.0001	<0.0001	0.002	0.15

Table 4 summarizes the effect of distance from jammer device on sperm motility and DNA fragmentation in fresh specimen after 2 hours. In the exposed sperm samples, 2 hours of exposure to electromagnetic fields at a distance of 1 meter from the Wi-Fi router produced a significant decrease in the proportion of fast progressive sperms ( $p < 0.001$ ). Furthermore, exposure to the Wi-Fi electromagnetic fields produced a significant increase in DNA fragmentation ( $p < 0.001$ ). On the other hand, sperm samples exposed for 2 hours at a distance of 3 meters from the Wi-Fi router produced a

significant decrease in the proportion of fast progressive sperms ( $p < 0.001$ ). However, exposure to the Wi-Fi electromagnetic fields could not produce a significant increase in DNA fragmentation ( $p = 0.095$ ). Furthermore, sperm samples exposed for 2 hours at a distance of 5 meters from the Wi-Fi router produced a significant decrease in the proportion of fast progressive sperms ( $p < 0.001$ ). However, exposure to the Wi-Fi electromagnetic fields produced a significant increase in DNA fragmentation ( $p = 0.04$ ).

**Table 4.** Effect of distance from jammer device of sperm motility and DNA fragmentation in fresh specimen after 2 hours.

Group B	Fast progressive % (Mean± SD)	Slow progressive % (Mean± SD)	Non-progressive % (Mean± SD)	Immotile % (Mean± SD)	DNA fragmentation % (Mean± SD)
Fresh sample control	51.1±18.5	34.4±13.8	10.6±6.5	4.0±4.4	13.2±9.3
1 meter distance	28.3±14.6	44.5±12.6	22.5±8.3	4.7±4.2	12.7±9.1
3 meters distance	31.6±15.1	40.6±12.4	22.3±9.5	5.4±4.5	13.2±9.9
5 meters distance	31.5±15.4	43.9±13.4	19.0±7.8	6.1±6.2	13.8±8.7
P-value	<0.0001	<0.0001	<0.0001	0.022	0.2

Table 5 summarizes the effect of distance from jammer device on sperm motility and DNA fragmentation in fresh specimen after 4 hours. In the exposed sperm samples, 4 hours of exposure to electromagnetic fields at a distance of 1 meter from the Wi-Fi router produced a

significant decrease in the proportion of fast progressive sperms ( $p < 0.001$ ). Furthermore, exposure to the Wi-Fi electromagnetic fields produced a significant increase in DNA fragmentation ( $p < 0.001$ ). On the other hand, sperm samples exposed for 4 hours at a distance

of 3 meters from the Wi-Fi router produced a significant decrease in the proportion of fast progressive sperms ( $p < 0.001$ ). However, exposure to the Wi-Fi electromagnetic fields could not produce a significant increase in DNA fragmentation ( $p = 0.095$ ). Furthermore, sperm samples exposed for 4 hours at a distance of 5

meters from the Wi-Fi router produced a significant decrease in the proportion of fast progressive sperms ( $p < 0.001$ ). However, exposure to the Wi-Fi electromagnetic fields produced a significant increase in DNA fragmentation ( $p = 0.04$ ).

Table 5. Effect of distance from jammer device of sperm motility and DNA fragmentation in fresh specimen after 4 hours.

Group B	Fast progressive % (Mean±SD)	Slow progressive % (Mean±SD)	Non-progressive % (Mean±SD)	Immotile % (Mean±SD)	DNA fragmentation % (Mean±SD)
Fresh sample control	46.7±18.9	38.3±15.5	11.6±6.3	3.6±2.7	12.4±8.1
1 meter distance	28.9±14.5	40.6±13.0	24.4±8.5	5.2±3.7	12.7±8.2
3 meters distance	29.0±14.6	43.5±12.0	22.6±9.7	5.2±6.3	12.6±8.6
5 meters distance	31.1±16.0	42.9±12.3	20.8±8.8	5.7±7.4	12.4±7.5
P-value	<0.0001	0.01	<0.0001	0.051	0.92

## DISCUSSION

In the present study, we aimed to investigate the effects of radiofrequency electromagnetic waves propagated from a mobile jammer device at frequency of 800-1900 Hz on 2 major characteristics of human semen; motility and DNA fragmentation. The results altogether strongly suggest a significant distance dependent influence of mobile jammer radiations on sperm motility. The highest influence was observed at a radius of 3 meters from the device. Regarding DNA fragmentation, significant difference was observed between exposed and sham-exposed specimens. Washed semen samples exposed to Wi-Fi radiation had significantly higher percentages of sperms with DNA damage. In fresh semen samples, DNA fragmentation was significantly higher in samples exposed for 4 hours compared to those of the controls. However, there was no difference between 2 hour exposed groups and the controls.

Generally speaking our results are in line with our previous findings. Mortazavi *et al.* for the first time reported that the exposure of rats to electromagnetic fields caused by laptop computers can decrease sperm count and motility which adversely affects male reproductive capabilities (7). Avendano *et al.* reported that human sperm samples exposed to

Wi-Fi internet-connected laptop for a short period of 4 hours exhibited a statistically significant decrease in progressive sperm motility and also an increase in sperm DNA fragmentation (8).

Numerous animal and human model studies have shown decreased sperm motility after exposure to radiofrequency radiation emitted from cell phones operating at 0.9/1.8GHz frequency. In a pilot human study performed in 2009, Agarwal *et al.* evaluated sperm motility and viability, ROS and DNA damage in fresh semen samples from 23 healthy donors and 7 infertile patients after 1 hour exposure to cell phone radiation in “talk” mode. The differences between exposed and control groups were significant for decreased motility and viability and increased ROS. Based on these observations, the authors linked impaired sperm characteristic to reactive oxygen stress caused by non-thermal damage due to radiofrequency electromagnetic radiation. On the other hand, for the DNA damage, there was no significant difference between radiation-exposed samples and controls, similar to the findings of this study (3). Same findings were reported by Eroglu *et al.* who exposed fresh human semen to 900 Hz radiofrequency electromagnetic radiation and evaluated the motility of sperms. The results showed a significant decrease in percentage of fast and slow progressive sperms and increased

percentages of immotile and non-progressive sperms<sup>(4)</sup>. Later in 2008, in a similar study by Falzone *et al.*, density-purified human sperm were exposed to 900 MHz cell phone radiation. The authors found no significant variation between exposed and control samples regarding sperm kinematic parameters. Also evaluation of mitochondrial membrane potential was performed and revealed no difference between exposed and control semen samples<sup>(9)</sup>. *In vitro* human epidemiologic studies on male individuals exposed to radiofrequency electromagnetic radiation have shown significant decrease in sperm count, motility and increased reactive oxidative stress<sup>(29)</sup>.

In the present study, results obtained by assessment of sperm motility and DNA fragmentation revealed significant decrease in sperm motility. Unfortunately due to limitations in funding, we were unable to perform reactive oxidative stress tests to investigate possible links between sperm motility impairments and ROS alterations. The main advantage in the protocol used in this study was performing the experiment using both fresh sperms and swim-up washed specimen. In this light, we were able to evaluate the radiofrequency effect on both fresh samples and motility gated sperms which revealed decreased motility in both samples.

For further investigations on this topic, we suggest performing this study with evaluation of ROS and total antioxidant capacity (TAC) in samples to shed light on cellular mechanisms which caused electromagnetic radiation induced decreased sperm motility in human sperms.

## CONCLUSION

Based on the findings of this study, exposure of human sperms to radiofrequency radiation emitted by a mobile phone jammer adversely affects the sperm quality. Semen samples exposed to jammer radiation showed a significant decrease in their motility and an increase in DNA fragmentation. The proportion of rapid progressive sperms in samples exposed to jammer radiofrequency radiation at different

distances was significantly lower than those of the sham-exposed samples. On the other hand, a significant increase was observed in DNA fragmentation in sperms exposed to jammer radiofrequency radiation compared to those of sham-exposed samples.

## ACKNOWLEDGEMENT

*This study was supported by the SUMS IVF center and Ionizing and Non-ionizing Radiation Protection Research Center (INIRPRC), Shiraz University of Medical Sciences (SUMS), Shiraz, Iran.*

**Conflict of interest:** Declared none.

## REFERENCES

1. De Luliis GN, Newey RJ, King BV, Aitken RJ (2009) Mobile phone radiation induces reactive oxygen species production and DNA damage in human spermatozoa *in-vitro*. *PLoS one*, **4(7)**:e6446.
2. Agarwal A, Deepinder F, Sharma RK, Ranga G, Li J (2008) Effect of cell phone usage on semen analysis in men attending infertility clinic: an observational study. (Translated from eng) *Fertility and sterility*, **89(1)**:124-128 (*in eng*).
3. Agarwal A, Desai NR, Makker K, Varghese A, Mouradi R, Sabanegh E, *et al.* (2009) Effects of radiofrequency electromagnetic waves (RF-EMW) from cellular phones on human ejaculated semen: an *in vitro* pilot study. *Fertility and sterility*, **92(4)**:1318-1325.
4. Eroglu O, Oztas E, Yildirim I, Kir T, Aydur E, Komesli G, *et al.* (2006) Effects of electromagnetic radiation from a cellular phone on human sperm motility: an *in-vitro* study. *Archives of Medical Research*, **37(7)**: 840-843.
5. Falzone N, Huyser C, Becker P, Leszczynski D, Franken DR (2011) The effect of pulsed 900-MHz GSM mobile phone radiation on the acrosome reaction, head morphometry and zona binding of human spermatozoa. *International Journal of Andrology*, **34(1)**: 20-26.
6. Mortazavi S, Parsanezhad M, Kazempour M, Ghahramani P, Mortazavi A, Davari M (2013) Male reproductive health under threat: Short term exposure to radiofrequency radiations emitted by common mobile jammers. *J Hum Reprod Sci*, **6(2)**: 124-128 (*in eng*).
7. Mortazavi SMJ, Tavassoli A, Ranjbari F, Moammaiee P (2010) Effects of Laptop Computers' Electromagnetic Field on Sperm Quality. *J Reprod Infertil*, **11(4)**: 251-258.
8. Avendano C, Mata A, Sanchez Sarmiento CA, Doncel GF

*Int. J. Radiat. Res., Vol. 15 No. 1, January 2017*

- (2012) Use of laptop computers connected to internet through Wi-Fi decreases human sperm motility and increases sperm DNA fragmentation. *Fertil Steril*, **97(1)**: 39-45 e32.
9. Falzone N, Huyser C, Fourie F, Toivo T, Leszczynski D, Franken D (2008) In vitro effect of pulsed 900 MHz GSM radiation on mitochondrial membrane potential and motility of human spermatozoa. *Bioelectromagnetics*, **29(4)**: 268-276.
  10. Mailankot M, Kunnath AP, Jayalekshmi H, Koduru B, Valsalan R (2009) Radio frequency electromagnetic radiation (RF-EMR) from GSM (0.9/1.8GHz) mobile phones induces oxidative stress and reduces sperm motility in rats. *Clinics*, **64(6)**: 561-565.
  11. Agarwal A, Desai NR, Makker K, Varghese A, Mouradi R, Sabanegh E, et al. (2009) Effects of radiofrequency electromagnetic waves (RF-EMW) from cellular phones on human ejaculated semen: an in vitro pilot study. *Fertility and Sterility*, **92(4)**: 1318-1325.
  12. Mortazavi SMJ, Motamedifar M, Namdari G, Taheri M, Mortazavi AR, Shokrpour N. (2014) Non-Linear Adaptive Phenomena which Decrease the Risk of infection after Pre-Exposure to Radiofrequency Radiation. *Dose-Response*, **12(2)**: 233-245.
  13. Mortazavi SMJ, Rouintan MS, Taeb S, Dehghan N, Ghaffarpanah AA, Sadeghi Z, et al. (2012) Human short-term exposure to electromagnetic fields emitted by mobile phones decreases computer-assisted visual reaction time. *Acta Neurologica Belgica*, **112(2)**: 171-175.
  14. Mortazavi SMJ, Mosleh-Shirazi MA, Tavassoli AR, Taheri M, Mehdizadeh AR, Namazi SAS, et al. (2013) Increased Radioresistance to Lethal Doses of Gamma Rays in Mice and Rats after Exposure to Microwave Radiation Emitted by a GSM Mobile Phone Simulator. *Dose-response : a publication of International Hormesis Society*, **11(2)**: 281-292.
  15. Mortazavi S, Mosleh-Shirazi M, Tavassoli A, Taheri M, Bagheri Z, Ghalandari R, et al. (2011) A comparative study on the increased radioresistance to lethal doses of gamma rays after exposure to microwave radiation and oral intake of flaxseed oil. *Int. JRR*, **9(1)**: 9-14.
  16. Mortazavi SMJ, Habib A, Ganj-Karimi AH, Samimi-Doost R, Pour-Abedi A, Babaie A. (2009) Alterations in TSH and Thyroid Hormones Following Mobile Phone Use. *OMJ* **24**: 274-278
  17. Mortazavi SMJ, Daiee E, Yazdi A, Khiabani K, Kavousi A, Vazirinejad R, et al. (2008) Mercury release from dental amalgam restorations after magnetic resonance imaging and following mobile phone use. *Pakistan Journal of Biological Sciences*, **11(8)**: 1142-1146.
  18. Mortazavi SMJ, Ahmadi J, Shariati M (2007) Prevalence of subjective poor health symptoms associated with exposure to electromagnetic fields among University students. *Bioelectromagnetics*, **28(4)**: 326-330.
  19. Mortazavi SMJ, Motamedifar M, Namdari G, Taheri M, Mortazavi AR (2013) Counterbalancing immunosuppression-induced infections during long-term stay of humans in space. *Journal of Medical Hypotheses and Ideas*, **7(1)**: 8-10.
  20. Mortazavi SMJ, Atefi M, Kholghi F (2011) The pattern of mobile phone use and prevalence of self-reported symptoms in elementary and junior high school students in Shiraz, Iran. *Iranian Journal of Medical Sciences*, **36(2)**: 96-103.
  21. Mortazavi SMJ (2013) Safety Issue of Mobile Phone Base Stations *Journal of Biomedical Physics & Engineering*, **3(1)**: 1-2.
  22. Mortazavi SMJ (2004) Adaptive responses after exposure to cosmic and natural terrestrial radiation. *Indian Journal of Radiation Research*, **104**-112.
  23. Mortazavi SMJ, Tavassoli AR, Ranjbari F, Moamaei P (2011) Effects of Laptop Computers' Electromagnetic Field on Sperm Quality. *Journal of Reproduction and Infertility* **11(4)**: 251-258.
  24. Mortazavi SMJ, Taeb S, Dehghan N (2013) Alterations of Visual Reaction Time and Short Term Memory in Military Radar Personnel. *Iranian J Publ Health*, **42(4)**: 428-435.
  25. Mortazavi SM, Vazife-Doost S, Yaghoobi M, Mehdizadeh S, Rajaie-Far A (2012) Occupational exposure of dentists to electromagnetic fields produced by magnetostrictive cavitrons alters the serum cortisol level. *Journal of Natural Science, Biology, and Medicine*, **3(1)**: 60-64.
  26. FCC (2011) Document: Enforcement bureau steps up education and enforcement efforts against cellphone and GPS jamming. Targeted education and outreach coupled with strict enforcement: *Federal Communications Commission*.
  27. Mortazavi SMJ, Neghab M, Anoshe SMH, Bahaeddini N, Mortazavi G, Neghab P (2014) High-field MRI and Mercury release from dental amalgam fillings. *THE IJOEM*, **5(2)**: 101-105.
  28. Organization WH (2010) *WHO laboratory manual for the examination and processing of human semen*. 5th ed. Geneva: World Health Organization. 2010 (WHO, Geneva).
  29. La Vignera S, Condorelli RA, Vicari E, D'Agata R, Calogero AE (2012) Effects of the exposure to mobile phones on male reproduction: a review of the literature. *Journal of Andrology*, **33(3)**: 350-356.