

Effects of Black Scorpion *Androctonus crasicauda* Venom on Striated Muscle Preparation *in vitro*.

Hossein Vatanpour*

Department of Pharmacology and Toxicology, School of Pharmacy, Shaheed Beheshti University of Medical Sciences.

Abstract

Effects of venom from black scorpion *Androctonus crasicauda* (AC) were determined on isolated chick biventer cervicis nerve-muscle and mouse hemidiaphragm preparations using twitch tension method. The isolated nerves were stimulated by electrical stimulator and response to each stimulus was recorded. The venom mainly acted prejunctionally to facilitate neuromuscular activity due to an increase in acetylcholine release.

Keywords: Scorpion; Muscle; Neuromuscular; Transmission; Acetylcholine.

Introduction

Over 1500 species of scorpions have been described. They are commonly subdivided into two groups: Buthidae (40% of known species) and Chactoids. All human hazardous scorpions belong to the Buthidae (Buthoid) family. Envenomation by scorpion stings is common in tropical and subtropical regions (1). Many investigators have shown that mammalian scorpion toxins are low molecular weight proteins of about 7000 daltons, as well as a single polypeptide chain cross-linked by four disulfide bridges (2, 3) which are toxic to mammals (4). The toxin can cause a variety of symptoms from pain at the site of sting to death (5). The venom of *Androctonus crasicauda* is a potent autonomic stimulator. Severity of symptoms depends on the size of the victim, season, and time lapse between sting and hospitalization. Vomiting, profuse sweating, priapism, mild pain at the sting site, local urticaria, and cool extremities are early signs of autonomic stimulation due to scorpion sting. Fatality after scorpion envenoming may be the result of cardiovascular failure complicated by

pulmonary oedema, as well as respiratory arrest (6). It has been proposed that the majority of the symptoms of envenoming are due to massive discharge of catecholamines during so-called autonomic storms (7). The underlying mechanism of transmitter release and the identity of the bioactive molecules in *Androctonus crasicauda* venom remain unknown. However, other scorpion venoms contain toxins that affect ion channels and synaptic transmission (8, 9). Scorpion venom toxicity to human has mainly been attributed to the pharmacological properties of toxic polypeptides active in mammals. Toxins that prolong the opening of Na⁺ channels by slowing the inactivation process (the so called a-type scorpion toxins) lead to a prolongation of the action potential. This effect, which is dependent on membrane potential, has been observed with the toxins mainly isolated from old world scorpion venoms, including *Androctonus australis* Hector, *Leiurus quinquestriatus quinquestriatus*, *Buthus tamulus* and *Buthus epeus* (10, 11). In contrast, there are toxins that cause Na⁺ channel to open at membrane potentials at which they would normally be closed (the b-type scorpion toxins) and provoke spontaneous and repetitive firing of action potentials. This effect has been

* Corresponding author:
E-mail: vatanpour@hotmail.com

observed with venom and toxins from new world scorpion, such as *Centruroides suffusus*, *Centruroides sculpturatus* and *Tityus serrulatus* (12, 13), although in two of these venoms (i.e. *Centruroides sculpturatus* and *Tityus serrulatus*) toxins representative of the two classes have been identified (13). These toxins can alter neuronal action potentials and cause release of neurotransmitter from cholinergic and adrenergic neurones. A few venoms have, in addition, minor components, e.g. iberiotoxin from *Buthus tamulus* venom (14), that act selectively on voltage-gated or Ca^{2+} -activated K^+ channels (15-19).

In vitro models of cholinergic transmission (chick biventer cervicis, mouse hemidiaphragm nerve-muscle preparations) were used in order to study changes in twitch tension and in acetylcholine release. Almost 30 years ago, Del Pozo while working with Mexican scorpion, has reported the action of the venom from the genus *Centruroides* on peripheral nervous system and muscle preparations (20). The venom of the same species applied at the frog neuromuscular junction can induce an increase in the duration of the presynaptic nerve action potential as well as an increase in the mean number of quanta released by a nerve impulse (21). Vital has also shown that the venom of the South American *Tityus serrulatus* induces the release of an acetylcholine-like substance, which could account for the twitches observed and also for the decurarizing activity of scorpion venom (22). It has been suggested that this scorpion toxin has two sites of action presynaptically: it depolarizes the nerve terminal facilitating the spontaneous release of transmitter (acetylcholine), and also acts on the membrane of the unmyelinated nerve terminal arborization prolonging the sodium current. It has been shown that scorpion *Centruroides noxius* Hoffman venom has a direct effect on K^+ channels. It reversibly blocks the delayed rectifier potassium current in the squid giant axon. Purified noxiustoxin blocks K^+ currents in the squid giant axon in a complex manner (23). At concentrations less than $1.5 \mu\text{M}$, block by noxiustoxin is independent of membrane potential, while at higher concentrations toxin block is voltage-dependent, and the toxin probably dissociates when the preparation is repetitively depolarized. These effects are

similar to the voltage-dependent properties of some Na^+ channel scorpion toxins.

The purpose of the present study is to investigate the effects of *Androctonus crasicauda* venom on the neuromuscular transmission. In vitro models of cholinergic transmission, including chick biventer cervicis and mouse hemidiaphragm nerve-muscle preparations, are used in order to study changes in muscle twitch tension, during the release of acetylcholine.

Experimental

Chick biventer cervicis nerve-muscle preparation and mouse phrenic nerve-hemidiaphragm preparation have been used for studying the effects of drugs and toxins on neuromuscular transmission. Both can be set up in small tissue baths (5 ml or less), and both are robust and stable over several hours in vitro. These features make them suitable for studying the effects of toxins, which are often available in only small quantities, and which are often slow in action.

Chick biventer cervicis preparations

The chick biventer cervicis muscle contain both focally innervated twitch fibres and multiply innervated contracture-producing fibres. Thus, it can be stimulated by exogenously applied cholinomimetic agonists, as well as by stimulation of its motor nerve. This enables prejunctional effects to be distinguished from postjunctional effects. Biventer cervicis nerve-muscle preparations (24) were isolated from 3-14 day old chicks that had been killed by exposure to anaesthetics (CO_2 for 2-7 days old and halotane for 7-12 days old chicks) and mounted (in pairs) with a resting tension of 0.5-1g in 2-10 ml tissue baths containing Krebs-Henseleit physiological salt solution of the following composition (mM): NaCl, 118.5; KCl, 4.7; MgSO_4 , 1.2; KH_2PO_4 , 1.2; CaCl_2 2.5; NaHCO_3 , 25.0; glucose, 11.1. The solution was maintained at 33°C and bubbled with 95% O_2 plus 5% CO_2 at pH 7.3.

For indirect stimulating, muscle contraction was induced by stimulation the motor nerve in the tendon, using ring electrodes every 10 sec with pulses of 0.2 ms duration and a voltage greater than that which produced a maximal twitch. In the absence of nerve stimulation, contractures to sub-maximal concentrations of

exogenously acetylcholine (1-3 mM), carbachol (20-40 μ M), and KCl (40-50 mM) were obtained prior to the addition of the test compound and at the end of the experiment. Acetylcholine and potassium chloride were allowed to bathe the preparation for 30 sec and carbachol for 60 sec. Preparations were then washed by overflow for 15 sec at a rate of 5-10 ml/s. Preparations were finally allowed to stabilize for 20-40 min before the addition of test compound.

For direct muscle stimulation, neuromuscular transmission was blocked with tubocurarine (10-20 μ M) and the stimulating electrodes were moved into contact with the belly of the muscle. Stimulation was performed directly at 0.1 Hz with pulses of 2 ms duration and a voltage greater than which was needed to induce maximum twitch height. Preparations were allowed to stabilize for approximately 30 min before the addition of test venoms or toxins.

Twitches and contractures were recorded isometrically on Narco-Trace polygraphs using F60 force displacement transducer.

Mouse hemidiaphragm nerve-muscle preparation

Mouse hemidiaphragms and attached phrenic nerve were isolated from 15-25 g male mice (Balb C strain) for the rat hemidiaphragm preparation. Preparations were mounted individually in 10 ml organ baths under the same conditions used for the biventer cervicis preparations, except that the temperature was maintained at 36.5°C.

For indirect stimulation, the phrenic nerve was stimulated (via a Burn and Rand electrode) at 0.1 Hz with pulses of 0.2 ms duration and voltage greater than that needed to produce maximal twitches.

For direct muscle stimulation, the preparation was impaled (just below the rib) upon two silver electrode. Pulses of 2 ms were applied at a frequency of 0.1 Hz with sufficient strength to produce a maximal twitch. Tubocurarine (10-20 μ M) was used to prevent neuromuscular transmission. Alternatively preparations can also be stimulated, indirectly via the phrenic nerve and directly via the hook electrodes at a frequency of 0.05 Hz. Under these conditions, tubocurarine was always added prior to the experiment to ensure that




Figure 1. Effects of *Androctonus crasicauda* venom on response of chick biventer cervicis preparation to direct (■) and indirect (○) stimulation. Each point represents the maximum response for that concentration (vertical bars are sem, n=4).

activation of nerve endings by the field stimulation do not contribute to the overall tension recorded in direct stimulation. In cases with more than a 10% decrease in response to direct stimulation in the presence of tubocurarine, the hook electrodes were repositioned to avoid this artifact. After washout of the tubocurarine, preparations was allowed 30-60 min to stabilize prior to the addition of toxin.

Results

Androctonus crasicauda venom (AC) was tested on directly and indirectly stimulated preparations at 0.3, 1, 3 and 10 μ g/ml




Figure 2. Effects of *Androctonus crasicauda* venom on responses of indirectly stimulated chick biventer cervicis nerve-muscle preparation (■) 0.3 μ g/ml, (□) 1 μ g/ml, (●) 3 μ g/ml and (○) 10 μ g/ml. Each point represents the mean \pm sem (n=4).




Figure 3. *Androctonus crasicauda* venom (10 $\mu\text{g/ml}$) induces a marked contracture followed by periodic oscillations of the resting tension.

(Figure 1). AC induced a large increase in the response of indirectly stimulated nerve-muscle preparation, but not directly stimulated preparation. The threshold concentration was around 1 $\mu\text{g/ml}$. Maximal twitch augmentation with indirect stimulation occurred after about 60 min and was $69 \pm 7\%$ and $300 \pm 40\%$ of control twitch height at 1 and 3 $\mu\text{g/ml}$, respectively (Figure 2). At 10 $\mu\text{g/ml}$, there was a transient augmentation ($34 \pm 22\%$) of twitch height followed by a contracture of $148 \pm 20\%$ of control twitch height. Thereafter, there was a series of large periodic (approximately one every 5 min) increase in resting tension, coinciding with the reduction of twitch height (Figure 3). The effect of venom was reversed by washing intermittently for about 45 min. AC (1-3 $\mu\text{g/ml}$) caused little change in post-junctional sensitivity, as assessed by contracture to exogenously applied acetylcholine, carbachol and KCl (Figure 4). In preparations without tubocurarine pretreatment, AC induced a $64 \pm 10\%$ increase in the contracture to exogenously applied KCl at 10 $\mu\text{g/ml}$, but no significant changes in contractures to acetylcholine and carbachol. However, contractures to KCl did not change when AC (10 $\mu\text{g/ml}$) was tested in preparations exposed to tubocurarine, indicating that the enhanced contracture to KCl may be due to an increase in the amount of acetylcholine released from nerve terminals by K^+ induced depolarization. In preparations stimulated




Figure 4. Effects of *Androctonus crasicauda* venom (1 $\mu\text{g/ml}$) on chick biventer cervicis preparation at 33°C . The muscle was stimulated indirectly once every ten seconds and, in the absence of twitches, by exogenous acetylcholine (1 mM), carbachol (40 μM) and potassium chloride (40 mM).

directly in the presence of tubocurarine, 10 $\mu\text{g/ml}$ AC caused a small increase in twitch responses ($29 \pm 13\%$) and a small contracture ($21 \pm 5\%$ of control twitch height). Additionally, in the absence of stimulation as well as in indirectly stimulated preparations, AC (10 $\mu\text{g/ml}$) caused contracture of around 15% of control twitch height. The contractures was abolished by higher concentrations of tubocurarine (10-100 μM), and was not seen in preparations pretreated with tubocurarine (Figure 5). In order to investigate whether the facilitatory effect was due to AC acting presynaptically or postsynaptically, AC (3 and 10 $\mu\text{g/ml}$) was tested on partially paralysed preparations (50-80%) with tubocurarine, MgCl_2 or procaine. In the presence of tubocurarine or Mg^{2+} , AC at 3 $\mu\text{g/ml}$ increased twitch height to about 120% greater than control after about 60 min; AC reversed the block induced by procaine by only 70%. At 10 $\mu\text{g/ml}$, AC reversed the reduction in twitch height induced by tubocurarine and procaine more rapidly, following by augmentation in the twitch height. The partial block induced by Mg^{2+} was incompletely reversed by AC (Figure 6).

AC at 3 and 10 $\mu\text{g/ml}$ increased twitch height of indirectly stimulated preparations by $53 \pm 11\%$ and $307 \pm 90\%$ of control twitch height, respectively. This increase did not cause contracture on mouse hemidiaphragm frenic nerve muscle preparations (Figure 7). Maximum augmentation was reached after 60 min. At 10 $\mu\text{g/ml}$, AC augmented responses of directly stimulated preparations by $34 \pm 9\%$ of the control twitch height.

Discussion

Envenoming by the Iranian scorpion *Androctonus crasicauda* can cause massive discharge of catecholamine. The relationship of

these effects to the clinical situation is unclear. Results from the present study show that venom from Iranian scorpion *Androctonus crasicauda* has some effects on neuromuscular transmission. Present results revealed some of the underlying mechanisms: AC venom can increase the release of acetylcholine at the neuromuscular junction by both increasing quantal content as well as the release after single shock stimulation. There is, however, no change in postsynaptic sensitivity. AC, at high concentration, causes very large periodic increases in resting tension on both indirectly stimulated and unstimulated chick biventer cervicis preparations. As these contractures are abolished by tubocurarine, they are probably caused by the release of acetylcholine induced by AC. Although toxins responsible for this action are still to be identified, but the initial steps toward their isolation have been undertaken in this study.

References

- (1) Goyffen M, Vachon M and Broglio N. Epidemiology and clinical characteristics of the scorpion envenomation in Tunisia. *Toxicon* (1982) 20: 337-44
- (2) Radmanesh M. Clinical study of hemiscorpion *Leptanus* in Iran. *J. trop. Med. Hyg.* (1990) 93: 327-332
- (3) El-Amin ED. Issues in management of scorpion sting in children. *Toxicon* (1992) 30: 111-115
- (4) Siemen D and Vogel W. Tetrodotoxin interferes with the reaction of scorpion toxin *Buthus tamulus* at the sodium channel of the excitable membrane. *Pflugers Arch.* (1983) 397: 306-311
- (5) Vargas O, Martin MF and Rochat H. Characterization of six toxin from the venom of Moroccan scorpion *Buthus occitanus mardochei*.




Figure 5. Effects of *Androctonus crasicauda* venom (10 $\mu\text{g/ml}$) on chick biventer cervicis preparation. A: It caused a large contracture in the absence of stimulation. B: tubocurarine pretreatment prevented the effect of venom in the lack of stimulation.




Figure 6. Reversal of tubocurarine (●), Mg^{2+} (■), procaine (○) blockade by *Androctonus crasicauda* venom on responses of stimulated chick biventer cervicis nerve-muscle preparation. (A) Experiments in the presence of 3 $\mu\text{g/ml}$ venom. (B) Experiments in the presence of 10 $\mu\text{g/ml}$ venom. (mean \pm sem, $n=4$).




Figure 7. *Androctonus crasicauda* venom (10 $\mu\text{g/ml}$) markedly increased twitch height evoked by indirectly (upper panel), but not direct (lower panel) stimulation of mouse hemidiaphragm nerve-muscle preparations.

- Eur. J. Biochem.* (1987) 162: 589-599
- (6) Balozet L. Scorpionism in the Old World. In: Bucheral W. and Buckley EE. (Eds) *Venomous Animals and their Venoms. Vol. III, Venomous Invertebrates*. Academic Press, New York (1971) 349-371
 - (7) Murthy KRK and Zolfagharian H. Increased osmotic fragility of red cells after incubation at 37°C for 24hr in dogs with acute myocarditis produced by scorpion (*Buthus tamulus*) venom. *Ind. J. exp. Biol.* (1986) 24: 464-467
 - (8) Narahashi T, Shapiro BI, Deguchi T, Scuka M and Wang CM. Effects of scorpion venom on squid axon membranes. *Am. J. Physiol.* (1972) 222: 850-857
 - (9) Marshall DL and Harvey AL. Block of potassium channels and facilitation of acetylcholine release at the neuromuscular junction by the venom of the scorpion, *Pandinus imperator*. *Toxicon* (1989) 27: 493-498
 - (10) Romey G, Chicheportiche R, Lazdunski M, Rochat H, Miranda F and Lissitzky S. Scorpion neurotoxin a presynaptic toxin which affects both Na and K channels in axons. *Biochem. Biophys. Res. Commun.* (1975) 64: 115-121
 - (11) Wang GW and Strichartz G. Simultaneous modifications of sodium channel gating by two scorpion toxins. *Biophys. J.* (1982) 40: 175-179
 - (12) Simard JM, Meves H and Watt DD. Effects of toxins VI and VII from the scorpion *Centruroides sculpturatus* on the Na currents of the frog node of ranvier. *Plugers Archiv. Eur. J. Physiol.* (1986) 406: 620-628
 - (13) Barhanin J, Schmid A, Lombet A, Wheeler KP, Lazdunski M and Ellory JC. Molecular size of different neurotoxin receptors on the voltage-sensitive Na channel. *J. Biol. Chem.* (1983) 258: 700-702
Galvez ML. Purification and characterization of a unique, potent, peptidyl probe for the high conductance calcium-activated potassium channel from venom of the scorpion *Buthus tamulus*. *J. biol. Chem.* (1990) 265: 11083-11090
 - (14) Possani LD, Martin BM and Svendsen I. The primary structure of noxiustoxin: a K channel blocking peptide, purified from the venom of the scorpion *Centruroides noxius* Hoffmann. *Carslberg Res. Commun.* (1982) 47: 285-289
 - (15) Strong PN. Potassium channel toxins. *Pharmac. Ther.* (1990) 46: 137-162
 - (16) Crest M, Jacquet G, Gola, Zerrouk H, Benslimane A, Rochat H, Mansuelle P and Martin-Eauclaire MF. Kalitoxin, a novel peptidyl inhibitor of neuronal Ca-activated K channels characterized from *Androctonus mauretanicus mauretanicus* venom. *J. biol. Chem.* (1992) 267: 1640-1647
 - (17) Vatanpour H, Rowan EG and Harvey AL. Effects of scorpion (*Buthus tamulus*) venom on neuromuscular transmission in vitro. *Toxicon* (1993) 31: 1373-1384
 - (18) Rowan EG, Vatanpour H, Furman BL, Harvey AL, Tanira MOM and Gopalakrishnakone P. The effects of Indian red scorpion *Buthus tamulus* venom in vivo and in vitro. *Toxicon* (1992) 30: 1157-1164
 - (19) Del Pozo EC, Salas M and Pacheco P. Effects of scorpion venom at the neuromuscular junction. *Mems Inst. Butantan Simp. Internac.* (1966) 33: 961
 - (20) Benoit PR and Mambrini J. Action du venin de scorpion sur la jonction neuromusculaire de la grenouille. *J. Physiol.* (1967) 59: 348
 - (21) Vital Brazil O, Neder AC and Corrado AP. Effects and mechanism of action of *Tityus serrulatus* venom on skeletal muscle. *Pharmacol. Res. Commun.* (1973) 5: 137-140
 - (22) Carbone E, Prestipino G, Spadavecchia L, Franciolini F and Possani LD. Blocking of the squid axon K channel by noxiustoxin: a toxin from the venom of the scorpion *Centruroides noxius*. *Pflugers Arch.* (1987) 408: 423-431
 - (23) Ginsborg BL and Warriner JN. The isolated chick biventer cervicis nerve-muscle preparation. *Br. J. Pharmac.* (1960) 15: 410-411