**Review** Article

# Mycotoxin Contamination of Foodstuffs and Feedstuffs in Iran

Hassan Yazdanpanah<sup>*a,b*\*</sup>

<sup>*a*</sup>Pharmacology and Toxicology Department, School of Pharmacy, <sup>*b*</sup>Pharmaceutical Sciences Research Center, Shaheed Beheshti University of Medical Sciences and Health Services, Tehran, Iran.

## Classification and Occurrence of Mycotoxins

Mycotoxins are organic and complex secondary metabolites produced by various fungi species. Aflatoxins (AF), aflatoxin M1 (AFM1), ochratoxin A (OTA), deoxynivalenol (DON), fumonisins (FM), zearalenone (ZEN) and patulin (PTN) are among the most important mycotoxins. They can contaminate human foods and animal feeds through fungal growth prior to and during harvest, or as result of improper storage. Mycotoxins are mainly produced by fungal genera including Aspergillus, Fusarium, Penicillium, Claviceps and Alternaria under proper conditions (1). Major food commodities affected are cereals, nuts, dried fruits, coffee, cocoa, spices, oil seeds, dried peas and beans and fruits, particularly apples (2).

There are a number of dimensions to the problems caused by mycotoxins. The first of these relates to their implications for both human and animal health. Mycotoxicosis is mainly acquired through ingestion of contaminated moldy food. Mycotoxins can induce a wide range of effects in animals such as immunosuppression, myelosuppresion, hepatotoxicity, neurotoxicity, dermatotoxicity, estrogenicity, mutagenicity, carcinogenicity and teratogenicity. Even though mycotoxicoses, like ergotism, have been known for more than a thousand years, the causal relationship between exposure and human

\* Corresponding author:

E-mail: hasyazdan@yahoo.com

diseases remains controversial. Experimental and epidemiological evidences suggest that certain mycotoxins are associated with different forms of human cancer. For certain types of mycotoxins, high doses can result in acute intoxication, and can be either sublethal or lethal, with numerous symptoms, including various organ pathologies or even cancer (1). Due to possible health risks to the consumers, many countries have set regulatory limits on the permissible levels of different mycotoxins in foods and feeds (3).

Mycotoxins also have a significant impact on economics, by causing losses in farm animals or giving rise to difficulties in their management, or by rendering commodities unacceptable for national or international trade (4).

In this article, contamination of Iranian foods and feeds with mycotoxins using published data are reviewed. Papers retrieved through searching databases including Entrez Pubmed, ScienceDirect, Chemical Abstract and IranDoc from 1960-2005.

### Occurrence of mycotoxins in Iran

In 1997, Institute of Standard and Industrial Research of Iran (ISIRI) set maximum tolerated limit (MTL) for mycotoxins in foods and feeds (3). MTLs came into force in 2002, and currently some of the imported commodities are analyzed for mycotoxins.

## 1. Aflatoxins (B and G groups)

AF are known mutagenic, teratogenic,

carcinogenic and immunosuppressive compounds. There are 4 main toxins namely aflatoxin B1 (AFB1), aflatoxin B2 (AFB2), aflatoxin G1 (AFG1) and aflatoxin G2 (AFG2). Different food commodities including cereals, nuts, dried fruit, cocoa, spices, oil seeds, pulses and beans are contaminated with AF. In Iran, most of the research regarding AF contamination of foods and feeds have been focused on cereals and nuts, especially pistachio. The history of AF contamination of foods in Iran dates back to 1967, where AF poisoning due to bread consumption was reported in Tehran (5).

## 1.1 Occurrence of AF in nuts

Pistachio nut, as an important member of tree nuts crop, is cultivated in several countries including Iran, USA, Turkey, Greece, Syria, Italy and other countries. Pistachio nuts are a rich source of fat and contain linoleic and linolenic fatty acids, essential for the human diet (6). However, they are very prone to AF contamination and may frequently contain an elevated levels of AF (7). *Aspergillus flavus* is dominant on tree nuts including pistachio, and usually produces only AFB1 and AFB2.

In 1971, for the first time, US Food and Drug Administration (FDA) reported AF contamination in pistachio nut consignments arriving into the USA. A more systematic testing on consignments arriving from Iran and Turkey led to rejection of 36 consignments out of total of 48 tested. For managing the problem, intensive research was initiated at mycotoxins research centers in Iran. Investigations revealed that AF contamination starts when pistachio nuts are still on the trees after ripening but before harvesting. It was also found that spraying trees with captan for a period of 30 days before harvesting could eliminate the problem (8). The fact that AF detected in pistachio nuts sampled from tree prior to harvest indicated that AF might not be solely from improper shipping and storage (8-9).

FDA inspection of imported pistachio nut consignments over a long period, revealed a wide yearly variation in the amount of AF contamination. Therefore, role of climate variations on the level of AF contamination was investigated. The results showed that yearly variation is directly related to the amount of rainfall during the period of ripening to harvesting of the nuts. It is during this crucial period that AF contamination of the pistachio nuts occurs (10-11).

Over the past years, the health authorities of European Union (EU) also reported a similar problem, regarding Iran's pistachio nuts arriving at EU entry ports, and in 1997, they imposed a temporary ban on importation of pistachio nuts from Iran for two months (12-13). This was a precautionary measure, following the reported findings of the Dutch authorities, indicating that a large number of pistachio nuts consignments from Iran were contaminated with unacceptable levels of AF. Subsequent analyses by other EU member states confirmed that up to 70% of the consignments contained AF, ranging from 11 to 400 ng/g (14). Although ban on Iran's pistachio nuts was removed later, based on the 1997 decision of European Commission (EC), a special condition was imposed on Iran's pistachio nuts entering EU (13). This decision was also re-stated again on 2000 (14) and 2003 (15).

Among the 47361 pistachio nut samples analysed during March 1998-March 2001 in Iran, 28227 samples were not contaminated (Lower than the limit of detection [LOD]), 7862 samples contained AFB1 between LOD-2 ng/g, 6583 samples contained AFB1 between 2-10 ng/g, and 4689 samples contained AFB1 higher than 10 ng/g. The maximum level detected was 1426 ng/g (16-17).

In the Food and Drug Control Labs of the Iran Ministry of Health, 7926 pistachio samples were analysed during March 2001-March 2002. The data indicated that 5390 samples were not contaminated (<LOD), 1324 samples contained AFB1 between LOD-2 ng/g, 451 samples contained AFB1 between 2-10 ng/g and 761 samples contained AFB1 higher than 10 ng/g. The maximum level detected was 261.8 ng/g (18).

During March 2001-March 2002, 3629 pistachio samples were also analysed by the ISIRI affiliated Food Quality Control Labs. The results showed that 2286 samples were not contaminated (<LOD), 201 samples contained AFB1 between LOD-2 ng/g, 689 samples contained AFB1 between 2-10 ng/g, and 453

samples contained AFB1 higher than 10 ng/g. The maximum level detected was 710 ng/g (19).

Pistachio nuts produced in Iran during March 2002 - February 2003 were analyzed for presence of AF (20). In this regard, 3356 pistachio nut samples were collected. After dividing samples to sub-samples, 10068 AF analyses were carried out. Among 10068 samples analyzed, AFB1 was detected in 3699 samples (36.7%), with the mean and median of 5.9 ng/g and 0.1 ng/g. respectively (Fig. 1). The total AF (AFT) was detected in 2852 samples (28.3%) with the mean and median of 7.3 ng/g and 0.4 ng/g, respectively (Fig. 2). AFB1 level in 1191 samples (11.8%) was above the MTL of AFB1 in the pistachio nut of Iran (5 ng/g) (3). Regarding AFT, the mean contamination level (7.3 ng/g) was lower than MTL of AFT in pistachio nut in Iran (3) as well as being lower than the proposed draft maximum level of the Codex Committee on Food Additives and Contaminants for AFT (15 ng/g), and only 7.5% of samples had levels above the MTL (20). In Iran, most of the contaminated pistachio nut samples contain only AFB1 and/or AFB2 (16-20).

In recent years, some measures have been taken by the Ministries of Health and Ministry of Jihad-e-Agriculture for the prevention and reduction of AF in pistachio. However, the results show that in spite of extensive efforts made by different parties in Iran, AF contamination in pistachio nuts is still substantial and a matter of concern for both producers and consumers. Despite the fact that Iran is the number one pistachio nut producing country in the world, very few research projects on different aspects of pistachio nuts contamination with AF and ways to control them have been performed. Therefore, the most resistant variety of pistachio tree to Aspergillus is not clearly known. There is no scientific evaluation of the implementation of GAP principles on the reduction of AF contamination. Even in cases that the present data proposed hull as a source of spore and mold contamination, no plan has been implemented yet to prevent returning hull to the orchard as fertilizer. No scientific published data presented to show the effect of shipping conditions on contamination and examined the

use of different conditions, such as packaging on preventing further contamination during shipment.

However, the results of an investigation on AF contamination of the Iranian peanuts showed that among 3 varieties, the Local Gilan-Iran variety, which resists seed infection by Aspergillus, was free from aflatoxin contamination (21).

#### 1.2 Occurrence of AF in other commodities

Epidemiologic studies were undertaken in the Caspian Littoral of Iran to investigate the geographic distribution of factors (including AF) might underlie the signal differences in the incidence of esophagus cancer. In this regard, AF were analyzed in typical dietary items and the results showed no unusual levels of AF (22-23). In another study, AF contamination of cottonseed and cottonseed cake was measured (24). The results showed Iranian cottonseed cake and cottonseed were contaminated with AFB1 in 2 processes including before harvest and during storage (24).

In a 4-year study, regarding AF contamination of rice and wheat samples, negligible contamination was observed (25). Rasti et al. analyzed AFB1 in 70 imported and/or locallyproduced corn samples collected from the central storage places in Isfahan. The results showed that almost all samples were contaminated in the range of LOD - 9.9 ng/g with a mean of 5.1 ng/g (26). In 1998, fourteen barley and nine corn samples, destined for animal feed, collected from the Golestan and Mazandaran provinces in the north of Iran, were analyzed for AF using high performance liquid chromatography In corn samples, AFB1 and (HPLC) (27). AFB2 were detected in 8 (88.8%) and 6 (66.6%) samples, at a mean level of 15.83 and 2.99 ng/g (median 1.72 and 1 ng/g), respectively (Fig. 3), and AFB1 level was above the Iranian MTL of AFB1 in corn for animal feed (5 ng/g) (3). None of the corn samples contained detectable amounts of AFG1 and AFG2 (27). In another study, Tayebi and Esavi reported that one out of twenty corn samples, collected from North of Iran, was contaminated with AFB1 and AFB2, at levels of 245 ng/g and 35 ng/g, respectively (28).

According to an unpublished data, among 44 corn samples collected from pre-harvest and



**Figure 1.** Incidence of AFB1 (ng/g) in Iran pistachio nut samples analyzed by food control labs during March 2002-February 2003.

post-harvest stages as well as storage places in Golestan province, 79.5% were contaminated with AFT at mean and median levels of 52.8 ng/g and 10.2 ng/g, respectively. The AFT mean level was higher than the recommended Iranian MTL of 20 and 30 ng/g in corn intended for animal feed and human consumption, respectively. These data are in agreement with the other studies done in Iran and showed that corn produced in Northern Iran is susceptible to AF contamination. Therefore, there is an urgent need for intervention strategies to prevent and control AF contamination in corn in Golestan and Mazandaran provinces.

#### 2. Aflatoxin M1

Dairy cattle fed rations containing AFB1 excrete, in their milk, a hydroxylated metabolite known as "milk toxin" or AFM1, which is specifically found in milk and milk products. There is a general consensus that approximately 1-3% of the AFB1 initially present in the animal feeds appears as AFM1 in milk (29). AFM1 is presumed to induce liver cancer in rodents, and its carcinogenic potency in sensitive species is about one order of magnitude less than that of AFB1 (30).

Occurrence of AFM1 in milk in Iran has been investigated (31-34). During 1973-1974, Suzangar et al. (31) analyzed AFM1 and aflatoxin M2 (AFM2) in 67 milk samples from village producers and in 28 milk samples from the commercial producers in Isfahan, Iran. Of the former, 50-56.7%, and of the latter, 0-2% were contaminated, respectively, with AFM1 and AFM2 (31). In another study, AFM1 was analyzed in 73 milk samples sold to Iran Pasteurization Milk Company (32). Sixty out of 73 samples were contaminated with AFM1 at



**Figure 2.** Incidence of AFT (ng/g) in Iran Pistachio nut samples analyzed by food control labs during March 2002-February 2003.

levels higher than 50 ng/l (32). Alborzi et al. (33) investigated AFM1 contamination in pasteurized milk samples in Shiraz city. During April to September 2003, 624 pasteurized milk samples were collected from different supermarkets. AFM1 was found in all milk samples, and 17.8% of the samples had levels higher than 50 ng/l (33).

In another study, among 111 raw milk samples obtained from dairy plants in Sarab city, 76.6% of the samples were contaminated with AFM1 in the range of 15 - 280 ng/l, and the levels in 40% of the samples was higher than 50 ng/l (34).

#### 3. Ochratoxin A

OTA is a nephrotoxic and nephrocarcinogenic mycotoxin. Although OTA has been found in many food commodities (such as dried vine fruits including raisins, currants and sultanas), most surveys indicate that the highest incidences and levels of OTA are found in cereals (2).

In one study in north eastern Iran, where esophageal cancer is common, grain was sampled from wheat and barley crops at harvest and after different periods of storage in underground pits or in buildings (35). OTA was detected in only one out of 141 samples and this had much *Penicillium* contamination. No other mycotoxins were detected, although occasional samples showed unidentified toxicity to brine shrimp larvae, and isolates of *A. flavus* and *A. ochraceus* produced AF and OTA, respectively, in culture. There was little evidence for mycotoxins being implicated in esophageal cancer in north-eastern Iran (35).

In 1998, Yazdanpanah et al. (27) analyzed OTA in fourteen barley and nine corn samples (destined for animal feed) collected from Golestan and Mazandaran provinces in the north of Iran. No contamination was observed in



Figure 3. Mycotoxin Levels in corn samples from Mazandaran Province, Iran in 1998

barley samples. In corn samples, only one of the AF-contaminated samples was co-contaminated with OTA at a concentration of 0.35 ng/g (Fig. 3). This is the first report concerning natural occurrence of OTA and co-occurrence with AF in corn samples of northern Iran (27).

In the first phase of the Iran national project, concerning analysis of mycotoxins in foods in Tehran, OTA was analyzed in 18 rice (imported and locally produced) and 18 bread samples collected from retail market in 2005 (36). Seven rice samples were positive for OTA, at levels varying from 0.06 to 1.66 ng/g with a mean ( $\pm$  SD) of 0.38 ( $\pm$  0.61) ng/g. None of the samples was above the Iranian MTL of 5 ng/g (3). Regarding bread, none of the samples was found to be contaminated with OTA. This is the first report of OTA contamination in rice in Iran (36).

According to the few published data, it seems that there is a low level of contamination with OTA in foods. But, further studies needed in order to confirm the findings. In addition, publishing the available data in Ministry of Health and ISIRI for raisins intended for export which have been analyzed in their laboratories may provide valuable information.

#### 4. Zearalenone

ZEN is an estrogenic mycotoxin which frequently colonizes cereal crops worldwide. Despite its low acute toxicity, the presence of ZEN in feeds has long been a problem in animal husbandry, notably by causing infertility and related disorders in swine and sheep. Most surveys indicate that ZEN occurs primarily in corn, as well as in certain other grains and mixed feeds (37). There are a few reports dealing with ZEN contamination in Iranian foods.

During 1991-1993, 99 wheat samples,

infected with Fusarium species, were collected from the Mazandaran province (38). Analysis of ZEN in 37 samples showed ZEN contamination in the range of 3000-42600 ng/g. It should be noted that this study was performed on samples contaminated with Fusarium species, and, therefore, the result does not show a real picture of ZEN contamination in wheat samples of the Mazandaran province (38). In another study, in September 2000, 40 pre-harvest corn samples from farms of the Dashte-naz region in Mazandaran province were randomly collected and analyzed for ZEN (39). The results showed that 7.5% corn samples contained ZEA in the range of 100–212 ng/g, with a mean of 141 ng/ g. This study, which is the first report of ZEA occurrence in the Iranian corn, showed that the ZEA level was lower than Iranian MTL for this mycotoxin (39).

Oveisi et al. analyzed ZEN in 38 samples of corn flour and a cheese snack derived from it. ZEN was detected in corn flour and cheese snack samples, with an average content of 377 ng/g (maximum, 889 ng/g) and 832 ng/g (maximum, 1471 ng/g), respectively (40).

#### 5. Deoxynivalenol

The trichothecenes are a group of closely related secondary metabolites produced essentially by a broad range of *Fusarium* molds (41). One of the trichothecenes most often associated with *Fusarium* head blight is DON. It is mainly associated with cereals and can cause outbreaks of acute illness in humans (30).

During 1991-1993, 99 wheat samples, infected with *Fusarium* species, were collected from the Mazandaran province (38). Analysis of DON in 37 samples showed contamination up to 10500 ng/g. Again, like ZEN, it should be noted that this study was conducted on samples contaminated with *Fusarium* species, and, therefore, the result does not show a real picture of DON contamination in wheat samples of Mazandaran province (38).

#### 6. Fumonisins

FM are a group of mycotoxins produced by a limited number of *Fusarium* molds (42). According to the International Agency for Research on Cancer, 'toxins produced by *Fusarium moniliforme*' (the old name for *F. verticillioides*) are possibly carcinogenic to humans (group 2B carcinogens) (43). Subsequently, this assessment has been refined and fumonisin B1 (FB1) itself was evaluated as a group 2B carcinogen (44). It has been shown that FM can be found almost wherever corn is grown.

During 1998-2000, the natural occurrence of FM in the Iranian corn was investigated (45-47). In 1998, corn collected in the Mazandaran and Isfahan provinces were analyzed for FB1, fumonisin  $B_2$  (FB<sub>2</sub>) and fumonisin  $B_3$  (FB<sub>3</sub>) (45). The samples from Mazandaran province were random samples obtained from farmers' corn lots collected in September 1998, whereas those from the Isfahan province, were bought as corn cobs in the local retail market during October 1998. All 11 samples from Mazandaran showed high levels of FM contamination, with FB1 levels between 1270 ng/g and 3980 ng/g, FB2 levels between 190 ng/g and 1175 ng/g and FB3 levels between 155 ng/g and 960 ng/g (Fig. 3). Samples from Isfahan showed lower levels of contamination with 100% samples had detectable FB1 (10 ng/g -590 ng/g), two of eight samples had detectable FB2 (50 ng/g -75 ng/g) and two of eight samples had detectable FB3 (50 ng/g -75 ng/g). This was the first report of FM contamination of corn from Iran (45). The overall results of FM analysis during 1998-2000 showed that all of the samples of Mazandaran province except one were positive for FB1. A comparison of the results indicated higher mean and maximum levels during 2000 (mean: 5670 ng/g, max: 12950 ng/g) than those found during 1999 (mean: 3180 ng/g, max: 7660 ng/g) and 1998 (mean: 2270 ng/g, max: 3980 ng/g), and the average concentration of FMB1 was higher than 2000 ng/g in all the years, which is higher than the recommended Iranian MTL of 1000 ng/g for FMB1 and FMB2 (45-47). However, the mean levels of FMB1 in samples collected from markets in the Isfahan province in 1998 and 1999 (170 ng/g and 220 ng/g, respectively) were significantly lower than the levels found in corn from Mazandaran (45-46).

To investigate the incidence of esophageal cancer in the Golestan province of North-East Iran, 1349 rural and urban inhabitants of Golestan province aged between 35-80 years were invited to undergo extensive lifestyle interviews and to provide biological samples (48). In addition, samples of rice, wheat and sorghum were tested for FM contamination. However, no FM were detected in the samples analyzed (48).

For the control of FM contamination of corn, application of proper agricultural and sanitary measures and HACCP (hazard analysis and critical control point) principles will lead to the production of a better quality corn in Iran.

## 7. Patulin

PTN is a secondary metabolite produced mainly in rotten parts of apples by a wide range of fungi. Due to its mutagenic and teratogenic nature and possible health risks to consumers, many countries have regulations to reduce the level of PTN in apple products to as low as practically possible. A survey on the presence of PTN was conducted in 2002, on 65 locally produced commercial apple juice and apple juice concentrates purchased from retail outlets or producers in Iran (49). In this study, 42 apple juice and 23 apple juice concentrate samples surveyed for PTN contamination. Results of the study showed in 69% of apple juice and 78% of the apple juice concentrates, level of PTN was found to be higher than 15  $\mu$ g/l. Overall, 33% of the apple juice samples had PTN levels higher than 50  $\mu$ g/l, with a maximum level of 285  $\mu$ g/l, and 56% of the apple juice concentrates had PTN levels of higher than 50 µg/l and a maximum level of 148 µg/l (49). Although the mean concentration of PTN in apple juice samples surveyed was slightly lower than Iran MTL, fairly high incidence of PTN in these samples indicates the need for improving production techniques by industry in order to reduce the incidence and level of PTN contamination in apple juices.

#### Conclusion

Considering suitable weather and temperature conditions for fungal growth in some parts of the country (such as the Northern Part), assessment of the occurrence of mycotoxins contamination of different kinds of commodities including nuts, cereals and foods should be a priority for health policymakers. The main current concern is contamination of corn with AF and FM, and PTN contamination of apple juices, although the risk of contamination of foods and feeds with other mycotoxins should be considered.

#### References

- Boutrif E. and Bessy C. Global significance of mycotoxins and phycotoxins. In: de Koe WJ, Samson SA, van Egmond HP, Gilbert J, Sabino M (eds) Mycotoxins and Phycotoxins in Perspective at the Turn of the Millenium; Proceedings of the Xth International IUPAC symposium on Mycotoxins and Phycotoxins, Brazil (2001) 3-16
- (2) Pittet A. Natural occurrence of mycotoxins in foods and feeds – an updated review. *Revue. Med. Vet. (1998)* 149: 479-492
- (3) Food and Agriculture Organization of the United Nations. Worldwide regulations for mycotoxins in food and feed in 2003. FAO Food and Nutrition Paper, No. 81, FAO, Rome (2004)
- (4) Moss MO. Economic importance of mycotoxins recent incidence. *Int. Biodeterior. (1991)* 27: 195-204
- (5) Tabibi M. and Salehian AA. Food poisoning in bread caused by Aspergillus flavatoxin: winter of 1967, Téhéran. Acta Medica Iranica (1974) 17: 63-9
- (6) International Nut Council. URL: http://www.nuthealth. org/nutrition/nutrient100g.html. (Accessed 17 January 2006).
- (7) Jelinek CF, Pohland AE and Wood GE. Worldwide occurrence of mycotoxins in foods and feeds – an update. J AOAC International (1989) 72: 223-30
- (8) Emami A, Suzangar m and Barnett RC. Contamination of pistachio nuts with aflatoxins while on the trees and in storage. Zesz Probl Postepow Nauki Rolniczych (1977) 189: 135-40
- (9) Dickens JW. and Welty RE. Fluorescence in pistachio nuts contaminated with aflatoxin. J. Am. Oil Chem. Soc. (1975) 52: 448-50
- (10) Mojtahedi H, Danesh D, Haghighi B and Barnett R. Postharvest pathology and mycotoxin contamination of Iranian pistachio nuts. *Phytopathol (1978)* 68: 1800-4
- (11) Danesh D, Mojtahedi H, Barnett R and Campbell A. Correlation between climatic data and aflatoxin contamination of Iranian pistachio nuts. *Phytopathol.* (1979) 69: 715-6
- (12) Commission Decision 97/613/EC of 8 September 1997 on the temporary suspension of imports of pistachios and certain products derived from pistachios originating in, or consigned from Iran, *Off. J. EU* (1997) L 248, 33
- (13) Commission Decision 97/830/EC of 11 December 1997 repealing Decision 97/613/EC and imposing special conditions on the import of pistachios and certain products derived from pistachios originating in

or consigned from Iran, Off. J. EU (1997) L 343, 30

- (14) Ministry of Agriculture, Fisheries and Foods, Iran pistachio nuts – imports suspended. Food Safety Inform. Bull., No. 89 (1997)
- (15) Commission Decision 2003/551/EC, amending Decision 97/830/EC repealing Decision 97/613/EC and imposing special conditions on the import of pistachios and certain products derived from pistachios originating in or consigned from Iran, Off. J. EU (2003) L 187, 43-6
- (16) Iranian Ministry of Health and Medical Education, The situation of aflatoxin contamination in pistachio during March 1998 - March 2001 (2001)
- (17) Institute of Standards and Industrial Research of Iran, The situation of aflatoxin contamination in pistachio during March 1998 - March 2001(2001)
- (18) Iranian Ministry of Health and Medical Education, Annual report: The situation of aflatoxin contamination in pistachio during March 2001 - March (2002)
- (19) Institute of Standards and Industrial Research of Iran, Annual report: The situation of aflatoxin contamination in pistachio during March 2001-March 2002 (2002)
- (20) Yazdanpanah H., Cheraghali AM, Doraki N., Abouhossain G., Hassibi M., Ali-abadi S., Aliakbarpoor M., Amirahmadi M., Askarian A., Fallah N., Hashemi T., Jalali M., Kalantari N., Khodadadi E., Maddah B., Mohit R., Mohseny M., Phaghihy Z., Rahmani A., Setoodeh L., Soleimany E. and Zamanian F. Incidence of Aflatoxins in Iran Pistachio Nuts. Submitted to *Food Chem. Toxicol. (2005)*
- (21) Tavasolian B. Native Iranian peanut resistance to seed infection by Aspergillus. J. Agric. Food Chem. (1977) 25: 1422-3
- (22) Joint IRAN-IARC Study Group. Esophageal cancer studies in the Caspian littoral of Iran: results of population studies - a prodrome. J. Nat. Cancer Inst. (1977) 59: 1127-38
- (23) Hormozdiari H, Day NE, Aramesh B and Mahboubi E. Dietary factors and esophageal cancer in the Caspian Littoral of Iran. *Cancer Res.* (1975) 35: 3493-8
- (24) Emami A, Suzangar M and Barnett RC. Contamination of cotton seeds and of cotton seed cake by aflatoxin. *Ann Nutr Aliment. (1977)* 31: 531-7
- (25) Emami A., Suzangar M and Barnett R. Aflatoxin contamination of pistachio nuts, village milk, wheat and rice. *Pazhoohandeh (1977)*16: 139-51 (persian)
- (26) Rasti M. Determination of aflatoxin B1 in corn of central storage places in Isfahan. *Chakavak (2001)* 10: 124-5 (Persian)
- (27) Yazdanpanah H, Miraglia M, Calfapietra FP and Brera C. Natural occurrence of aflatoxins and ochratoxin A in corn and barley from Mazandaran and Golestan provinces in north of I. R. Iran. *Mycotoxin Res. (2001)* 17: 21-30
- (28) Tayebi, J and Esavi C. Investigation on aflatoxins B and G contamination in corn in the field. Proceedings of the 13<sup>th</sup> Iran Plant Protection Congress, Islamic Republic of IRAN: Karaj Junior College of Agriculture,

(1998) p. 92

- (29) van Egmond HP. Aflatoxin M1: occurrence, toxicity, regulation. In: Mycotoxins in dairy products. van Egmond HP. (Ed.), Elsevier, Amsterdam, (1989) p. 11-55.
- (30) Bolger M, Coker RD, DiNovi M, Gaylor D, Gelderblom W, Olsen M, Paster N, Riley RT, Shephard G, Speijers GJA. Safety evaluation of certain mycotoxins in food. Fifty-sixth meeting of the Joint FAO/WHO Expert Committee on Food Additives, FAO Food and Nutrition Paper 74, (2001)
- (31) Suzangar M, Emami A and Barnett R. Aflatoxin contamination of village milk in Isfahan, Iran. *Tropical Science (1976)* 18: 155-9
- (32) Bokaei S., Determination of aflatoxin M1 in milk sold to Iran pasteurized Milk Factory. *Pazhoohesh va Sazandeghi (1999)* 12: 163-5 (persian)
- (33) Alborzi S, Pourabbas B, Rashidi M and Astaneh B. Aflatoxin M1 contamination in pasteurized milk in Shiraz (south of Iran). *Food Cont. (2006)* 17: 582-4
- (34) Kamkar A. A study on the occurrence of aflatoxin M1 in raw milk produced in Sarab city of Iran. Food Cont. (2005) 16: 593-9
- (35) Lacey J. The microbiology of cereal grains from areas of Iran with a high incidence of oesophageal cancer. J. Stored Prod. Res. (1988) 24: 39-50
- (36) Yazdanpanah H, Aboul-fathi F, Khoddam A and Foroutan SM. Rapid determination of ochratoxin A in rice and bread by immunoaffinity column clean-up and liquid chromatography using a monolithic column with fluorescence detection. Abstracts of The World Mycotoxin Forum: The Third Conference, Noordwijk, The Netherlands, (2005) p. 118.
- (37) Kuiper-Goodman T, Scott PM and Watanabe H. Risk assessment of the mycotoxin zearalenone. *Biomed. Environ. Sci.* (1989) 2:179-248
- (38) Zamani-Zadeh, H. R., and Khoursandi, H. Occurrence of Fusarium species and their mycotoxins in wheat in Mazandaran province. (1995) Iran J. plant Pathol., 31: 23-37
- (39) Hadiani MR, Yazdanpanah H,\_Ghazi Khansari M, Cheraghali AM and Goodarzi M. Survey of the natural occurrence of zearalenone in maize from northern Iran by thin-layer chromatography densitometry. *Food Addit. Contam. (2003)* 20: 380-5
- (40) Oveisi M, Hajimahmoodi M, Memarian S, Sadeghi

N and Shoeibi S. Determination of zearalenone in corn flour and a cheese snack product using highperformance liquid chromatography with fluorescence detection. *Food Addit. Contam. (2005)* 22:443-8

- (41) Moss MO. Mycotoxins. *Mycol. Res. (1996)* 100: 523-33
- (42) Shephard GS, Thiel PG, Stockenstrom S and Sydenham EW. Worldwide survey of fumonisin contamination of corn and corn-based products. J. AOAC International (1996) 79: 671-87
- (43) IARC. Toxins derived from *Fusarium moniliforme*: Fumonisin  $B_1$  and  $B_2$  and fusarin C. Lyon: IARC, (1993)
- (44) IARC, 2002. Fumonisin B<sub>1</sub>. In: IARC Monographs on the Evaluation of Carcinogenic Risks to Humans, Some Traditional Herbal Medicines, Some Mycotoxins, Naphthalene and Styrene, vol. 82. IARC, Lyon, (2002) 301-66
- (45) Shephard GS, Marasas WFO, Leggott NL, Yazdanpanah H, Rahimian H and Safavi N. Natural occurrence of fumonisins in corn from Iran. J. Agric. Food Chem. (2000) 48: 1860-4
- (46) Shephard GS, Marasas WFO, Yazdanpanah H, Rahimian H, Safavi N, Zarghi A, Shafaati AR and Rasekh HR. Fumonisin B1 in maize harvested in Iran during 1999. *Food Addit. Contam. (2002)* 19: 676-9
- (47) Yazdanpanah H, Eskandari Gheidari P, Zarghi A, and Mirkarimi SK. Survey of fumonisin B1 contamination of corn in Northern Iran during 2000. *Iranian J. Pharm. Res. (2006)* **In Press.**
- (48) Pourshams A, Saadatian-Elahi M, Nouraie M, Malekshah AF, Rakhshani N, Salahi R, Yoonessi A, Semnani S, Islami F, Sotoudeh M, Fahimi S, Sadjadi AR, Nasrollahzadeh D, Aghcheli K, Kamangar F, Abnet CC, Saidi F, Sewram V, Strickland PT, Dawsey SM, Brennan P, Boffetta P, and Malekzadeh R. Golestan cohort study of oesophageal cancer: feasibility and first results. *Br. J. Cancer (2005)* 92: 176-81
- (49) Cheraghali AM, Mohammadi HR, Amirahmadi M, Yazdanpanah H, Abouhossain G, Zamanian F, Ghazi Khansari M and Afshar M. Incidence of patulin contamination in apple juice produced in Iran. *Food Cont.* (2005) 16: 165-7

This article is available online at http://www.ijpr-online.com