

Uranyl Acetate Induces Oxidative Stress and Mitochondrial Membrane potential collapse in the Human Dermal Fibroblast Primary Cells

Bahram Daraei^{c*}, Jalal Pourahmad^{a,b}, NedaHamidi-Pour^c,
Mir-Jamal Hosseini^{a,d}, Fatemeh Shaki^a and Masoud Soleimani^c

^aSchool of Pharmacy, Shaheed Beheshti University of Medical Sciences, Tehran, Iran.
^bPharmaceutical Sciences Research Center, Shaheed Beheshti University of Medical Sciences, Tehran, Iran. ^cDepartment of Toxicology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran. ^dStudent Research Committee, School of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Abstract

Cytotoxicity of depleted uranium, as a byproduct of military has been came to spotlight in recent decades. DU is known as a chemical rather than radioactive hazard and efforts to illustrating its mechanism is undergo, but the precise complete molecular mechanisms are still unclear. Recent studies showed that uranium induces biological changes in many different target tissues, such as the kidney, brain and skin. The aim of this study was to assess the impact of depleted uranium exposure at the cellular level in the human dermal fibroblast primary cells. The human dermal fibroblast primary cells incubated with different concentration (250-750 μ M) of depleted uranium.

Cytotoxicity and mitochondrial function in this cell lines were determined with the LDH leakage assay and the MTT test respectively. MDA levels were measured for determination of Lipid peroxidation in DU treated cells. Besides glutathione depletion and apoptosis phenotype detection were also assessed to complete the mechanistic screening. Results showed that the cell viability ameliorates in concentration and time dependent manners following in 24, 48 and 72 h incubation with DU. Moreover the significant increase in lipid peroxidation and significant decrease in cellular GSH recorded in DU treated human dermal fibroblast primary cells suggesting the preoxidant effect of uranyl ions. Cytoprotective effects of N-acetylcysteine (NAC) and dramatic decrease of cell viability in buthionin sulfoxamid (BSO) pretreated cells indicated the possibility of a critical role for glutathione system in DU detoxification. Death pattern, in fibroblast cells following DU treatment was varied from apoptosis to necrosis while the time and concentration increased.

Since ROS formation is the initiation step for cell apoptosis, the present studies suggest Uranyl-induced toxicity in the human dermal fibroblast primary cells originated from oxidative stress and lead to occurrence of programmed cell death.

Keywords: Uranium; Fibroblast; Lipid peroxidation; Oxidative stress.

* Corresponding author:

E-mail: bdaraei@modares.ac.ir

Introduction

Uranium (U) is a ubiquitous environmental trace metallo-element which is found in small amounts in food and water supplies as a non-essential inorganic component (1). The U remaining after removal of the enriched fraction is referred to as depleted U (DU) (2, 3).

It has been documented that inhalation is the most common route of uranium exposure. But uranium can also be absorbed through the skin or wounds, ingested with food or drink, and injected into the system intravenously (4). In recent years, it has been noticed that depleted uranium (DU) in munitions is also another route of exposure to the heavy metal through contact with the skin (5).

The most important toxic mechanism that suggested for uranium is involvement of reactive oxygen species (ROS). ROS are known to play a dual role in biological systems, since they can be either harmful or beneficial to living systems (6). The harmful effects of ROS are balanced by the antioxidant action of non-enzymatic antioxidants in addition to antioxidant enzymes. A number of studies have focused on metal-induced toxicity and carcinogenicity, emphasizing their role in the generation of ROS in biological systems and the significance of this therein (7, 8).

Previous studies also showed that oral uranyl acetate (UA) administration increased TBARS in kidneys and testes (9). Other studies demonstrated that chronic uranyl nitrate (UN) ingestion resulted in an increase in the levels of free radicals (10), brain lipid peroxidation (LPO) (11) and also uranium induce oxidative stress in lung epithelial cells and loss of antioxidant response (5). In our previous study, UA caused rapid glutathione oxidation, ROS formation, lipid peroxidation and decreased mitochondrial membrane potential in isolated rat hepatocytes (12). These may be based upon uranium related induction of cellular oxidative stress (13). Taken together with the studies on uranium toxicity it reveals that DU could be an environmental health hazard and requires further investigation to figure out the precise mechanism of its action on its many different target organs. Considering that skin contact is a way of the exposure to uranium, we have investigated the effect of DU

on human dermal fibroblast primary cells and to study the response due to induction of oxidative stress.

Experimental

Chemicals

Uranyl acetate and all other chemicals were obtained from the Sigma Chemical Co. with the highest commercial grade available unless otherwise stated.

Cell Cultures and treatments

The human dermal fibroblast primary cells obtained from National Cell Bank of Iran, Pasteur Institute of Iran (Tehran, Iran) was cultured in DMEM medium supplemented with 10% heat inactivated FBS, 100 IU/mL penicillin, 100 µg/mL streptomycin and 2 mM L-glutamine and maintained in a humidified atmosphere with 5% CO₂ at 37°C. The cultured cells were sub-cultured twice each week and the exponentially growing cells were used for all treatments. Experiments were performed in 4 groups of cells as follow: Group I: control group, Group II, III and IV were pretreated with different dose of uranyl acetate (250-500-750 µM). Only one dose of uranyl acetate was added to each medium.

Lactate dehydrogenase (LDH) leakage assay

Cytotoxicity induced by uranyl acetate was assayed by LDH leakage into the culture medium. Following exposure to the uranyl acetate the fibroblast cells medium was aspirated and centrifuged at 3000 rpm for 5 min for obtain a cell free supernatant. LDH activity in the medium was measured by specific commercial kit from Sigma-Aldrich Co. The viability was measured spectrophotometrically in absorbance at 340 nm. Cell viability was determined relative to the untreated control cells (14).

Lipid peroxidation

For this assay human dermal fibroblast cells in exponential growth phase (1×10^6 cells) were seeded in 75 cm² cell culture flasks. After reaching to 80% confluence, cells were treated with 250, 500, 750 µM uranyl acetate. In the time intervals of 24, 48 and 72 h cells were harvested

Table 1. Mitochondrial function assessment for the human dermal fibroblast primary cells following exposure to uranyl acetate for 24, 48 and 72 h based on the dose-response curves as derived from the MTT assay. Succinate dehydrogenase activity inhibited by U in a dose- and time-dependent manner (* $p < 0.05$).

Treated cells	LDH leakage		
	24 h	48 h	72 h
Control	0 ± 3	3 ± 4	4 ± 6
U (250 μM)	*12 ± 4	34 ± 5*	43 ± 5*
U (500 μM)	23 ± 8*	48 ± 9*	59 ± 8*
U (750 μM)	31 ± 3*	52 ± 6*	82 ± 5*

Data represent the means of three separate experiments (±SD).

* $p < 0.05$, significantly different when compared with control.

with a rubber policeman and collected cells were centrifuged at 1000 g for 10 min at 40°C. The supernatant were removed and cell pellet were suspended in cold PBS and sonicated. 2 mL TBA solution (3.75 g TCA, 92.5 g TBA and 0.7 mL HCl in 100 mL distilled water) were added to 1 mL cell suspension and were heated at 90°C for 30 min. Calorimetric absorption was measured at 530 nm (15).

MTT assay

Fibroblast cells were suspended in RPMI 1640 at 10^6 cells/mL. A sample of 100 μL of the cell suspension was seeded into a 96-well plate. After an overnight culture, cells were exposed to uranyl acetate (0–750 μM) for 24, 48 and 72 h. After incubation with MTT (0.5 mg/mL) for 24 h, the medium was removed and 150 μL of DMSO was added to dissolve the formazan crystals. Absorbance was measured at 540 nm using an ELISA microplate reader. Mitochondrial function was determined relative to the untreated control cells. Data represent the mean value and standard deviation of triplicate assays in at least one experiment (16).

Intracellular GSH assessment

GSH content was determined according to the spectrophotometer method for the human dermal fibroblast cells. The content of GSH was determined using DTNB as indicator (17).

Apoptosis phenotype detection test

Cells were plated in 96-well plates at a concentration of 5×10^3 cells/well. 24 h after plating, they were washed twice with 200 μL of serum-free medium and were starved by incubation in serum-free medium for 15 h at

37 °C in a 5% CO₂ incubator. After starvation, cells were treated with 250, 500, 750 μM uranyl acetate 24, 48 and 72 h. Apoptosis was analyzed using Biovision Annexin-FITC detection kit. At the end of treatment, media from control and uranyl-treated cells were discarded and the cells were washed with 1× phosphate-buffered saline (PBS). Then 200 μL buffer solution containing 5 μg/mL annexin-FITC was added to 3 well. In order to detection necrosis, 200 μL buffer solution containing 5 μg/mL PI was added to 3 other wells. After 5 min the supernatant discarded and cells were washed with 1× PBS. 100 μL PBS was added to each well. The fluorescence emission at 530 nm (for annexin v-FITC) and 640 nm (for PI) was measured using a synergy fluorimeter. When calculating the apoptosis and necrotic cell percents it was considered that cells that are apoptotic bind annexin V and necrotic cells with a disrupted plasma membrane display uptake of both annexin V and propidium iodide (18, 19).

Statistical analysis

Levene's test was used to check the homogeneity of variances. Data were analyzed using one-way analysis of variance (ANOVA) followed by Tukey's HSD as the post-hoc test. Results were presented as mean ± SD of triplicate samples. The minimal level of significance chosen was $p < 0.05$.

Results

The human dermal fibroblast primary cells were exposed to different concentration of uranyl acetate (0-750 μm) for 24, 48 and 72 h

Table 2. TBARs level in the human dermal fibroblast primary cells at different concentration and times after uranyl acetate exposure.

Treated cells	MDA($\mu\text{mol}/\text{mg}$ protien)		
	Incubation time		
	24 h	48 h	72 h
Control	3.26 \pm 0.2	3.21 \pm 0.1	3.41 \pm 0.1
U (250 μM)	3.59 \pm 0.1	5.7 \pm 0.17*	12.55 \pm 0.38*
U (500 μM)	4.13 \pm 0.12	6.9 \pm 0.21*	19.33 \pm 0.59*
U (750 μM)	4.87 \pm 0.15	8.96 \pm 0.2*	32.86 \pm 1.01*

Data represent the means of three separate experiments (\pm SD).

* $p < 0.05$, significantly different when compared with control.

and cytotoxicity was determined with the LDH leakage assay. The viability results were shown in Table 1. Our results showed the EC_{50} value of 400 μM using the LDH leakage assay when fibroblast cells were exposed to uranyl acetate in 72 h. On the other hand, pretreatment of the human dermal fibroblast primary cells with NAC significantly increased in viability by DU (750 μM) in 24 and 72 h ($p < 0.05$, Figure 1).

The extent of lipid peroxidation was measured using of thiobarbituric acid reactive substances (TBAR_s) formation assay (15). The MDA concentration was significantly raised in the human dermal fibroblast primary cells while treated with different concentration of uranyl acetate at different sampling times (24, 48 and 72 h). ($p < 0.05$; Table 2).

As shown in Table 3, GSH levels of DU-treated cells were significantly decreased as concentration and time dependent manner ($p < 0.05$, Table 3) that our result is similar with previous studies. On the other hand, rapid decrease of cell viability in buthionin

sulfoxamid (BSO) pretreated cells that depleted cellular content of GSH, indicated the possibility of a critical role for glutathione system in DU detoxification.

The MTT dye is metabolized by viable mitochondria to a colored product and can be detected photometric instrument. Thus, the extent of MTT metabolism is an indicator of mitochondrial function (20). So we used MTT test for assessment of mitochondrial function in fibroblast cells after exposure of different concentration of uranyl acetate. As shown in Table 3, mitochondrial function significantly decreased as concentration and time dependent manner.

In detection of apoptosis, DU also increased apoptosis signaling in the human dermal fibroblast primary cells following 24, 48 and 72 h after incubation (Table 5). After initiation of apoptosis, phosphatidyl serine gradually appears on the outer leaflet of the plasma membrane of primary cells due to inhibition of Mg-ATP dependent amino phospholipid translocase

Table 3. Effect of uranyl acetate on mitochondrial function (succinate dehydrogenase activity) in the human dermal fibroblast primary cells for 24, 48 and 72 h.

Treated cells	MTT assay		
	24 h	48 h	72 h
Control	0	2 \pm 1	3 \pm 1
U(250 μM)	9 \pm 5	30 \pm 7*	40 \pm 6*
U(500 μM)	21 \pm 3*	43 \pm 8*	54 \pm 5*
U(750 μM)	27 \pm 7*	49 \pm 6*	76 \pm 9*
NAC + U(750 μM)	0 \pm 6†	46 \pm 3	59 \pm 9†

Data represent the means of three separate experiments (\pm SD).

* $p < 0.05$, significantly different when compared with control.

† $p < 0.05$, significantly different when compared with U (750 μM)

Table 4. Effect of Uranyl acetate on GSH level in the human dermal fibroblast primary cells at different times.

Treated cells	GSH ($\mu\text{mol}/\text{mg}$ protien)		
	Incubation time		
	24 h	48 h	72 h
Control	2.01 \pm 0.04	2. \pm 0.01	2 \pm 0.01
U (250 μM)	1.81 \pm 0.05*	1.41 \pm 0.07*	0.69 \pm 0.02*
U (500 μM)	1.7 \pm 0.06*	1.16 \pm 0.05*	0.44 \pm 0.01*
U (750 μM)	1.48 \pm 0.05*	0.98 \pm 0.03*	0.3 \pm 0.01*

Data represent the means of three separate experiments (\pm SD).

*Significant difference in comparison with control cells ($p < 0.05$).

giving off red fluorescence. The human dermal fibroblast primary cells that bind to Annexin V, which is representative of cells undergoing apoptosis and necrotic cells with a disrupted plasma membrane display uptake of both annexin V and propidium iodide. Du in different concentration showed significant increase in the number of apoptotic phenotype in the human dermal fibroblast primary cells compared to control (Table 5).

Discussion

Oxidative stress, an imbalance between free radical generation and the antioxidant defense system, represents a common threat and danger for all aerobic organisms. ROS can be generated by endogenous physiological mechanisms or after exposure to exogenous compounds (21, 22). DU is one of the external sources that invariably showed a number of indicators of oxidative stress such as elevation of ROS formation, induction of lipid peroxidation and GSH depletion (5, 9-11, 23). In order to determine if oxidative stress could be involved in DU cytotoxicity in fibroblast cells, we have attempted to elucidate the molecular events that occurred during U-induced oxidative stress in the human dermal fibroblast primary cells after exposure to different concentration of DU.

The LDH leakage assay is based on release of the enzyme (14). In the present work the viability assays performed using LDH leakage assay test. It noted that in order to avoid overestimation or underestimation of the toxicity of a substance, incubations with various concentrations at many time points are required. Mortality that were observed after 24, 48 and 72 h incubation were

31, 52 and 82 percent respectively that we found a LC_{50} of 400 μM after 72 h incubation with DU. Also, as demonstrated in results, we confirmed that DU could induce oxidative stress in fibroblast cells when cells treatment with at least 250 μM DU. Among oxidative damage induced by ROS, the lipid membrane is a preferential target, causing lipid peroxidation which we showed that the MDA levels in fibroblast cells were increased as time and concentration dependent manner that again confirmed the role of ROS in DU toxicity and were parallel to cytotoxicity associated with uranyl acetate. The induction of lipid peroxidation by DU already has been reported in several studies on various cell types (9, 11).

It has been demonstrated that reductive activation of U (VI) by reduced CYP 2E1, or NADPH- P_{450} reductase and glutathione to U(V) and finally U (IV), provides enough redox cycling to generate considerable amounts of ROS. On the other hand, U(VI) can also cause mitochondrial membrane damage and significant collapse of mitochondrial membrane potential (12). So, mitochondria can be considered as a target for DU cytotoxicity, therefore we performed MTT assay test for evaluation of mitochondrial function. The present data showed that 24-72 h exposure of the human dermal fibroblast primary cells with DU significantly decline the mitochondrial functionality that showed using MTT dye reduction, which reveals the adverse impact of DU in mitochondrial function.

The balance between free radical generation and the antioxidant systems is the key for maintaining a physiological intracellular redox status. Glutathione (GSH) is one of the main components of the cellular antioxidant systems

Table 5. Effect of Uranyl acetate on apoptosis and necrosis phenotype in the human dermal fibroblast primary cells at different times.

Treated cells	Incubation time								
	24 h			48 h			72 h		
	Cell death%	A %	N%	Cell death%	A %	N%	Cell death%	A %	N%
Control	0	0	0	3	97 ± 1	3 ± 1	8	93 ± 3	7 ± 2
U (250 µM)	11	98 ± 5	2 ± 0	30	82 ± 3*	18 ± 2*	42	74 ± 3*	26 ± 2*
U (500 µM)	20	92 ± 3	8 ± 1*	45	80 ± 1*	20 ± 1*	54	52 ± 2*	48 ± 1*
U (750 µM)	32	89 ± 3	11 ± 3*	48	73 ± 4*	27 ± 3*	72	20 ± 1*	80 ± 1*

A: Apoptosis, N: Necrosis. Data represent the means of three separate experiments (±SD).

*Significant difference in comparison with control cells ($p < 0.05$).

that is critical to defense against reactive oxygen species (24). In this basis, we assayed the GSH content of the human dermal fibroblast primary cells exposed to different concentration of DU. As our data showed, GSH oxidation were significantly increased after 24, 48 and 72 h treatment with DU. Our results are in agreement with previous studies that showed glutathione oxidation was occurred as a consequence of DU induced-ROS formation (12). Generation of ROS has been widely viewed as being a common pathway to cell death from a variety of injurious agents (25), and apoptosis can be triggered by ROS (26, 27). Present work showed that the rate of apoptosis in fibroblast cells exposed to lower concentration of DU was significantly higher than the rate of necrosis. DU induced apparently more apoptotic death than necrotic death in fibroblast cells in low concentration but in longer incubation time and/or higher concentration, necrosis was the prominent mode of cell death in fibroblast cells.

In conclusion, our results suggest that oxidative stress could be involved in cytotoxicity of DU in fibroblast cells. Glutathione and other antioxidants could protect the fibroblast cells against DU- induced cytotoxicity.

References

- (1) Bleise A, Danesi PR, and Burkart W. Properties, use and health effects of depleted uranium (DU): a general overview. *J. Environ. Radioact.* (2003) 64: 93-112.
- (2) WHO, *Depleted Uranium: Sources, Exposure and Health Effects - Full Report.* In WHO, in *Department of Protection of the Human Environment*. 2001, World Health Organization: Geneva.
- (3) U.S.EPA. *EPA facts about U*. United States Environment Protection Agency 2002 [cited; Available from: <http://www.epa.gov/superfund/resources/radiation/pdf/U.pdf>].
- (4) Sanchez DJ, Belles M, Albina ML, Sirvent JJ, and Domingo JL. Nephrotoxicity of Simultaneous Exposure to Mercury and Uranium in Comparison to Individual Effects of These Metals in Rats. *Biol Trace Elem Res.* (2001) 84: 139-54.
- (5) Periyakaruppan A, Kumar F, Sarkar S, Sharma CS, and Ramesh GT. Uranium induces oxidative stress in lung epithelial cells. *Arch Toxicol.* (2007) 81: 389-95.
- (6) Valko M, Izakovic M, Mazur M, Rhodes CJ, and Telser J. Role of oxygen radicals in DNA damage and cancer incidence. *Mol. Cell. Biochem.* (2004) 266: 37-56.
- (7) Ibrahim D, Froberg B, Wolf A, and Rusyniak DE. Heavy metal poisoning: clinical presentations and pathophysiology. *Clin. Lab. Med.* (2006) 26: 67-97.
- (8) Valko M, Rhodes CJ, Moncol J, Izakovic M, and Mazur M. Free radicals, metals and antioxidants in oxidative stress induced cancer. *Chemico-Biol. Interact.* (2006) 160: 1-40.
- (9) Linares V, Bell'es M, Albina ML, Mayayo E, Sanchez DJ, and Domingo JL. Combined action of U and stress in the rat. II. Effects on male reproduction. *Toxicol Lett.* (2005) 158: 186-95.
- (10) Taulan M, Paquet F, Maubert C, Delissen O, Demaille J, and Romey MC. Renal toxicogenomic response to chronic uranyl nitrate insult in mice. *Environ. Health Perspect.* (2004) 112: 1628-35.
- (11) Briner W and Murray J. Effects of short-term and long-term depleted U exposure on open-field behavior and brain lipid oxidation in rats. *Neurotoxicol. Teratol.* (2005) 27: 135-44.
- (12) Pourahmad J, Ghashang M, Ettehadi HA, and Ghalandari R. A search for cellular and molecular mechanisms involved in depleted uranium (DU) toxicity. *Environmental Toxicology.* (2006) 21: 349-54.
- (13) Taulan M, Paquet F, Maubert C, Delissen O, Demaille J, and Romey MC. Renal toxicogenomic response to chronic uranyl nitrate insult in mice. *Environ Health Perspect.* (2004) 112: 1628-35.
- (14) Decker T and Lohmann-Matthes ML. A quick

- and simple method for the quantitation of lactate dehydrogenase release in measurements of cellular cytotoxicity and tumor necrosis factor (TNF) activity. *J. Immunol. Methods.* (1988) 15: 61-9.
- (15) Reilly CA and Aust SD. Measurement of Lipid Peroxidation. *Current Protocols in Toxicology: John Wiley & Sons.* (1999)
- (16) Mosmann T. Rapid colorimetric assay for cellular growth and survival :Application to proliferation and cytotoxicity assay. *J. Immunol. Methods.* (1983) 65: 55-63.
- (17) Sadegh C and Schreck RP. The Spectroscopic Determination of Aqueous Sulfite Using Ellman's Reagent. *MURJ* (2003) 8: 39-43.
- (18) Tait JF, Smith C, and Wood BL. Measurement of Phosphatidylserine Exposure in Leukocytes and Platelets by Whole-Blood Flow Cytometry with Annexin V. *Blood Cells, Molecules, and Diseases* (1999) 25: 271-8.
- (19) Zhivotovsky B, Samali A, and Orrenius S. Assessment of Cell Toxicity. *Current Protocols in Toxicology; John Wiley & Sons.* (1999)
- (20) Wang C, Sadovova N, Ali HK, Duhart HM, Fu X, Zou X, Patterson TA, Binienda ZK, Virmani A, Paule MG, jr WS, and Ali ASF. L-Carnitine Protects Neurons From 1-Methyl-4-Phenylpyridinium-Induced Neuronal Apoptosis In Rat Forebrain Culture. *Neuroscience.* (2007) 144: 46-55.
- (21) Valko M, Morris H, and Cronin MT. Metals, toxicity and oxidative stress. *Curr. Med. Chem.* (2005) 12: 1161-208.
- (22) Pourahmad J, Hosseini MJ, Eskandari MR, and Rahmani F. Involvement of Four Different Intracellular Sites in Chloroacetaldehyde- Induced Oxidative Stress Cytotoxicity. *Iranian J. Pharm. Res.* (2012) 11: 265-76.
- (23) Pourahmad J, Shaki F, Tanbakosazan F, Ghalandari R, Etehadhi HA, and Dahaghin E. Protective effects of fungal β -(1 \rightarrow 3)-D-glucan against oxidative stress cytotoxicity induced by depleted uranium in isolated rat hepatocytes *Hum. Exp. Toxicol.* . (2011) 30: 173-81.
- (24) McCord JM. The evolution of free radicals and oxidative stress. *Am. J. Med.* (2000) 108: 652-9.
- (25) Ames BN, Shigenaga MK, and Hagen TM. Oxidants, antioxidants, and the degenerative diseases of aging. *Proc. natn. Acad. Sci. U.S.A.* (1993) 90: 7915-22.
- (26) Wood KA and Youle RI. The role of free radicals and p53 in neuron apoptosis in vivo. *J. Neurosci.* (1995) 15: 5851-7.
- (27) Pourahmad J, Mortada Y, Eskandari MR, and Shahraki J. Involvement of Lysosomal Labilisation and Lysosomal/mitochondrial Cross-Talk in Diclofenac Induced Hepatotoxicity. *Iranian J. Pharm. Res.* (2011) 10: 877-87.

This article is available online at <http://www.ijpr.ir>

Archive