

## Biotechnological Approach to Evaluate the Immunomodulatory Activity of Ethanolic Extract of *Tinospora cordifolia* Stem (Mango Plant Climber)

Vaibhav Aher<sup>a\*</sup> and Arun Kumar Wahi<sup>b</sup>

<sup>a</sup>Institute of Pharmacy, NIMS University, Jaipur, 303121, India. <sup>b</sup>College of Pharmacy, MIT, Moradabad-244001, India.

### Abstract

The present study was designed to investigate the immunomodulatory activity of the ethanolic extract of *Tinospora cordifolia* (Family: Menispermaceae) stem (climbing shrub, mango plant) at cellular level. For antioxidant study, the liver mitochondria were separated and the concentration of enzymes like lipid peroxidation (LPO), reduced glutathione (GSH), catalase (CAT) and superoxide Dismutase (SOD) were estimated; melatonin secretion characterization was carried out through SDS-PAGE. The spleen lymphocyte proliferation assay was performed through measuring its optical density at 570 nm using Elisa Reader. The cytokines viz. IL-2, IL-10 and TNF- $\alpha$  expression in spleen cells were determined through Real Time PCR. *Tinospora cordifolia* (*Tc*) ethanolic extract (100 mg/Kg/p.o.) increased the level of liver mitochondrial enzymes like GSH, CAT and SOD but decreased the level of LPO in liver as compared to the vehicle, SRBC and cyclophosphamide-treated groups. The secretion of melatonin via pineal gland was enhanced with *Tc* treatment. The extract also increased the spleen lymphocyte proliferation. In RT-PCR analysis, the expression of cytokines viz. IL-2, IL-10 and TNF- $\alpha$  was more in *Tc*-treated animals than vehicle and cyclophosphamide treatment. Hence, the study confirms the immunomodulatory activity of *Tc* stem through altering the concentration of antioxidant enzymes, increasing T and B cells and antibody which play an important role in immunity, enhancing the concentration of melatonin in pineal gland and increasing the level of cytokines like IL-2, IL-10 and TNF- $\alpha$  which plays an important role in immunity.

**Keywords:** *Tinospora cordifolia* (*Tc*); Immunomodulator; Antioxidant enzymes; Lymphocyte proliferation; Real time PCR.

### Introduction

In recent times, focus on plant research has been intensified all over the world and large amount of evidence has been collected to show the immense potential of medicinal plant used in various traditional systems (1). The immune system is involved in the etiology as well as pathophysiologic mechanism of many diseases

(2). The modulation of immune responses to alleviate the diseases has been of interest for many years and the concept of 'Rasayana' in Ayurveda is based on the related principles (3). Further immunomodulation using medicinal plants can provide an alternative to conventional chemotherapy for a variety of diseases especially when the host defense mechanism has to be acquired under the conditions of impaired immune responsiveness (4). Indian medicinal plants are rich sources of substances which are claimed to induce para-immunity, the

\* Corresponding author:

E-mail: vaibhav2020@gmail.com

non-specific immunomodulation of especially granulocytes, macro-phages, natural killer cells and competent functions (5).

Among the suppressive synthetic substances, cyclophosphamide has been extensively studied (6). However, the major drawback of this drug is myelosuppression, which is undesirable (7). Immunostimulation and immunosuppression both need to be tackled in order to regulate the normal immunological functioning. Therefore, the stimulatory or suppressive agents have been shown to possess activity to normalize or modulate the pathophysiological processes and are hence called immunomodulatory agents (8).

In recent years, immunostimulatory activity has been reported in a number of Ayurvedic plants like *Terminalia chebula* (9, 10), *Curcuma longa* (11), *Lawsonia alba* (12), and *Tinospora cordifolia* (5, 13-16). *Tc* stem is stomachic, diuretic (17), cures jaundice, skin diseases (18), etc.

*Tinospora cordifolia* (*Tc*) itself possesses a bitter, pungent and astringent taste but it acts through the virtue of its madhur (sweet) vipak. The bitter taste is said to improve metabolic activity, even at a cellular level. All the three tastes destroy toxic metabolic products and purify the tissues. Rasayana drugs with madhur vipak are considered to be immunostimulants (13). The root and stem of *Tc* are prescribed in combination with other drugs as an anti-dote to snake bite and scorpion sting (19), dry barks has anti-spasmodic, antipyretic (20), anti-allergic (21), anti-inflammatory (22), anti-leprotic properties (23), Immunomodulatory effect in human Immuno deficiency virus positive patients (24).

In earlier studies, the immunomodulatory activity of *Tc* was studied through different parameters viz. differential leukocyte count (25), phagocytic activity (26), leukocyte migration inhibition (27), delayed type hypersensitivity (28), bone marrow cellularity (29), etc. Although there are many therapeutic effects, no data is available on the immunomodulatory effect of *Tc* stem (mango plant climber) at cellular level until now. In this context, the objective of the present study was to study the immunomodulator potential of ethanolic extract of *Tc* at cellular and molecular level using male rats.

## Experimental

### *Plant material and preparation of extracts*

The stems of *Tc* (mango plant climber) were identified by Dr G. C. Joshi (Botanist), Ranikhet, Uttarakhand, India. A voucher specimen has been preserved in the herbarium at the Pharmacognosy laboratory under the voucher specimen number of PHG/H/2132 for the future reference. Then, the stems of *Tc* were shade dried and coarsely powdered, extracted in soxhlet apparatus with alcohol for 32 h. The extract was then concentrated to dryness under reduced pressure using rotary evaporator at 45°C and preserved in dessicator for further use.

### *Animals*

Male Wister rats weighing 150-180 g were procured from Laboratory Animals Resources, Division of Animal Genetics, Indian Veterinary Research Institute (IVRI), Izatnager, Bairelly, India, Reg No. CPC-196 and acclimatized to laboratory condition at Animal House, IFTM, Moradabad at room temperature with a 12 h/12 h light/dark cycle and relative humidity (60 ± 5%) and had free access to standard pellet diet; water was provided *ad libitum*. The Institutional Animal Ethical Committee reviewed the animal protocol prior to the experiment. All rats were treated in accordance with the guideline for the Care and Use of Laboratory Animals (NIH Publication No. 86-23, revised 1985) with the permission of Institute Animal Ethical Committee (Proposal No. 11). All experiments were carried out with strict adherence to ethical guidelines and were conducted as approved protocol by the Institutional Animal Ethics Committee (IAEC) and as Indian norms laid down by the Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA), New Delhi; vide approval number CPCSEA/IAEC/837/AC/04.

### *Animal grouping*

For experimental procedure, male rats were divided into four groups of each six as below: Group I: Negative control: Rats treated with 2 mL of 1% gum acacia solution in distilled water. Group II: Positive control: Sensitized rats (through administrating  $1 \times 10^8$  SRBCs, IP)

treated with 1% gum acacia solution orally. Group III: Rats treated with cyclophosphamide 100 mg/Kg/p.o. Group IV: Sensitized rats treated with *Tc* ethanolic extract 100 mg/Kg/p.o.

The administrations of drugs were in the following regimens:

a) Four days prior to the sensitization (days -3, -2, -1, 0)

b) Seven days after the sensitization (days +1, +2, +3, +4, +5, +6, +7)

#### *Preparation of sheep red blood cells (SRBC)*

Blood of healthy sheep was collected from local butcher house and mixed with sterile Alsever's solution (1 : 1). Then, it was thoroughly mixed and centrifuged at 3000 rpm for 5 min. Supernatant was discarded and SRBC pellets were washed twice with sterilized phosphate buffer saline (pH = 7.2). Then, the SRBC pellets were prepared in phosphate buffer saline (pH = 7.2), total SRBC was counted using Neubauer chamber and finally  $1 \times 10^8$  SRBCs (0.5 mL) were injected intraperitoneally for sensitization and challenging the rats (30).

#### *Immunological activity*

The immunostimulating activity was examined through the selection of *in-vivo* and *in-vitro* immunological tests. Six hours after the last dose, the animals were sacrificed and their livers, pineal glands, spleens, kidneys, bladders and adrenal glands were separated and the following parameters were determined.

#### *Estimation of liver mitochondrial antioxidant marker enzymes*

For the isolation of mitochondria, 1 g of liver tissue was weighed and homogenized with 5 mL of 0.35 M sucrose buffer (pH = 7.0) at 4°C and centrifuged at 10,000 g for 5 min. The supernatant was discarded and resultant mitochondrial pellets were suspended in a mixture of 1 mL of 10 mM Tris-HCl (pH = 7.4) solution and 0.2 mL of 1 mM EDTA, finally volume was made up to 2 mL with 0.25 M sucrose solution (31). This solution was used for determination of various mitochondrial antioxidant enzymes.

#### *Estimation of lipid peroxidation (LPO) by homogenization of liver tissue*

After sacrificing the animals, the liver was

isolated, weighed, washed with chilled ice-cold sterile 0.9% NaCl solution. Then, the tissue homogenates were prepared in a ratio of 1 g of wet tissue to 9 mL of 1.15% KCl by using a Teflon Potter-Elvehjem homogenizer (32).

#### *Assay procedure*

A volume of 0.2 mL tissue homogenate was added to the mixture of 0.2 mL of 1% SDS, 1.5 mL of 20% acetic acid solution adjusted to the pH of 3.5 with NaOH, and 1.5 mL of 0.8% aqueous solution of thiobabutaric acid. The mixture was made to 4.0 mL with distilled water and then heated in an oil bath at 95°C for 60 min using a glass ball as a condenser. After cooling in tap water, 1.0 mL of distilled water, and 5.0 mL of the mixture of *n*-butanol and pyridine (15:1 v/v) were added to the mixture and shaken vigorously. Then, it was centrifuged at 4000 rpm for 10 min, the organic layer was taken and the absorbance at 532 nm was measured. 1, 1, 3, 3- tetramethoxypropane (TMP) was used as a standard and the level of LPO was expressed as nmol of malondialdehyde (MDA) (33).

#### *Reduced glutathione (GSH)*

Mitochondrial GSH was estimated through adding 0.2 mL of mitochondrial enzyme solution to 1.8 mL distilled water followed by the addition of 3.0 mL precipitating mixture (0.0501 g metaphosphoric acid, 0.006 g EDTA and 0.9 gm NaCl in 3 mL distilled water). It was centrifuged at 5000 g for 5 min, 1.0 mL of the supernatant was collected and 1.5 mL of the phosphate solution was added to it and then, 0.5 mL of 5,5'-Dithio-Bis (2-nitrobenzoic acid) (DTNB) reagent was added. The optical density was measured at 412 nm spectrophotometrically (Schimadzu 1601). The absorbance of reduced GSH was noted and represented as percentage  $\mu\text{Mol/gm Hb}$  (34).

#### *Catalase (CAT)*

A volume of 0.2 mL mitochondrial enzyme solution was added to a cuvette containing 2.0 mL of phosphate buffer (pH = 7.0) and 1.0 mL of 30 mM  $\text{H}_2\text{O}_2$ . Catalase activity was measured at 240 nm for 1 min using spectrophotometer. The molar extinction coefficient of  $\text{H}_2\text{O}_2$ , 43.6  $\text{M cm}^{-1}$ , was used to determine the catalase activity.

**Table 1.** The sequences of primers used in RT-PCR.

Primer	Sequence (5' to 3')	Length (bases)
<b>IL-10Rat-F</b>	ACCAGCTGGACAACATACTGCTGA	24
<b>IL-10Rat-R</b>	CCTTGATTCTGGGCCATGGTTCT	24
<b>IL-2Rat-F</b>	CTGCAGCGTGTGTTGGATTTGACT	24
<b>IL-2Rat-R</b>	TTGCTGGCTCATCATCGAATTGGC	24
<b>BactRat-F</b>	TGAGAGGGAAATCGTGCCTGACAT	24
<b>BactRat-R</b>	ACCGCTCATTGCCGATAGTGATGA	24
<b>TNF Rat-F</b>	CTGGCCAAATGGCATGGATCTCAA	24
<b>TNF Rat-R</b>	ATGAAATGGCAAATCGGCTGACGG	24
<b>RT-BGF</b>	CATGTTTGTGATGGGCGTGAACCA	24
<b>RT-BGR</b>	TAAGTCCCTCCACGATGCCAAAGT	24

One unit of activity is equal to 1 mmol of H<sub>2</sub>O<sub>2</sub> degraded per minute and is expressed as units/mg of protein (35).

#### *Superoxide dismutase (SOD)*

Fifteen µL of the mitochondrial enzyme solution was added to a mixture of 0.5 mL of 75 mM Tris-HCl buffer (pH = 8.2), 1 mL of 30 mM EDTA and 1 mL of 2 mM pyrogallol. The absorbance was recorded at 420 nm for 3 min using spectrophotometer. One unit of enzyme activity is calculated as the 50% inhibition of pyrogallol autooxidation rate as determined through the change of absorbance/min at 420 nm. The activity of SOD is expressed as units/mg protein (36).

#### *SDS-PAGE*

Protein fractions were isolated from the pineal glands of animals from each group and subjected to the SDS-PAGE analysis for the detection of melatonin secretion. Therefore, the resolving gel mixture and the stacking gel mixtures used were 12% and 5% respectively. The gels were stained using Coomassie Brilliant Blue R-250 solution for 4 h. Further gels were kept in destaining solution for 8 h until the background gets colourless. Finally, after taking the photographs, the gels were stored in distilled water containing 20% glycerol (37).

#### *Lymphocyte proliferation assay*

Splenocyte single cell suspension was prepared through up-downing 4 mL of RPMI-

1640 in spleen and after omitting RBCs using 0.75% NH<sub>4</sub>Cl in Tris buffer (0.02%, pH = 7.2) (adding 6 mL buffer to 2 mL cell suspension after 3 min of centrifugation at 1000 g for 2 min). The concentration was adjusted to 2×10<sup>6</sup> Cells/mL in RPMI-1640 supplemented by 10% fetal calf serum, 2 mM L-Glutamine, 25 mM (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid) (HEPES). A hundred µL of diluted cell suspension was dispensed into a 96-well flat bottom culture plate. Mitogen phytohemmagglutinin A (PHA) was added at 5 µg/mL of final concentration to each well and the volume was adjusted to 0.2 mL. After incubating for 72 h at 37°C and 5% CO<sub>2</sub> humid atmosphere air, the cell proliferation was determined using MTT assay method (37). The 10% of (3-(4, 5-dimethyl-2-thiazolyl) 2,5-diphenyl-2H-tetrazolium) (MTT) (5 mg/mL) was added to each well and the plates were incubated at 37°C in CO<sub>2</sub> humid atmosphere for 4 h. The blue formazan precipitate was dissolved in acidic isopropanol and its optical density was measured at 570 nm using Elisa Reader. The stimulation index was calculated as follows (39):

Stimulation index = Optical density of stimulated cells/Optical density of unstimulated cells.

#### *Real-time PCR for the determination of IL-2, IL-10 and TNF-α expression in spleen cells*

Spleen cell suspensions were prepared and used for the determination of cytokines such as IL-2, IL-10 and TNF-α. The sequences of primers used in RT-PCR are given in Table 1.



**Table 2.** Effect of *Tc* on rat's liver mitochondrial enzymes.

Treatment group	Liver Mitochondrial Enzymes			
	LPO (MMDA/g Hb)	GSH ( $\mu\text{Mol/g Hb}$ )	SOD (units/mg protein)	Catalase (units/mg protein)
Vehicle	83.85 $\pm$ 2.12	4.22 $\pm$ 0.21	22.22 $\pm$ 0.10	271.31 $\pm$ 3.2
SRBC sensitized	98.38 $\pm$ 1.14	2.13 $\pm$ 0.18	19.48 $\pm$ 0.16	252.22 $\pm$ 2.5
Cyclophosphamide	90.42 $\pm$ 2.13	2.31 $\pm$ 0.38	20.38 $\pm$ 0.18	256.29 $\pm$ 3.8
<i>Tc</i>	62.52 $\pm$ 1.13*	4.82 $\pm$ 0.31*	25.32 $\pm$ 0.14*	278.41 $\pm$ 9.2*

\* $p < 0.05$  when compared to SRBC sensitized and cyclophosphamide-treated groups.

#### RNA isolation

Trizol LS reagent (Invitrogen cat No. 10296-010) was used for the RNA isolation and the method prescribed by the manufacturer was followed. Briefly, spleen cells were washed with PBS lysed through adding Trizol LS reagent onto them and repeated pipetting. The organic and aqueous phases were separated with the help of chloroform and centrifugation. Total RNA from the aqueous phase was precipitated with isopropyl alcohol, washed in ethanol, air dried and diluted in nuclease free water.

#### cDNA preparation

Revert Aid M-MuLV Reverse Transcriptase (Fermentas cat No. EP0441) was used for this purpose. About 100 ng of RNA and 0.5  $\mu\text{g}$  of oligo dT (13-19) primer (Invitrogen cat No. 18418-012) were mixed and snap chilled. Four  $\mu\text{L}$  of 5 $\times$  reaction buffer, 20 U of RNase inhibitor, 2  $\mu\text{L}$  of 10 mM dNTP and 200 U of Revert Aid M-MuLV Reverse Transcriptase were added to this mixture. The final reaction volume was kept 20  $\mu\text{L}$ . The reaction mixture was incubated at 42°C for 1 h. The reaction was then stopped through heating the mixture at 70°C for 10 min. The cDNA was divided into aliquots of suitable size and stored at -20°C (40).

#### Real-time PCR

Brilliant SYBR Green QPCR Core Reagent Kit (Stratagene cat No. 600546) was used for this purpose. One  $\mu\text{L}$  of cDNA was added to 11  $\mu\text{L}$  of nuclease free water, 2.5  $\mu\text{L}$  of 10 $\times$ core PCR buffer, 1.25  $\mu\text{L}$  of 50 mM  $\text{MgCl}_2$  solution, 1  $\mu\text{L}$  of 20 mM dNTP mix, 1  $\mu\text{L}$  (12.5 pM/ $\mu\text{L}$ ) of forward primer, 1  $\mu\text{L}$  (12.5 nM/ $\mu\text{L}$ ) of reverse primer, 4  $\mu\text{L}$  of 50% glycerol solution, 0.75  $\mu\text{L}$  of 100% DMSO, 1.25  $\mu\text{L}$  of 1:3000 diluted SYBR Green I dye and 0.25  $\mu\text{L}$  (1.25 U)

of SureStart Taq DNA polymerase. Real-time PCR was carried out in M $\times$ 3000p (Stratagene). The thermal profile used for this was as follows: 95°C for 10 min then 40 cycles of 95°C for 30 sec, 64°C for 30 sec and 72°C for 30 sec with fluorescence recording at the end of each cycle, followed by denaturation of products from 55°C to 95°C with fluorescence recording throughout the step. Fold changes in target transcript levels were determined using reported method (40), considering  $E_{\text{target}} = E_{\text{ref}} = 2$ . The GAPDH gene was used as a reference.

#### Statistical analysis


The results were expressed as mean  $\pm$  SD and the statistical evaluation of the data was done using the prism software (version 5.00) and Student's t-test. P-values less than 0.05 were considered as significant.

## Results and Discussion

#### Liver mitochondrial enzyme estimation

Table 2 shows the increased lipid peroxidation level in liver (98.38  $\pm$  1.14 MMDA/g Hb) in SRBC-treated group as compared to the vehicle-treated rat (83.85  $\pm$  2.12 MMDA/g Hb). Furthermore, cyclophosphamide-treated rats showed slight increase (90.42  $\pm$  2.13 MMDA/g Hb) in the lipid peroxidation level in the liver and after the treatment of *Tc* ethanolic extract, the elevated lipid peroxidation was decreased to 62.52  $\pm$  1.13  $\pm$  1.16 MMDA/g Hb.

The levels of reduced glutathione, superoxide dismutase and catalase were decreased through the SRBC sensitization and were increased via the treatment with *Tc* ethanolic extract. The levels of reduced glutathione, superoxide dismutase and catalase, treated by SRBC sensitization and *Tc* ethanolic extract, were 2.13  $\pm$  0.18



**Figure 1.** SDS-PAGE of rat pineal gland.

Lane 1: Indicates the protein marker of a standard; Lane 2: *Tc* ethanolic extract treatment shows thick band of protein expression (248 kDa) indicating the increased secretion of melatonin via pineal gland due to the immunomodulatory action; Lane 3: Control group; Lane 4: SRBC sensitized group; Lane 5: Immunosuppressant action of cyclophosphamide.

and  $4.82 \pm 0.31$   $\mu\text{Mol/g}$  Hb,  $19.48 \pm 0.16$  and  $25.32 \pm 0.21$  units/mg protein and  $252.22 \pm 2.5$  and  $278.41 \pm 9.2$  units/mg protein respectively. Cyclophosphamide showed immunosuppression through increasing the lipid peroxidation level and decreasing the level of reduced glutathione, superoxide dismutase and catalase as compared with the control group.

The oxidation of membrane's lipid molecules damages the development of several physiological and pathological disorders. The inhibition of lipid peroxidation by any means is the best way to avoid these disorders in the body (41). Present study showed the effect of *Tc* ethanolic extract on lipid peroxidation which was inhibitor as compared with SRBC and cyclophosphamide-treated groups.

Glutathione is the major endogenous antioxidant produced by the cells, participating directly in the neutralization of free radicals and reactive oxygen compounds, as well as maintaining the exogenous antioxidants such as vitamins C and E in their reduced (active) forms. Through the direct conjugation, it detoxifies many xenobiotics (foreign compounds) and carcinogens, both organic and inorganic. It is

essential for the immune system to exert its full potential viz. modulating the antigen presentation to lymphocytes, and thereby, influencing the cytokine production; enhancing the proliferation of lymphocytes thereby increasing magnitude of response; enhancing the killing activity of cytotoxic T-cells and natural killer cells and regulating apoptosis thereby maintaining the control of immune response. Glutathione plays a fundamental role in numerous metabolic and biochemical reactions such as DNA synthesis and repair, protein synthesis, prostaglandin synthesis, amino acid transport and enzyme activation. Thus, every system in the body can be affected through the state of glutathione system, especially the immune system, the nervous system, the gastrointestinal system and the lungs (42-45).

Superoxide dismutase induces the activation of endogenous system of the antioxidant defenses which fights against free radicals (46). It is known that superoxide dismutase plays an important role in the detoxification of superoxide anion and  $\text{H}_2\text{O}_2$ , and thereby promotes the cell against the free radical-induced damage (47).

Catalase is an enzymatic antioxidant and

**Table 3.** Proliferation of rat spleen lymphocytes in response to antigen prepared from *Tc* ethanolic extract using MTT dye reduction test.

Treatment	Stimulation index	Phytohaemagglutinin-stimulated
Vehicle	0.787 ± 0.256	1.222 ± 0.841
Cyclophosphamide	0.832 ± 0.471	0.936 ± 0.412
<i>Tc</i>	1.428 ± 0.361*	1.386 ± 0.361*

\*p < 0.05 when compared with control and cyclophosphamide-treated group.

helps in neutralizing the toxic effect of H<sub>2</sub>O<sub>2</sub>. Hydrogen peroxide is not reactive enough to cause a chain of lipid peroxidation reactions, but its combination with superoxide radical produces hydroxyl radicals that are highly reactive and thus initiates the lipid peroxidation reaction (48). This lipid peroxidation damages the cell membrane resulting into the development of several physiological and pathological disorders. Catalase prevents this through the conversion of hydrogen peroxide to water and non-reactive oxygen species and thus, prevents from the generation of hydroxyl radical and protect the cell from the oxidative damage (49, 50). Our studies showed distinct increase in the level of glutathione, superoxide dismutase and catalase in case of *Tc*-treated groups as compared to SRBC and cyclophosphamide-treated groups. Therefore, *Tc* has antioxidant activity as well as immunomodulator property through protecting the cells from the oxidative damage.

#### Separation of pineal proteins through SDS-PAGE

A thick band at 248 kDa appeared in the group treated with *Tc* ethanolic extract indicating higher secretion of melatonin (Mol. Wt. 248 kDa) (lane 2). In SRBC-sensitized group, the expression of proteins (lane 4) was less than that of control group (lane 3). The immunosuppressant cyclophosphamide showed the least expression of proteins as compared to other groups (lane 5). The increased expression of melatonin in lane 2 confirms the immunomodulator activity of *Tc* (Figure 1).

Melatonin plays an important role as immunomodulatory (51-53) properties, as well as many other physiological functions (54). Melatonin was reported to be an integral part of the immune system, through exerting direct and/or indirect stimulatory effect on both cellular and humoral immunity (51). The present study showed that *Tc* treatment shows the thick band of protein expression indicating

the increased production of the melatonin as compared to SRBC sensitized, vehicle and cyclophosphamide-treated groups. Hence, *Tc* exhibits immunostimulant action through enhancing melatonin secretion in pineal gland.

#### Lymphocyte proliferation assay

The administration of *Tc*-ethanolic extract increased the spleen lymphocytes proliferation (1.428 ± 0.361) as compared to the vehicle-treated group (0.787 ± 0.256) and cyclophosphamide-treated rats (0.832 ± 0.471) (Table 3). Phytohaemagglutinin used as a standard shows proliferation of lymphocytes in all groups.

The earlier works have revealed that drugs having immunomodulatory activity show enhanced proliferation of splenocytes (55). In present study, *Tc* was found to significantly enhance the proliferation of splenocytes as compared with control and cyclophosphamide treatment. The proliferation of lymphocytes indicated the increase in the number of B and T cells, which release cytokines and growth factors that regulate other immune cells and secretion of antibodies in the blood (56).

#### Comparison of IL-2, IL-10 and TNF- $\alpha$ gene expression in rat spleen mRNA

Measuring and assessing cytokine profile provides a useful method for the accurate study of cytokine such as IL-2, IL-10 and TNF- $\alpha$  and immunomodulatory effect was investigated through RT-PCR. *Tc* treatment showed more fold changes in IL-2, IL-10 and TNF- $\alpha$  mRNA (17.63, 42.52 and 5.39 respectively) levels as compared to the vehicle, cyclophosphamide and SRBC sensitized groups as shown in Table 4. Increased expression of fold change in different mRNA levels using *Tc* confirms the immunostimulant action.

Cytokines are released by living cells of the

**Table 4.** Effect of different treatments on IL-2, IL-10 and TNF- $\alpha$  gene expression.

Treatment	Fold change in IL-2, IL-10 and TNF- $\alpha$ mRNA level		
	IL-2	IL-10	TNF- $\alpha$
Vehicle	1.00	1.00	1.00
Cyclophosphamide	6.41	8.63	-1.78
<i>Tc</i>	17.63	42.52	5.39
SRBC sensitized	-2.87	22.63	1.18

host in a highly regulated fashion to regulate the cell functions via specific receptors (57) which participate in the control of all immunologically relevant events, whether they concern the activation, differentiation, maturation, proliferation, apoptosis, or acquisition of effectors' functions. Cytokines influence the quantitative, as well as the qualitative outcome of all the immune responses (58). In our study, after the administration of ethanolic extract of *Tc* in male Wister rats, significant immunomodulatory activities were shown through fold changes in IL-2, IL-10 and TNF- $\alpha$  mRNA levels. IL-2 shows immunomodulatory activities by T and B lymphocytes proliferation, natural killer cell activation (59, 60) and also increasing in the immunoglobulin G production whereas IL-10 promotes elevation of Th<sub>2</sub> cells (58, 60) and TNF- $\alpha$  modulate the cytokine gene expression (59). Hence, the immunomodulatory activity of *Tc* may be due to the above mechanism(s).

### Conclusion

From the above results, it can be concluded that *Tc* stem (mango plant climber) has potent immunomodulatory activity and the effect may be due to the specific chemical constituents or the synergistic effect of chemical compounds of the drug.

### Acknowledgments

Authors are thankful to Dr. Sameer Shrivastava, Scientist and Dr. Sonal Shrivastava, Scientist, Peptide Synthesis and Protein Engineering Lab, Division of Animal Biotechnology, Dr. Abhijit Pawde, Scientist, Division of Surgery, Dr. Arvind Sonawane, Scientist, Division of Genetics, Indian Veterinary

Research Institute (IVRI), Izatnagar, Bareilly, India for their cooperation.

### References

- (1) Dahanukar SA, Kulkarni RA and Rege NN. Pharmacology of medicinal plants and natural products. *Indian J. Pharmacol.* (2000) 32: S81-S118.
- (2) Liu Y, Jiao F, Qiu Y, Li W, Lao F, Zhou GQ, Sun BY, Xing GM, Dong JQ, Zhao YL, Chai ZF and Chen CY. The effect of Gd@C<sub>82</sub>(OH)<sub>22</sub> nanoparticles on the release of Th1/Th2 cytokines and induction of TNF- $\alpha$  mediated cellular immunity. *Biomaterials* (2009) 30: 3934-3945.
- (3) Sharma P. *Charaka Samhita, Chikitsasthana*. Varanasi Chaukhamba Orientalia, India (1983) 2.
- (4) Lily G, Padwad Y, Singh R, Karan D, Chanda S, Chopra MK, Bhatnagar P, Kashyap R and Sawhney RC. Antiinflammatory activity of seabuckthorn (*Hippophae rhamnoides*) leaves. *Int. J. Immunopharmacol.* (2005) 5: 1675-1684.
- (5) Sainis KB, Sumariwalla PF, Goel A, Chintalwar GJ, Sipa- himalani AT and Banerji A. Immunomodulatory properties of stem extracts of *Tinospora cordifolia*: cell targets and active principles. In: Upadhyay SN. (ed.) *Immunomodulation*. Narosa Publishing House, New Delhi (1997) 95.
- (6) Shand FL and Howard JG. Induction *in-vitro* of reversible immunosuppression and inhibition of B cell receptor regeneration by defined metabolites of cyclophosphamide. *Eur. J. Immunol.* (1979) 9: 17-21.
- (7) Hardman JG, Limbird LE, Molinoff PB, Ruddon RW and Gilman AG. Alkylating agents. In: New York Academy of Sciences (eds.) *Goodman and Gilman's the Pharmacological Basis of Therapeutics*. McGraw-Hill, New York (1996) 1216-1218.
- (8) Wagner H. Immunomodulatory agents. *Proc. Alfred Benzon Symp.* (1983) 20: 559.
- (9) Chattopadhyay RR and Bhattacharyya SK. *Terminalia chebula*: an update pharmacognosy reviews. *Phcog. Rev.: Plant Review* (2007) 1: 151-157.
- (10) Prakash SS, Niranjana Patra CH, Santanu CH, Hemant Kumar P, Jagannath Patro V and Vimala Devi V. Studies on flowability, compressibility and *in-vitro* release of *Terminalia chebula* fruit powder tablets. *Iranian J. Pharm. Res.* (2011) 10: 393-401.
- (11) Deodhar SD, Sethi R and Srimal RC. Preliminary



- study on antirheumatic activity of curcumin (diferuloyl methane). *Ind. J. Med. Res.* (1980) 71: 632-634.
- (12) Dev S. Ethanotherapeutics and modern drugs development: The potential of Ayurveda. *Curr. Sci.* (1997) 73: 909-928.
- (13) Dahanukar SA, Thatte UM and Rege NN. Immunostimulants in Ayurveda medicine. In: Wagner H. (ed.) *Immunomodulatory Agents from Plants*. Berkhauer Verlag, Basel (1999) 289-323.
- (14) Thatte UM, Rao SG and Dhanukar SA. *Tinospora cordifolia* induces colony stimulating activity in serum. *J. Postgrad. Med.* (1994) 40: 202-203.
- (15) Sainis KB, Ramakrishnan R, Sumariwalla PF, Goel A, Sipahimalani AT and Banerji A. Further studies on immunomodulation by natural products from *Tinospora cordifolia*. In: Upadhyay SN. (ed.) *Immunopharmacology: Strategies for Immunotherapy*. Narosa Publishing House, India (1999) 96-104.
- (16) Aher Vaibhav D and Wahi Arunkumar. Pharmacological study of *Tinospora cordifolia* as an immunomodulator. *Int. J. Curr. Pharma. Rev.* (2010) 2: 52-54.
- (17) Nayampalli SS, Ainapure SS, Samant BD, Kudtarkar RG, Desai NK and Gupta KC. A comparative study of diuretic effects of *Tinospora cordifolia* and hydrochloro-thiazide in rats and a preliminary phase I study in human volunteers. *J. Postgrad. Med.* (1988) 34: 233-236
- (18) Raghunathan K and Mittra R. (eds.) *Pharmacognosy of Indigenous Drugs*. Central Council for Research in Ayurveda and Siddha, New Delhi (1982) 484-510.
- (19) Zhao TF, Wang X, Rimando AM and Che C. Folkloric medicinal plants: *Tinospora sagittata* var. *cravaniana* and *Mahonia bealei*. *Planta Med.* (1991) 57: 505.
- (20) Ikram M, Khattak SG and Gilani SN. Antipyretic studies on some indigenous Pakistani medicinal plants: II. *J. Ethnopharmacol.* (1987) 19: 185-192.
- (21) Nayampalli SS, Desai NK and Ainapure SS. Anti-allergic properties of *Tinospora cordifolia* in animal models. *Indian J. Pharm.* (1986) 18: 250-252.
- (22) Rai M and Gupta SS. The deposition of the secondary salts over the five pellets in rats was inhibited by the aqueous extract of *T. cordifolia*. *J. Res. Ind. Med.* (1966) 10: 113-116.
- (23) Asthana JG, Jain S, Mishra A and Vijaykanth MS. Evaluation of antileprotic herbal drug combinations and their combination with Dapsone. *Indian Drugs* (2001) 38: 82-86.
- (24) Thawani VR, Kalikar MV, Varadpande VK, Sontakke SD, Singh RP and Khiyani RK. Immunomodulatory effect on *T. cordifolia* in human Immunodeficiency virus positive patients. *Indian J. Pharmacol.* (2008) 40: 107-110.
- (25) Sheela D, Ramasundaram S and Jaya PN. Immunomodulatory activity of triphala on neutrophil functions. *Biol. Pharm. Bull.* (2005) 28: 1398-1403.
- (26) Furthvan R and Bergvanden BM. *Clinical Immunology*. Gower Medical Publishing, London (1991) 67.
- (27) Sorberg M, Bendisen G. Human leukocyte migration as a parameter of hypersensitivity. *Acta Med. Scand.* (1967) 18: 247-252.
- (28) Benencia F, Courreges M, Cecialia C and Felix C. *In-vivo* and *in-vitro* immunomodulatory activities of *Trichilia glabra* aqueous leaf extracts. *J. Ethnopharmacol.* (2000) 69: 199-205.
- (29) Sredni B, Albeck M, Kazimirsky G and Shalet F. Immunomodulator AS101 administered orally as a chemoprotective and radioprotective agent. *Int. J. Immunopharmacol.* (1992) 14: 613-619.
- (30) Patel S, Basnji D, Otilia JF, Patel MM and Shah KK. Scrutinizing the role of aqueous extract of *Trapa bispinosa* as an immunomodulator in experimental animals. *Int. J. Res. Pharm. Sci.* (2010) 1: 13-19.
- (31) Jonshon D and Lardy H. Isolation of liver, kidney mitochondria. *Methods in Enzymology* (1947) 10: 94-96.
- (32) Hogeboom GH. Fractionation of cell components of animal tissues. In: Colowick SP and Kaplan NO. (eds.) *Methods in Enzymology*. Academic Press, New York (1955) 16-19.
- (33) Ohkawa H, Ohishi N and Yagi K. Assay for lipid peroxidation in animal tissue by Thiobarbituric acid reaction. *Anal. Biochem.* (1979) 95: 351-358.
- (34) Ellman GL. Tissue sulphhydryl group elucidation of the biologically active components of *Terminalia chebula* Retzius. *Arch. Biochem. Biophys.* (1959) 82: 70-77.
- (35) Aebi H. Catalase *in-vitro*. *Methods Enzymol.* (1984) 105: 121-126.
- (36) Marklund S and Marklund C. Involvement of the superoxide anion radical in the autooxidation of pyrogallol and a convenient assay for superoxide dismutase. *Eur. J. Biochem.* (1984) 7: 469-474.
- (37) Sambrook J and Russell D. *Molecular Cloning. A Laboratory Manual*. 3<sup>rd</sup> ed. (2000) 3, A8: 40-49.
- (38) Mosmann TR and Coffman RL. Th1 and Th2 cells. Different pattern of lymphokines secretion lead to different functional properties. *Ann. Rev. Immunol.* (1989) 7: 145-173.
- (39) Singh GK, Kumar P, Chauhan RS and Singhal LK. Cow urine upregulates lymphoblastogenesis in chicks. *ISAH. Warsaw, Poland* (2005) 2: 90-92.
- (40) Pfaffi MW. A new mathematical model for relative quantification in Real-Time RT-PCR. *Nucleic Acids Res.* (2001) 29: 2003-2007.
- (41) Janero DR. Malonyldialdehyde and thiobarbituric acid reactivity as diagnostic indices of lipids peroxidation and peroxidation tissue injury. *Free Radic. Biol. Med.* (1990) 9: 515-540.
- (42) Kinschereff R. Effect of glutathione depletion and oral *n*-acetyl-cysteine treatment on CD4<sup>+</sup> and CD8<sup>+</sup> cells. *FASEB J.* (1994) 8: 448-451.
- (43) Wu D. *In-vitro* glutathione supplementation enhances interleukin-2 production and mitogenic response of peripheral blood mononuclear cells from young and old subjects. *J. Nutr.* (1994) 124: 655-663.
- (44) Fidelus M. Glutathione and lymphocyte activation: a function of aging and auto immune disease. *Immunol.* (1987) 61:503-508.
- (45) Buhl R. Systemic glutathione deficiency in symptom free HIV-1 seropositive individuals. *Lancet* (1989)

- 334: 1294-1298.
- (46) Bourg M and Joanny F. Superoxide dismutase, a powerful antioxidant, is now available orally. *Phytotherapie* (2005) 3: 1-4.
- (47) Goyal BR, Mahajan S, Mali R, Ramesh GK and Anita MA. Beneficial effect of *Achyranthes aspers* Linn. in toluene di-isocyanate induced occupational asthma in rats. *Global J. Pharmacol.* (2007) 1: 6-12.
- (48) Lubac B, Hayr M, Denk W and Bauer G. Brain lipid peroxidation and hydroxyl radical attack following the intravenous infusion of hydrogen peroxide in an infant. *Free Radic. Biol. Med.* (1996) 21: 219-223.
- (49) Chelikani P, Fita I and Loewen P. Diversity of structures and properties among catalases. *Cell Mol. Life Sci.* (2004) 61: 192-208.
- (50) Zamocky M and Koller F. Understanding the structure and function of catalases: clues from molecular evolution and *in-vitro* mutagenesis. *Prog. Biophys. Mol. Biol.* (1999) 72: 19-66.
- (51) Guerrero JM and Reiter RJ. Melatonin-immune system relationships. *Curr. Top. Med. Chem.* (2002) 2: 167-179.
- (52) Carrillo-Vico A, Garcia-Maurino S, Calvo JR and Guerrero JM. Melatonin counteracts the inhibitory effect of PGE<sub>2</sub> on IL-2 production in human lymphocytes via its mt1 membrane receptor. *FASEB J.* (2003) 17: 755-757.
- (53) Carrillo-Vico A, Lardone PJ, Fernandez-Santos JM, Martin-Lacave I, Calvo JR, Karasek M and Guerrero JM. Human lymphocyte-synthesized melatonin is involved in the regulation of the interleukin-2/interleukin-2 receptor system. *J. Clin. Endocrinol. Metab.* (2005) 90: 992-1000.
- (54) Sallinen P, Saarela S, Ilves M, Vakkuri O and Leppaluoto J. The expression of MT1 and MT2 melatonin receptor mRNA in several rat tissues. *Life Sci.* (2005) 76: 1123-1134.
- (55) Kijjoa A, Amat N, Upur H, Ablimit A and Matsidik A. Immunomodulatory effect of Abnormal Savda Munsiq, a traditional Uighur medicine, on the combined stress mice. *J. Ethnopharmacol.* (2009) 122: 42-47.
- (56) Berrington JE, Barge D, Fenton AC, Can AJ and Spickett GP. Lymphocyte subsets in term and significantly preterm UK infants in the first year of life analysed by single platform flow cytometry. *Clin. Exp. Immunol.* (2005) 140: 289-292.
- (57) Kroemer G, Alboran M, Gonzalo JA, and Martinez C. Immunoregulation by cytokines. *Crit. Rev. Immunol.* (1993) 19: 163-191.
- (58) Kresina F. *Thomas Immune Modulating Agents*. Marcel Dekker, New York (1998) 1-19.
- (59) Kroemer GJ, Anderu-Sanchez L, Gonzalo JA, Gutierrez-Ramos JC and Martinez AC. Interleukin-2 autotolerance and autoimmunity. *Adv. Immunol.* (1991) 150: 147-235.
- (60) Plaut M, Pierce JH, Watson CJ, Hanley-Hyde Nordan RP and Paul WE. Mast cell lines produce lymphokines in response to cross-linkage of FcRI or to calcium ionophores. *Nature* (1989) 339: 64.
- 
- This article is available online at <http://www.ijpr.ir>