

Possible Mechanism(s) of the Relaxant Effects of *Achillea wilhelmsii* on Guinea-Pig Tracheal Chains

Mohammad Hossein Boskabady*, Naeima Eftekhari and Mahsa Kaveh

Applied Physiology Research Center and Dept. of Physiology, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran.

Abstract

Achillea wilhelmsii have been used in folk remedies. The relaxant effects of the extract of *A. wilhelmsii* on tracheal chains of guinea pigs were examined. The relaxant effects of four cumulative concentrations of the extract, theophylline and saline were examined by their relaxant effects on precontracted tracheal chains of guinea pig by KCl (group 1), 10 μ M methacholine (group 2), incubated tissues by atropine, propranolol and chlorpheniramine and contracted by KCl (group 3) and incubated tissues by propranolol and chlorpheniramine and contracted by methacholine (group 4). In group 1 and 2, all concentrations of theophylline and three higher concentrations (4, 6 and 8 mg/mL) of the extract showed significant relaxant effects compared to that of saline. In groups 3 and 4 experiments also all concentrations of the extract showed significant relaxant effects compared to that of saline. The relaxant effect of three higher concentrations (4, 6 and 8 mg/mL) of the extract in group 1 were significantly greater than those of group 2 and in group 3 were significantly lower than those of group 1. There were significant positive correlations between the relaxant effects and concentrations for theophylline in groups 1 and 2 and the extract in all four groups of experiments. These results showed a potent relaxant effect for the extract from *A. wilhelmsii* on tracheal chains of guinea pigs. A muscarinic receptor blockade was also suggested for the extract.

Keywords: *Achillea wilhelmsii*; Hydroalcoholic extract; Possible mechanism; Relaxant effects; Guinea pig; Trachea.

Introduction

The herb *Achillea*, which belongs to the family Compositae (Asteraceae), is a genus with more than 100 species all around the world (1). These plants are medicinal perennial rhizomatous herbs that are native to Europe and Western Asia, although they are also found in Australia, New Zealand and North America (2, 3). *Achillea millefolium* C. Koch is the best known species,

commonly named yarrow, and was used as traditional medicine in Europe and by native Americans for treating swollen tissues and wounds (4-6).

The aerial parts of different species of the genus are widely used in folk medicine due to numerous medicinal properties, such as anti-inflammatory, antispasmodic, antihemorrhoidal, stomachic and antiseptic (7-9).

The main components of the oil of *A. wilhelmsii* are carvacrol (25.1%), linalool (11.0%), 1,8-cineol (10.3%), E-nerolidol (9.0%) and borneol (6.4%) (10). The main components of the monoterpene fraction of the

* Corresponding author:

E-mail: boskabady@ums.ac.ir

oil are camphor, borneol, linalool, 1,8-cineole, chrysanthenol acetate and carvacrol (9).

The analgesic (11) and anticonvulsant (12) effects of the plant have been demonstrated. Several investigators (13, 14) reported significant effects from the extracts of *A. wilhelmsii* or *A. talagonica* grown in Iran as being antilipidemic, antihypertensive or immunosuppressive for humans or laboratory animals respectively. The relative radical scavenging activity of the plant is also documented (15).

The antispasmodic activity of total extract of *A. nobilis* subsp. *sipylea* on rat duodenum was observed (16). Some flavonoids can act as spasmolytic agents by relaxing various smooth muscles (17, 18). They have also reduced the tone of guinea-pig isolated trachea, main pulmonary artery, rat uterus and rat vas deferens (19-21). It has been also suggested that the inhibitory effects of cirsiol on smooth muscle are attributed to inhibition of transmembrane Ca^{2+} influx (22). In addition, the relaxant effect of carvacrol on tracheal smooth muscle was observed in our previous study (23).

Therefore in the present study, the possible mechanism(s) of relaxant effects of hydro alcoholic extract from *A. wilhelmsii* on tracheal chains of guinea pigs were examined.

Experimental

Plant and extraction

Achillea wilhelmsii was collected from north east region of Iran and identified by MR Joharghi. A voucher specimen was preserved in the Herbarium of the School of Agriculture, Ferdowsi University (Herbarium No: 40377, FUMH). The plant was dried in the absence of sun light. The hydro-alcoholic extract of the plant was prepared as follows: fifty grams of *Achillea millefolium* were grinded, added to 700 mL of ethanol 50% (350 mL of pure ethanol was diluted with water to final volume of 700 mL) using the Soxhlet apparatus. The solvent was then removed under reduced pressure. The plant ingredient concentration in the final extract was adjusted to 0.1 g/mL by adding distilled water to the dried extract. Three and half (3.5) grams of extract was collected from 50 grams of the grinded plant.

Tissue preparation

Guinea pigs (400-700 g, both sexes) were killed by a blow on the neck and tracheas were removed. Each trachea was cut into 10 rings (each containing 2-3 cartilaginous rings). The study was approved by Ethical Committee of Mashhad University of Medical Sciences. Before experiments, animals were housed in individual cages with access to food and water ad libitum and were maintained at $22^{\circ} \pm 2^{\circ}C$ on a 12 h light/dark cycle (light period 0700 and 1900 h). All the rings were then cut open opposite the trachealis muscle, and sutured together to form a tracheal chain (24, 25). Tissue was then suspended in a 10 mL organ bath (organ bath 61300, Bio Science Palmer-Washington, Sheerness, Kent U.K.) containing Krebs-Henseliet solution of the following composition (mM): NaCl 120, $NaHCO_3$ 25, $MgSO_4$ 0.5, KH_2PO_4 1.2, KCl 4.72, $CaCl_2$ 2.5 and dextrose 11.

The Krebs solution was maintained at $37^{\circ}C$ and gassed with 95% O_2 and 5% CO_2 . Tissue was suspended under an isotonic tension of 1 g and allowed to equilibrate for at least 1 h while it was washed with Krebs solution every 15 min.

Protocols

The relaxant effects of four cumulative concentrations of the extract of *A. wilhelmsii* (2, 4, 6 and 8 mg/mL) and theophylline anhydrous (Sigma Chemical Ltd UK) (0.2, 0.4, 0.6 and 0.8 mM) as positive control, and saline (1 mL) as negative control were examined. To produce the first concentration of the extract, 0.2 mL of 10 g% was added to a 10 mL organ bath and for other three concentrations; 0.2 mL of 10 g% was added to organ bath three times respectively. For theophylline, 0.2 mL of 0.01 M concentrated solution was added to organ bath 4 times. The consecutive volumes were added to organ bath at five minutes intervals.

In each experiment, the effect of saline, four cumulative concentrations from the extract and theophylline on contracted tracheal smooth muscle was measured after exposing tissue to each concentration of the solution for 5 min. A decrease in tone was considered to be a relaxant (bronchodilatory) effect and expressed as positive percentage change in proportion to the maximum contraction. An increase in tone was considered

Table 1. Relaxant effect of the extract of *Achillea millefolium* (percentage change in proportion to the maximum contraction) in comparison with negative control (saline) and positive control (theophylline) in group 1 experiments (contracted tracheal chains with 60 mM KCl, n = 7)

Different Concentration	Saline	Theo.	St. Dif. vs Saline	Extract	St. Dif. vs Saline	St. Dif. vs Theo.
1	-	28.25 ± 3.24	p < 0.001	0.24 ± 7.29	NS	p < 0.001
2	-	49.04 ± 4.19	p < 0.001	42.32 ± 6.84	p < 0.001	NS
3	-	75.62 ± 5.31	p < 0.001	80.3 ± 11.13	p < 0.001	NS
4	0.43 ± 0.2	88.33 ± 8.16	p < 0.001	108.71 ± 13.71	p < 0.001	NS

Values are presented as mean±SEM. St. Dif.: Statistical deference. The four different concentrations for the extract were 2, 4, 6 and 8 mg/mL, and for theophylline, 0.2, 0.4, 0.6 and 0.8 mM. Theo.: theophylline.

as a contractile (bronchoconstrictory) effect which was expressed as negative percentage change (24, 25).

The relaxant effect of different solutions was tested with four different experimental designs, as follows:

I: On tracheal chains contracted by 60 mM KCl (group 1 experiment, n= 7).

II: On tracheal chains contracted by 10 µM methacholine hydrochloride (Sigma Chemical Ltd UK), (group 2 experiment, n= 9).

III: On tracheal chains incubated with 1 µM atropine, propranolol and chlorpheniramine and contracted by 60 mM KCl (group 3 experiment, n= 7).

IV: On tracheal chains incubated with 1 µM propranolol and chlorpheniramine and contracted by 10 µM methacholine (group 4 experiment, n= 8).

The relaxant effects in four groups of experiments were examined in four different series of tracheal chains. All of the experiments were performed randomly with a 1 h resting period of tracheal chains between each two experiments while washing the tissues every

15 min with Krebs solution. In all experiments contractions were measured using an isotonic transducer (Harvard APP LTD, 50-6360 SINO. 0210) and measured by using a software by a computer)Acer model NO.: G781) recording.

Statistical analysis

All data were expressed as mean±SEM. The data of relaxant effects obtained in four groups of experiments were compared using one way analysis of variance (ANOVA) and Tukey-Kramer post test. The relaxant effect of the extract and theophylline were related to the concentrations using least square regression. Significance was accepted at p < 0.05.

Results

Relaxant effect

In group 1 (non-incubated tissues contracted by KCl) and 2 (non-incubated tissues contracted by methacholine), all concentrations of theophylline and three higher concentrations (4, 6 and 8 mg/mL) of the extract showed relaxant effects compared to that of saline (p < 0.001 for

Table 2. Relaxant effect of the extract of *Achillea millefolium* (percentage change in proportion to the maximum contraction) in comparison with negative control (saline) and positive control (theophylline) in group 2 experiments (contracted tracheal chains by 10 µM methacholie, n = 9)

Different Concentration	Saline	Theo.	St. Dif. vs Saline	Extract	St. Dif. vs Saline	St. Dif. vs Theo.
1	-	30.33 ± 3.26	p < 0.001	0.58 ± 3.28	NS	p < 0.001
2	-	48.24 ± 4.93	p < 0.001	22.6 ± 2.55	p < 0.001	p < 0.001
3	-	68.02 ± 5.08	p < 0.001	49.27 ± 5.94	p < 0.001	p < 0.05
4	0.6 ± 0.23	90.86 ± 6.01	p < 0.001	87.98 ± 8.60	p < 0.001	NS

For abbreviations see Table 1.

Table 3. Relaxant effect of the extract of *Achillea millefolium* (percentage change in proportion to the maximum contraction) in comparison with negative control (saline) and positive control (theophylline) in group 3 experiments (contracted tracheal chains by 60 mM KCl-Inc, n = 7) and group 4 experiments (contracted tracheal chains by 10 μ M methacholine-Inc, n = 8)

Different Concentration	Saline	KCl-Inc	St. Dif. vs Saline	M-Inc	St. Dif. vs Saline	St. Dif. KCl-Inc vs M-Inc
1	-	6.47 \pm 1.64	p < 0.01	7.84 \pm 1.81	p < 0.01	NS
2	-	12.54 \pm 3.12	p < 0.01	25.75 \pm 3.92	p < 0.001	p < 0.05
3	-	32.00 \pm 6.04	p < 0.005	60.27 \pm 8.62	p < 0.001	p < 0.05
4	0.72 \pm 0.18	50.17 \pm 6.11	p < 0.001	79.18 \pm 6.91	p < 0.001	p < 0.01

For abbreviations see Table 1.

all cases) (Tables 1 and 2).

In groups 3 and 4 experiments also all concentrations of the extract showed relaxant effects compared to that of saline (p < 0.01 to p < 0.001; Table 3).

Comparison of the relaxant effect of theophylline with that of the extract

In groups 1, the relaxant effect of only lowest concentration (2 mg/mL) and in group 2 the relaxant effect of three lower concentrations (2, 4 and 6 mg/mL) of the extract were significantly less than those of theophylline (p < 0.05 to p < 0.001; Tables 1 and 2).

Comparison of the relaxant effect between group 1 and 2

The relaxant effect of three higher concentrations (4, 6 and 8 mg/mL) of the extract in group 1 (contracted tracheal muscle with KCl) were greater than those of group 2 (contracted tracheal muscle with methacholine), (p < 0.01 to p < 0.001). However, the relaxant effect of theophylline between two groups was not statistically significant (Figure 1).

Comparison of the relaxant effect between incubated and non-incubated tissues

The relaxant effect of three higher concentrations (4, 6 and 8 mg/mL) of the extract in group 3 (tracheal muscle incubated with atropine, propranolol and chlorpheniramine and contracted with KCl) were significantly lower than those of non incubated tissues contracted by KCl (p < 0.05 to p < 0.001). However there was no significant difference in relaxant effect between non incubated (group 2) and incubated

(incubated tissues with propranolol and chlorpheniramine, group 4) tissues contracted by methacholine (Figure 2b).

Comparison of the relaxant effect between incubated tissues contracted by KCl and methacholine

The relaxant effect of three higher concentrations (4, 6 and 8 mg/mL) of the extract in group 3 (tracheal muscle incubated with atropine, propranolol and chlorpheniramine) contracted with KCl were lower than group 4 (incubated tissues with propranolol and chlorpheniramine) contracted by methacholine; (p < 0.05 to p < 0.01; Table 3).

Correlation between concentrations of solutions and their relaxant effect

There were significant positive correlations between the relaxant effects and concentrations for theophylline in groups 1 and 2 and the extract in all four groups of experiments (p < 0.001 for all cases; Table 4).

Discussion

In this study, the relaxant effect of the extract from *A. wilhelmsii* in comparison with saline as negative control and theophylline as positive control was studied. In group 1 (contracted tracheal chains with KCl), the extract and theophylline showed relaxant effect on tracheal smooth muscle. The relaxant effect of only lower concentration of the extract was significantly less than that of theophylline.

The results of this study confirmed those of the study of Asgari *et al.* (14) indicating the




Figure 1. Concentration response curves of the relaxant effect of theophylline (a) and the extract of *Achillea millefolium* (b) in four groups of experiments. group 1; KCl induced contraction of tracheal chains (○, n = 7), group 2; methacholine induced contraction of tracheal chains (□, n = 9), group 3; KCl induced contraction of incubated tracheal chains with atropine, propranolol and chlorpheniramine (▲, n = 7) and group 4; methacholine induced contraction of incubated tracheal chains with propranolol and chlorpheniramine (◆, n = 8). Statistical differences in the relaxant effect of different concentrations of each solution between group 1 and 2; ns: non-significant difference, +; p < 0.05, +++; p < 0.001. Statistical differences in the relaxant effect of different concentrations of the extract between group 1 and 3; ns, non significant difference, ***; p < 0.001. Statistical differences in the relaxant effect of different concentrations of the extract between group 2 with those of group 4; NS, non significant difference. The concentration unit for the extract was mg/mL and for theophylline, mM.

antihypertensive effect of the plant which is perhaps due to its relaxant effect on vascular smooth muscle.

In group 2 (contracted tracheal chains with methacholine), the extract and theophylline showed relaxant effect on tracheal smooth muscle. The relaxant effects of three lower concentrations of the extract were significantly less than that of theophylline. In addition, the relaxant effects of three higher concentrations of the extract in this group were significantly lower than those of group 1.

The greater relaxant effect of the extract on KCl induced contraction of tracheal muscle (group 1), indicates that the extract may have inhibitory effect on calcium channels and/or opening effect on potassium channels. In fact, the relaxant effect of potassium channels openers (26) and calcium channel blockers (27) has been shown. In addition, it is well known that KCl can affect calcium channels. However, the greater relaxant effect on KCl induced contraction (group 1) compared to methacholine induced contraction (group 2) is in favour of inhibitory effect of the extract on calcium channels. In fact, these results supported by those of previous studies suggesting that the inhibitory effects of cirsiolol, a flavone isolated from *Achillea fragrantissima*, on smooth muscle are attributed to inhibition of transmembrane Ca^{2+} influx (22).

It is known that the agonists of β -adrenergic receptors are the most potent relaxant agents of tracheal smooth muscle (28). However, no difference in the relaxant effect of the extract between non incubated (group 2) and incubated tracheal smooth muscle with propranolol and chlorpheniramine (group 4) and contracted with methacholine indicates that the extract has no stimulatory effect on β -adrenergic receptors. These results also suggest that the relaxant effect is not mediated by the inhibitory effect of the extract on histamine (H1) receptors.

However, the significant lower relaxant effect on incubated tracheal smooth muscle with propranolol, chlorpheniramine and atropine and contracted with KCl (group 3) compared to non incubated tissues (group 1) indicated a muscarinic inhibitory effect for the extract. In addition, the relaxant effects of muscarinic receptors inhibitory agents have

Table 4. Correlation (r) between the relaxant effects of the extract of *Achillea millefolium* and theophylline with their concentrations in four groups of experiments.

Different Solutions	KCl		Metha		KCl-Inc		Meth-Inc	
	R	p-value	R	p-value	R	p-value	r	p--value
Extract	0.8507	p< 0.001	0.891	p< 0.001	0.664	P= 0.001	0.735	p< 0.001
Theo.	0.8396	p< 0.001	0.850	p< 0.001				

been documented previously (29). However, more studies are required to examine the exact mechanism(s) of the relaxant effect seen for the extract of the plant.

The extract contained several substances, which could be lead to relaxation response by several mechanisms. Therefore, in group 3, tracheal smooth muscle were incubated by atropine, chlorpheniramine and propranolol to inhibit muscarinic, histamine (H1) and adrenergic receptors and evaluate the possible contribution of these three sets of receptors in observed relaxation response. In this group a significant reduction in relaxation response was observed. In group 4, tracheal smooth muscle was incubated with chlorpheniramine and propranolol to inhibit histamine (H1) and adrenergic receptors and leave muscrinic receptor intact. In this group the relaxation response did not changed significantly which suggest that main mechanism of the relaxant effect of the extract is its blocking effect on muscarinic receptors. However, the exact mechanism(s) of the relaxant effect of the plant should be investigated in further studies.

One of the constituent of *Achillea wilhelmsii* is carvacrol which its relaxant effect was shown previously (23, 30). Therefore, the relaxant effect of the plant could be due to its main constituent, carvacrol. However, the relaxant effect of carvacrol was higher on methacholine induced contraction of tracheal smooth muscle compared to KCl induced contraction but the effect of the extract of *Achillea wilhelmsii* was more potent on KCl induced contraction. These results indicate that the mechanism(s) of the relaxant effect of the extract is not exactly the same as that of carvacrol. Therefore, the relaxant of the extract of *Achillea wilhelmsii* is not just due to its constituent carvacrol.

There were positive correlations between concentrations and the relaxant effects of

both theophylline and the extract indicating a concentration dependent relaxant effect for the extract as well as for theophylline. Although the relaxation effect of the extract of other species of this plants family was shown in different types of smooth muscle (19-21), the relaxant effect of *A. wilhelmsii* on tracheal smooth muscle which is in some way different from other smooth muscle had been studied and this is the first report in this regard. The relaxant effects of the extract observed in the present study can be translated to bronchodilation and might have potential benefits in asthmatic patients.

In conclusion the results of this study indicated a potent relaxant effect for the extract from *A. wilhelmsii* on tracheal chains of guinea pigs. The results also suggest that the H1 histamine receptors and Beta-adrenergic receptors are not involved in the relaxant effect of the extract.

Acknowledgment

This study was financially supported by the Research Council of Mashhad University of Medical Sciences. This paper is the results of a MSc thesis.

References

- (1) Gaji M. In: Josifovi M Ed. Flore de la Republique Socialiste de Serbie. *Academie serbe des sciences et des arts, Belgrade* (1975) 7: 95-96.
- (2) Chevallier A. The Encyclopedia of Medicinal Plants. London: Dorling Kindersley Publishing Inc. (1996) p 54.
- (3) Könemann. Botanica: The Illustrated A-Z of over 10,000 Garden Plants and How to Cultivate Them. Gordon Cheers Publication, Hong Kong, 1999 pp. 51-53.
- (4) Chandler RF, Hooper SN and Harvey MJ. Ethnobotany and phytochemistry of yarrow, *Achillea millefolium*, Compositae. *Econ. Bot.*(1982) 36: 203-223.
- (5) Small E and Catling PM. Canadian Medicinal Crops.

- NRC Research Press, 1999 pp. 7-12.
- (6) Saeidnia S, Gohari AR, Kiuchi F and Honda G. Effects of some fractions from *Achillea biebersteinii* and *A. millefolium* on the epimastigotes of *Trypanosoma cruzi*. *Iranian J. Pharm. Res.* (2004) 78-79.
 - (7) Goldberg AS, Evangelynne C and Eigen E. Isolation of the anti inflammatory principles from *Achillea millefolium* (compositae). *J. Pharm. Sci.* (2006) 58: 938-941.
 - (8) Falk AJ, Smolenski SJ, Bauer L and Bell CL. Isolation and identification of three new flavones from *Achillea millefolium* L. *J. Pharm. Sci.* (1975) 64: 1838-1842.
 - (9) Afsharypour S, Asgary S and Lockwood GB. Constituents of the essential oil of *Achillea wilhelmsii* from Iran. *Planta Med.* (1996) 62: 77-78.
 - (10) Javidnia K, Miri R and Sadeghpour H. Composition of the volatile oil of *Achillea wilhelmsii* C. Koch from Iran. *Daru* (2004) 12: 63-66.
 - (11) Arzi A and Akhavan M. The analgesic effect of hydroalcoholic extract of *Achillea wilhelmsii* on mice. *Med. J. Babol Univers. Med. Sc.* (2001) 4:11-14.
 - (12) Hydari MR, Ebrahimi S, Mehrabani M, Pardakhti A and Vafazadeh J. The anticonvulsant effect of methanolic extract of *Achillea wilhelmsii* C. Koch on mice. *ed. J. Babol Univers. Med. Sc.* (2005) 7: 7-13.
 - (13) Rezaeipoor R, Saeidnia S and Kamalinejad M. Immunosuppressive activity of *Achillea talagonica* on humoral immune responses in experimental animals. *J. Ethnopharmacol.* (1999) 65: 273-276.
 - (14) Asgari S, Naderi GH, Sarrafzadegan N, Mohammadifard N, Mostafavi S and Vakili R. Antihypertensive and antihyperlipidemic effects of *Achillea wilhelmsii*. *Drugs. Exp. Clin. Res.* (2000) 26: 89-93.
 - (15) Nickavar B, Kamalinejad M, Haj-yahya M and Shafaghi B. Comparison of the free radical scavenging activity of six iranian *Achillea* species. *Pharmac. Boil.* (2006) 44: 208-212.
 - (16) Canan K and Sebnem A. Antispasmodic effect of *Achillea nobilis* L. subsp. *Sipylea* (O. Schwarz) Baessler on the rat isolated duodenum. *J. Ethnopharmacol.* (2003) 84: 175-179.
 - (17) Harborne JB and Williams CA. Advances in flavonoid research since 1992. *Phytochem.* (2000) 55: 481-504.
 - (18) Hammad HM and Abdalla SS. Pharmacological effects of selected flavonoids on rat isolated ileum: structure-activity relationship. *Gen. Pharmacol.* (1997) 28: 767-771.
 - (19) Van Den Broucke CO and Lemli JA. Spasmodic activity of the flavonoids from *Thymus vulgaris*. *Pharmac. Weekblad* (1983) 5: 9-14.
 - (20) Abdalla S, Zarga MA, Afifi F, al-Khalil S, Mahasneh A and Sabri S. Effects of 3-O-methylquercetin on guinea-pig isolated smooth muscle. *J. Pharm. Pharmacol.* (1989) 41: 138-141.
 - (21) Rojas A, Cruz S, Ponce-Monter H and Mata R. Smooth muscle relaxing compounds from *Dodonaea viscosa*. *Planta Med.* (1996) 62: 154-159.
 - (22) Mustafa EH, Abu Zarga M and Abdalla S. Effects of cirsiliol, a flavone isolated from *Achillea fragrantissima*, on rat isolated ileum. *Gen. Pharmacol.* (1992) 23: 555-560. <http://www.ncbi.nlm.nih.gov/pubmed/1511863>.
 - (23) Boskabady MH. and Jandaghi P. Relaxant effects of carvacrol on guinea pig tracheal chains and its possible mechanisms. *Pharmazie* (2003) 58: 661-663.
 - (24) Boskabady MH., Khatami A. And Nazari A. Possible mechanism(s) for relaxant effects of *Foeniculum vulgare* on guinea pig tracheal chains. *Pharmazie* (2004) 59: 561-564.
 - (25) Holroyde MC. The influence of epithelium on the responsiveness of guinea pig isolated trachea. *Br. J. Pharmacol.* (1986) 87: 501-507.
 - (26) Buckle DR, Arch JRS, Boering NE, Foster KA, Taylor JF, Taylor SG and Shaw DJ. Relaxation effect of potassium channel activators BRL 38227 and Pinacidil on guinea pig and human airway smooth muscle, and blockade of their effects by Glibenclamide and BRL 31660. *Pulmon. Pharmacol.* (1993) 6: 77-86.
 - (27) Miyahara Y, Kizawa Y, Sano M and Murakami H. Effect of organic and inorganic Ca²⁺ antagonists on acetylcholine induced contraction in molluscan (*Mytilus edulis*) smooth muscle. *Gen. Pharmacol.* (1993) 24: 1419-1423.
 - (28) Linden A, Bergendal A, Ullman A, Skoogh BE and Lofdahl CG. Salmeterol, formoterol, and salbutamol in the isolated guinea-pig trachea: differences in maximum relaxant effect and potency but not in functional antagonism. *Thorax* (1993) 48: 547-553.
 - (29) Lronards B, Rampart M and Herman AG. Selective M₃ muscarinic receptor inhibit smooth muscle contraction in rabbit trachea without increasing the release of acetylcholine. *J. Pharmacol. Exp. Ther.* (1992) 263: 773-770.
 - (30) Boskabady MH, Ramezani M and Tabei T. Relaxant Effects of Different Fractions of Essential Oil from *Carum copticum* on Guinea Pig Tracheal Chains. *Phytother. Res.* (2003) 17: 1145-1149.

This article is available online at <http://www.ijpr.ir>