

***N*-(5-Mercapto-1,3,4-Thiadiazol-2-yl)-2-Phenylacetamide Derivatives: Synthesis and *In-vitro* Cytotoxicity Evaluation as Potential Anticancer Agents**

Ahmad Mohammadi-Farani^a, Neda Heidarian^{b,c} and Alireza Aliabadi^{c*}

^aDepartment of Pharmacology, Toxicology and Medical Services, Faculty of Pharmacy, Kermanshah University of Medical Sciences, Kermanshah, Iran. ^bStudents Research Committee, Kermanshah University of Medical Sciences, Kermanshah, Iran. ^cDepartment of Medicinal Chemistry, Faculty of Pharmacy, Kermanshah University of Medical Sciences, Kermanshah, Iran.

Abstract

A new series of *N*-(5-Mercapto-1,3,4-thiadiazol-2-yl)-2-phenylacetamide derivatives (3a-3j) were synthesized via an amidation reaction using EDC and HOBt in acetonitrile solvent at room temperature condition. Chemical structures were characterized by ¹H NMR, IR and MS spectroscopic methods and related melting points were also determined. The anticancer activity was evaluated using MTT procedure *in-vitro*. All compounds were tested against SKNMC (Neuroblastoma), HT-29 (Colon cancer) and PC3 (Prostate cancer) cell lines. According to the toxicological data, none of the synthesized derivatives exerted superior activity than doxorubicin as reference drug. Derivatives with *Ortho* chlorine (compound 3d), *meta* methoxy (compound 3h) and *meta* fluorine (compound 3b) substituents on the phenyl ring exhibited the best cytotoxic activity against SKNMC (IC₅₀ = 4.5 ± 0.035 μM), HT-29 (IC₅₀ = 3.1 ± 0.030 μM) and PC3 (IC₅₀ = 12.6 ± 0.302 μM) cell lines respectively.

Keywords: Synthesis; 1,3,4-Thiadiazole; Anticancer; Amidation.

Introduction

Cancer is a major problem throughout the world and is the second leading cause of mortality in developed countries. Since, many of the current treatments have problems with adverse effects and drug-resistance; there is a strong and vital demand for the discovery and development of effective new anticancer therapeutics. Currently, cancer is the second leading cause of death in the developed countries. Tremendous progress has been made in the war against cancer with the development of many novel chemotherapy

agents. The main treatments for neoplastic diseases involve surgery, chemotherapy and radiotherapy. Chemotherapy involves the use of low-molecular-weight drugs (doxorubicin, methotrexate, paclitaxel, ...) to selectively destroy tumor cells or at least limit their proliferation. Disadvantages of many anticancer agents include myelosuppression, gastrointestinal side effects (nausea and vomiting), hair loss (alopecia), and also the development of clinical resistance (1-7).

During recent years a wide and intense investigation of different pharmacophores and chemical classes containing 1,3,4-thiadiazole have been carried out. Many of these derivatives possess interesting and potential biological effects such as antimicrobial, antitubercular,

* Corresponding author:

E-mail: aliabadi.alireza@gmail.com




Figure 1. Structures of acetazolamide (carbonic anhydrase inhibitor) and sulfamethizole (antibacterial agent) as examples of drugs containing 1,3,4-thiadiazole ring.

antiviral, anti-inflammatory, anticonvulsant, antihypertensive, antioxidant, antifungal and anticancer activity and now there are in the market as common used drugs (Figure 1) (8-16).

Diverse chemical structures containing 1,3,4-Thiadiazole nucleus have been reported with potential anticancer activity (Figure 2). The 1,3,4-thiadiazole ring in anticancer agents performs its role in pharmacophores of apoptosis inducers and caspase activators, tyrosine kinase inhibitors, carbonic anhydrase inhibitors and *etc* (17-25). Hence, various mechanisms could be imagined for anticancer chemical structures that containing the 1,3,4-thiadiazole ring.

In the present study, we focused on the synthesis of novel thiol containing 1,3,4-thiadiazole derivatives and assessed their anticancer activity against three cancerous cell lines consist of PC3 (Prostate cancer), HT-29 (Colon cancer) and SKNMC (Neuroblastoma).

Results and Discussion

Chemistry

According to the scheme 1, 5-amino-1,3,4-thiadiazole-2-thiol (1) was treated directly with various derivatives of phenylacetic acid for amide bond formation. The reaction was carried out in the presence of EDC and hydroxybenzotriazole (HOBt) in acetonitrile as solvent. The completion of reaction was confirmed by thin

layer chromatography (TLC). After completion, the solvent was evaporated under reduced pressure and ethyl acetate and water were added. The aqueous phase was separated and the organic phase was washed two times by sodium bicarbonate 5%, diluted sulfuric acid and brine (26, 27). Anhydrous sodium sulfate was added and filtration was done. Ethyl acetate was removed using rotatory evaporator and the intended product was obtained as powder.

MTT assay

According to the Table 1, cytotoxicity of all synthesized compounds 3a-3j was evaluated against three cancerous cell lines using MTT method. PC3 (Prostate cancer), HT-29 (Colon cancer) and SKNMC (Neuroblastoma) was applied in this investigation. None of the synthesized compounds showed superior activity than doxorubicin as reference drug. Totally, all compounds exhibited a higher activity against PC3 and HT-29 cell lines in comparison with SKNMC. Fluorine substituent at position *meta* (compound 3b) demonstrated the best anticancer activity against SKNMC cell line. This pattern was also observed for PC3 cell line. But, *para* substitution (compound 3c) of fluorine afforded a better activity against HT-29 cell line. Chlorine moiety at position 2 of the phenyl ring rendered a high anticancer potency against SKNMC cell line ($IC_{50} = 4.5 \pm 0.035 \mu M$). Substitution of




Figure 2. Structures of two 1,3,4-thiadiazole based compounds with potential anticancer activity.

General procedure for synthesis of N-(5-Mercapto-1,3,4-thiadiazol-2-yl)-2-phenylacetamide derivatives (3a-3j)

According to the scheme 1, in a flat bottom flask appropriate phenylacetic acid derivative (1a-1j) was mixed with equimolar quantities of *N*-ethyl-*N*-dimethylaminopropyl carbodiimide (EDC) and hydroxybenzotriazole (HOBt) in acetonitrile solvent, and was stirred for 30 minutes at room temperature. Then, Equimolar quantity of 5-amino-1,3,4-thiadiazole-2-thiol (2) was added to the reaction medium. The stirring was continued for 24 h. After confirmation of the reaction end by TLC, the acetonitrile was evaporated and 20 mL of water and 20 mL of ethyl acetate were added to the residue. The organic phase was separated and was washed two times by sodium bicarbonate 5%, diluted sulfuric acid and brine. Anhydrous sodium sulfate was added, filtered and the ethyl acetate was evaporated. The obtained powder was washed by diethyl ether, *n*-hexane and purified by column chromatography (EtOAc/Petroleum ether).

2-(2-Fluorophenyl)-N-(5-mercapto-1,3,4-thiadiazol-2-yl)acetamide (3a)

mp. 224 °C, Yield: 63%, IR (KBr, cm⁻¹) $\bar{\nu}$: 3151, 2931, 2804, 1712, 1570, 1492, 1465, 1342, 1311, 1234, 1180, 1145, 1068, 759, 694. MS (*m/z*, %): 296 (M⁺, 55), 133 (48), 109 (100), 83 (15).

2-(3-Fluorophenyl)-N-(5-mercapto-1,3,4-thiadiazol-2-yl)acetamide (3b)

mp. 162 °C, Yield: 78%, IR (KBr, cm⁻¹) $\bar{\nu}$: 3147, 2924, 1708, 1570, 1485, 1458, 1307, 1253, 1138, 1068, 964, 767. MS (*m/z*, %): 296 (M⁺, 30), 133 (55), 109 (100), 83 (10).

2-(4-Fluorophenyl)-N-(5-mercapto-1,3,4-thiadiazol-2-yl)acetamide (3c)

mp. 158-160 °C, Yield: 72%, ¹H NMR (DMSO-d₆, 400 MHz) δ : 3.1 (brs, 1H, -SH), 3.71 (s, 2H, -CH₂CO-), 7.01 (t, 4-fluorophenyl), 7.30 (t, 4-fluorophenyl), 12.3 (brs, NH). IR (KBr, cm⁻¹) $\bar{\nu}$: 3140, 3100, 2924, 1963, 1554, 1504, 1296, 1220, 1120, 1062. MS (*m/z*, %): 269 (M⁺, 55), 133 (55), 109 (100), 83 (30), 59 (10).

2-(2-Chlorophenyl)-N-(5-mercapto-1,3,4-thiadiazol-2-yl)acetamide (3d)

mp. 212-216 °C, Yield: 82%, ¹H NMR (DMSO-d₆, 400 MHz) δ : 3.86 (s, 2H, -CH₂CO-), 3.98 (s, SH), 7.26-7.58 (m, 2-Chlorophenyl), 7.85-7.91 (m, 2-Chlorophenyl), 10.15 (brs, NH). IR (KBr, cm⁻¹) $\bar{\nu}$: 3147, 2924, 2862, 1708, 1570, 1469, 1307, 1145, 1064, 744.

2-(3-Chlorophenyl)-N-(5-mercapto-1,3,4-thiadiazol-2-yl)acetamide (3e)

mp. 210 °C, Yield: 46%, ¹H NMR (DMSO-d₆, 400 MHz) δ : 3.68 (s, 2H, -CH₂CO-), 3.81 (s, SH), 7.48-7.52 (m, 1H, H₅-3-Chlorophenyl), 7.49 (s, H₂-3-Chlorophenyl), 7.81 (d, 1H, *J* = 8 Hz, H₆-3-Chlorophenyl), 7.87 (d, 1H, *J* = 8 Hz, H₄-3-Chlorophenyl), 11.22 (brs, NH). IR (KBr, cm⁻¹) $\bar{\nu}$: 3151, 2927, 2804, 1701, 1573, 1469, 1311, 1222, 1153, 1064, 783, 682.

2-(4-Chlorophenyl)-N-(5-mercapto-1,3,4-thiadiazol-2-yl)acetamide (3f)

mp. 174 °C, Yield: 65%, ¹H NMR (DMSO-d₆, 200 MHz) δ : 3.73 (s, -SH), 4.86 (s, -CH₂CO-), 7.14-8.29 (m, aromatic), 13.21 (s, NH). IR (KBr, cm⁻¹) $\bar{\nu}$: 3444, 3137, 3095, 2850, 2718, 1687, 1551, 1519, 1492, 1352, 1089, 960, 818, 712. MS (*m/z*, %): 285 (M⁺, 3), 280 (45), 167 (94), 149 (100), 125 (12), 113 (15), 71 (20), 57 (20).

N-(5-Mercapto-1,3,4-thiadiazol-2-yl)-2-(2-methoxyphenyl)acetamide (3g)

mp. 200 °C, Yield: 47%, ¹H NMR (DMSO-d₆, 200 MHz) δ : 3.82 (s, 2H, -CH₂CO-), 3.89 (s, SH), 3.95 (s, 3H, -OCH₃), 7.02 (m, 2H, 2-Methoxyphenyl), 7.27-7.4 (m, 2H, 2-Methoxyphenyl), 11.44 (brs, NH). IR (KBr, cm⁻¹) $\bar{\nu}$: 3468, 3147, 2931, 2835, 1701, 1570, 1492, 1462, 1303, 1246, 1145, 1064, 1026, 798, 752. MS (*m/z*, %): 281 (M⁺, 73), 148 (75), 133 (15), 121 (100), 91 (90), 65 (15).

N-(5-Mercapto-1,3,4-thiadiazol-2-yl)-2-(3-methoxyphenyl)acetamide (3h)

mp. 114 °C, Yield: 65%, ¹H NMR (DMSO-d₆, 400 MHz) δ : 3.67 (s, SH), 3.79 (s, 2H, CH₂CO-), 3.84 (s, 3H, -OCH₃), 6.88 (m, 3-methoxyphenyl), 7.26-7.33 (m, 3-methoxyphenyl), 7.29 (s, H₂-3-methoxyphenyl), 11.24 (s, NH). IR (KBr, cm⁻¹) $\bar{\nu}$: 3410, 3147, 2927, 1701, 1570, 1489, 1465,

1307, 1261, 1149, 1064, 779, 759, 690.

N-(5-Mercapto-1,3,4-thiadiazol-2-yl)-2-(4-methoxyphenyl)acetamide (3i)

mp. 152 °C, Yield: 79%, ¹H NMR (DMSO-d₆, 200 MHz) δ: 3.35 (s, 2H, -CH₂CO-), 3.38 (s, -SH), 3.79 (s, 3H, -OCH₃), 7.76 (d, *J* = 8 Hz, Phenyl), 8.02 (d, *J* = 8 Hz, Phenyl), 13.1 (brs, NH). MS (*m/z*, %): 281 (M⁺, 45), 238 (25), 148 (90), 121 (100), 91 (25), 78 (30).

2-(4-Bromophenyl)-*N*-(5-mercapto-1,3,4-thiadiazol-2-yl)acetamide (3j)

mp. 198 °C, Yield: 68%, ¹H NMR (CDCl₃, 400 MHz) δ: 3.67 (s, SH), 3.75 (s, 2H, -CH₂CO-), 7.18 (d, H_{2,6}-4-Bromophenyl), 7.47 (d, H_{2,6}-4-Bromophenyl), 10.6 (brs, NH). IR (KBr, cm⁻¹) $\bar{\nu}$: 3150, 2980, 2825, 1699, 1571, 1558, 1487, 1400, 1319, 1296, 1064, 1012, 788. MS (*m/z*, %): 331 (M⁺+2, 15), 329 (M⁺, 15), 198 (30), 196 (30), 171 (60), 169 (60), 133 (90), 89 (100), 63 (30).

Cytotoxicity assay

Synthesized derivatives of 1,3,4-thiadiazole (compounds 3a-3j) were tested for cytotoxic activity at 0.1-500 mcg/mL concentration in three human cancer cell lines of PC3 cell (prostate cancer), HT-29 (Colon cancer) and SKNMC (Neuroblastoma). Cells from different cell lines were seeded in 96-well plates at the density of 8000–10,000 viable cells per well and incubated for 48 hours to allow cell attachment. The cells were then incubated for another 48-96 hours (depends to cell cycle of each cell line) with various concentrations of compounds 3a-3j. Cells were then washed in PBS, and 20 μL of MTT (3-(4, 5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide solution (5 mg/mL) were added to each well. An additional 4 hours of incubation at 37 °C were done, and then the medium was discarded. Dimethyl sulfoxide (60 μL) was added to each well, and the solution was vigorously mixed to dissolve the purple tetrazolium crystals. The absorbance of each well was measured by plate reader (Anthous 2020; Austria) at a test wavelength of 550 nm against a standard reference solution at 690 nm. The amount of produced purple formazan is proportional to the number of viable cells (26).

Acknowledgement

Authors appreciate from the research council of Kermanshah University of Medical Sciences for financial supports. This work was performed in partial fulfillment of the requirement for PharmD of Mrs Neda Heidarian.

References

- (1) Nussbaumer S, Bonnabry P, Veuthey J and Fleury-Souverain S. Analysis of anticancer drugs: A review. *Talanta*. (2011) 85: 2265-2289.
- (2) Kaushik G, Kaushik T, Khanduja S, Pathak CM and Khanduja K. Cigarette smoke condensate promotes cell proliferation through disturbance in cellular redox homeostasis of transformed lung epithelial type-II cells. *Cancer Lett*. (2008) 270: 120-131.
- (3) Albrecht T, Mecke M, Alexe DM, Coleman MP and Martin-Moreno JM. Making progress against cancer in Europe in 2008. *Eur. J. cancer* (2008) 44: 1451-1456.
- (4) Ferlay J, Autier P, Boniol M, Heanue M, Colombet M and Boyle P. Estimates of the cancer incidence and mortality in Europe in 2006. *Ann. Oncol*. (2007) 18: 581-592.
- (5) Jemal A, Siegel R, Ward E, Murray T, Xu J, Smigal C and Thun MJ. Cancer Statistics, 2006. *Cancer J. Clin*. (2006) 56: 106-130.
- (6) Bhuvha HA and Kini SJ. Synthesis, anticancer activity and docking of some substituted benzothiazoles as tyrosine kinase inhibitors. *J. Mol. Graph. Model* (2010) 29: 32-37.
- (7) Madhusudan S, Trivadi S and Ganesan TS. Tyrosine kinase inhibitors in cancer therapy. *Clinical. Biochem*. (2004) 37: 618-635.
- (8) Rajak H, Deshmukh R, Aggarwal N, Kashaw S, Kharya MD and Mishra P. Synthesis of Novel 2,5-Disubstituted 1,3,4-Thiadiazoles for Their Potential Anticonvulsant Activity: Pharmacophoric Model Studies. *Arch. Pharm. Chem. Life Sci*. (2009) 342: 453-461.
- (9) Abdel-Aziz M, Aly OM, Khan SS, Mukherjee K and Bane S. Synthesis, Cytotoxic Properties and Tubulin Polymerization Inhibitory Activity of Novel 2-Pyrazoline Derivatives. *Arch. Pharm. Chem. Life Sci*. (2012) 345: 535-548.
- (10) Deng XQ, Dong ZQ, Song MX, Shu B, Wang SB and Qua ZS. Synthesis and Anticonvulsant Activities of Some Triazolothiadiazole Derivatives. *Arch. Pharm. Chem. Life Sci*. (2012) 345: 565-573.
- (11) Rostom SAF, Badr MH, Abd El Razik HA, Ashour HMA and Abdel Wahab AE. Synthesis of Some Pyrazolines and Pyrimidines Derived from Polymethoxy Chalcones as Anticancer and Antimicrobial Agents. *Arch. Pharm. Chem. Life Sci*. (2011) 344: 572-587.
- (12) Mishra G, Singh AK and Jyoti K. Review article on 1, 3, 4-Thiadiazole derivatives and it's Pharmacological activities. *Int. J. Chem. Tech. Res*. (2011) 3: 1380-1393.
- (13) Singh AK, Mishra G and Jyoti K. Review on Biological

- Activities of 1,3,4-Thiadiazole Derivatives. *J. App. Pharm. Sci.* (2011) 5: 44-49.
- (14) Kharb R, Kaur P, Sharma PC and Yar MS. Significance of Thiadiazole derivatives as Antimicrobial agents. *Int. J. Res. Pharm. Biomed. Sci.* (2011) 4: 1520-1540.
- (15) Kalidhar U and Kaur A. 1,3,4-Thiadiazole Derivatives and their Biological Activities: A Review *Res. J. Pharm. Biol. Chem.* (2011) 4: 1091-1106.
- (16) Siddiqui N, Ahuja P, Ahsan W, Pandeya SN and Alam MS. Thiadiazoles: Progress Report on Biological Activities. *J. Chem. Pharm. Res.* (2009) 1: 19-30.
- (17) Kumar D, Reddy Vaddula B, Chang KH and Shah K. One-pot synthesis and anticancer studies of 2-arylamino-5-aryl-1,3,4-thiadiazoles. *Bioorg. Med. Chem. Lett.* (2011) 21: 2320-2323.
- (18) Farshori NN, Banday RM, Ahmad A, Khan AU and Rauf A. Synthesis, characterization, and *in vitro* antimicrobial activities of 5-alkenyl/hydroxyalkenyl-2-phenylamine-1,3,4-oxadiazoles and thiadiazoles. *Bioorg. Med. Chem. Lett.* (2010) 6: 1933-1938
- (19) Matysiak J and Opolski A. Synthesis and antiproliferative activity of *N*-substituted 2-amino-5-(2,4-dihydroxyphenyl)-1,3,4-thiadiazoles. *Bioorg. Med. Chem.* (2006) 14: 4483-4489.
- (20) Yusuf M and Jain P. Synthesis and biological significances of 1,3,4-thiadiazolines and related heterocyclic compounds. *Arab. J. Chem.* (2011) in press.
- (21) Rzeski W, Matysiakb J and Kandefer-Szerszen M. Anticancer, neuroprotective activities and computational studies of 2-amino-1,3,4-thiadiazole based compound. *Bioorg. Med. Chem.* (2007) 15: 3201-3207.
- (22) Chou JY, Lai SY, Pan SL, Jow GM, Chern JM and Guh JH. Investigation of anticancer mechanism of thiadiazole-based compound in human non-small cell lung cancer A549 cells. *Biochem. Pharmacol.* (2003) 66: 115-124.
- (23) Ibrahim DA. Synthesis and biological evaluation of 3,6-disubstituted [1,2,4]triazolo[3,4-b][1,3,4]thiadiazole derivatives as a novel class of potential anti-tumor agents. *Eur. J. Med. Chem.* (2009) 44: 2776-2781.
- (24) Supuran CT and Scozzafava A. Carbonic anhydrase inhibitors - Part 94. 1,3,4-Thiadiazole-2-sulfonamide derivatives as antitumor agents. *Eur. J. Med. Chem.* (2000) 35: 867-874.
- (25) Yang XH, Wen Q, Zhao TT, Sun J, Li X, Xing M, Lu X and Zhu HL. Synthesis, biological evaluation, and molecular docking studies of cinnamic acyl 1,3,4-thiadiazole amide derivatives as novel antitubulin agents. *Bioorg. Med. Chem.* (2012) 20: 1181-1187.
- (26) Aliabadi A, Hasanvand Z, Kiani A and Mirabdali SS. Synthesis and *in-vitro* cytotoxicity assessment of *N*-(5-(Benzylthio)-1,3,4-thiadiazol-2-yl)-2-(4-(trifluoromethyl)phenyl)acetamide with potential anticancer activity. *Iran. J. Pharm. Res.* (2012) 12: 687-693.
- (27) Aliabadi A, Andisheh S, Tyarani-Najaran Z and Tayarani-Najaran M. 2-(4-Fluorophenyl)-*N*-phenylacetamide derivatives as anticancer agents: synthesis and *in-vitro* cytotoxicity evaluation. *Iran. J. Pharm. Res.* (2012) 12: 267-271.

This article is available online at <http://www.ijpr.ir>
