Original Article

# **Comparative Binding Affinities of Flavonoid Phytochemicals with Bovine Serum Albumin**

Shuqing Liu<sup>*a*</sup>, Chunmei Guo<sup>*b*</sup>, Yimeng Guo<sup>*a*</sup>, Hongshan Yu<sup>*c*</sup>, Frederick Greenaway <sup>*d*</sup> and Ming-Zhong Sun<sup>*b*\*</sup>

<sup>a</sup>Department of Biochemistry, Dalian Medical University, Dalian, 116044, China. <sup>b</sup>Department of Biotechnology, Dalian Medical University, Dalian, 116044, China. <sup>c</sup>School of Bioengineering, Dalian Polytechnic University, Dalian 116034, China. <sup>d</sup> Carlson School of Chemistry and Biochemistry, Clark University, Worcester, MA 01610, USA

#### Abstract

Dietary flavonoids show beneficial effects in the prevention of chronic diseases. However, flavonoid bioavailability is poor, probably due to their interaction with serum albumins. In the current work, the binding interactions of eight related flavonoids, sharing a similar core structure, with bovine serum albumin (BSA) were investigated by fluorescence spectroscopy. The binding affinities of the flavonoids with BSA were in the order hesperetin ( $K_A=5.59 \times 10^5$ )> quercetin ( $4.94 \times 10^5$ ) > naringenin ( $3.04 \times 10^5$ ) > isoquercitrin ( $4.66 \times 10^4$ ) > icariin ( $3.60 \times 10^4$ ) > rutin ( $1.65 \times 10^4$ ) > hesperidin ( $2.50 \times 10^3$ ) > naringin ( $8.70 \times 10^2$ ). The associations of specific structural components of the flavonoids with their binding properties to BSA were also explored and hydrophobicity, functional group substituents, steric hindrance effects and the spatial arrangements of substituents seem to be the key factors for the affinities of flavonoids towards BSA. The results from the current work contribute to a better understanding of the transport of flavonoids in plasma and helping predict their physiological functions based on their intrinsic structures.

Keywords: Flavonoids; Bovine serum albumin; Binding capacity; Structure; Function.

# Introduction

Serum albumin (SA) is the most abundant protein in plasma. It is a single non-glycosylated protein of 585 amino acids with three homologous domains I, II and III. Each domain is composed of two separate subdomains A and B connected by random coils (1-3). SA acts as a depot and carrier of endogenous and exogenous compounds such as fatty acids, amino acids and pharmaceuticals (4-6). The binding capabilities of physiological molecules with SA are largely due to two major hydrophobic binding sites located within particular cavities in subdomains IIA and IIIA (7, 8). The drug-SA interaction plays a critical role in drug disposition and efficacy. The bound drug acts as a depot while the unbound drug produces the desired pharmacological effect. The drug-SA complexes dissociate to replenish the free drug that is removed by metabolic or elimination processes, which prolongs the duration of the drug action (9-11). The drug-SA binding affinity is the determining factor for the therapeutic, pharmacodynamic and toxicological properties of drugs (12-15). Bovine serum albumin (BSA) is one of the most extensively studied proteins due to its high structural

<sup>\*</sup> Corresponding author:

E-mail: smzlsq@163.com

homology with human serum albumin (HSA). The stable BSA-drug complex is an ideal model for gaining fundamental insights into HSA-drug and plasma-drug binding.

Flavonoids are important phytonutrient components widely distributed in plant foods. They have a wide range of biochemical and pharmacological properties, including antiplatelet, antioxidant, antiradical, anticarcinogenic, antiviral, antimicrobial, antithrombotic and antimutagenic activities (16, 17). Flavonoids are divided into flavonol, flavone, isoflavone, flavanone, flavanol and anthocyanidin classes depending on their molecular substitutes. The polyphenolic flavonoids share a 15-carbon skeleton core structure (represented as C<sub>6</sub>- $C_3$ - $C_6$ ) consisting of two phenyl (chromanol) rings linked through a pyran ring (18-20). Considerable epidemiological evidence indicates that the intake of dietary flavonoid phytochemicals reduces the incidence of and the mortality of degenerative diseases (21, 22). More attention has been attracted to flavonoids as they show great potential value in human health care. However, their bioavailability is often limited due to interaction with their transporter, SA (23, 24). The distribution, duration and intensity of physiological actions of flavonoids in the human body, as well as their metabolism and elimination, are correlated with their affinities towards SA (25, 26). Thus, investigation of the binding of flavonoids with SA is of fundamental importance and can help provide insights for new drug development.

Structural variations of flavonoids can affect their binding capacibilities to SA (27, 28). In this study, we investigated the binding affinities of quercetin, rutin, isoquercitrin, icariin, hesperetin, hesperidin, naringenin and naringin, which have similar structures, with BSA by fluorescence spectroscopy. The results indicated that minor modifications and/ or arrangements of the functional groups of flavonoids affected their binding affinities with BSA. The results provide bases for estimating/ evaluating the contributions of different functional groups of flavonoids to their binding with BSA. Results from current work give a better understanding of the importance of various structural components of flavonoids on

their transport in plasma, and could help in the selection of effective dietary flavonoid drugs for the prevention and treatment of human diseases.

## Experimental

# Materials and instruments

Quercetin, isoquercitrin, rutin, hesperetin, hesperidin, naringenin, naringin and icariin were generous gifts from Prof. Fengxie Jin of the School of Bioengineering, Dalian Polytechnic University, Dalian, China. Bovine serum albumin (BSA) with purity >98% was from Acros (Spain). All other chemicals were analytical grade from commercial sources. The HITACHI 650-60 fluorescence spectrometer was from Hitachi, Japan.

Purity analyses of eight flavonoid phytochemicals by HPLC

High performance liquid chromatography (HPLC) was performed to analyze the purity of flavonoid phytochemicals. Each of the eight flavonoids (0.55 mg) was dissolved separately in 2 mL of methanol and filtered through a 0.2 µm filter. For purity analysis, 10 µL of each flavonoid sample was analyzed by a Waters 2690/996 HPLC system using an Intersil ODS-3 C18 column (4.6  $\times$  250 mm) at 25 °C. The mobile phase was 60% (v/v) methanol containing 0.2% phosphoric acid. HPLC elution was performed with a flow rate of 1.0 mL/min. The absorbance wavelength was 254 nm for quercetin, isoquercitrin, rutin, hesperetin and hesperidin, 283 nm for naringenin and naringin, and 270 nm for icariin. Triplicate measurements were performed for each sample. HPLC plots were collected and processed using Millennium 32 chromatographic software.

Binding assays of eight flavonoid phytochemicals with BSA

A 10  $\mu$ M BSA stock solution was prepared in 0.1 M Tris-HCl buffer (pH 7.4). Stock solutions (0.2 mM) were prepared of the eight flavonoids in dimethyl sulfoxide (DMSO) and were stored at 4 °C prior to use.

The binding capacities of flavonoids with BSA were investigated (2, 14). Briefly, a series of

assay solutions was prepared by mixing 0.5 mL of 10 µM BSA with various volumes of the stock solution of each flavonoid, and diluting with 0.1 M Tris-HCl buffer (pH 7.4) to a total volume of 5 mL. The concentrations of quercetin, rutin, isoquercitrin, icariin, hesperetin, naringenin, hesperidin and naringin were in the ranges of 5.30×10<sup>-7</sup>-5.30×10<sup>-6</sup> M, 8.30× 10<sup>-7</sup>-3.73×10<sup>-6</sup> M, 1.55×10<sup>-6</sup>-5.18×10<sup>-6</sup> M, 1.04×10<sup>-6</sup>-5.18×10<sup>-6</sup> M, 1.04×10<sup>-6</sup>-5.18×10<sup>-6</sup> M, 5.18× 10<sup>-7</sup>-5.18×10<sup>-6</sup> M, 4.15×10<sup>-7</sup>-3.73×10<sup>-6</sup> M and 8.30×10<sup>-7</sup>-4.15×10<sup>-6</sup> M, respectively. In each tube, the concentration of BSA was constant while the concentration of flavonoids was different. The resultant mixtures were incubated at room temperature for 1 h and transferred into a quartz cuvette for analysis by fluorescence spectroscopy. Fluorescence spectroscopy was carried out at 25 °C with a HITACHI 650-60 spectrometer (Hitachi, Japan) interfaced to a microcomputer. The excitation wavelength was 280 nm and the emission wavelength was 340 nm, and the excitation and emission slit widths were both set at 5 nm.

The stereospecific interaction between flavonoid and BSA is quantified by the apparent association constant ( $K_A$ ) as previously reported (5, 14). For the binding of BSA with a flavonoid:

$$nD + P \xleftarrow{K_A} D_n P$$
 Equation (1)

Where P is BSA, D is flavonoid,  $D_n P$  is the complex and n is the number of flavonoid binding sites, and the equilibrium constant,  $K_A$  is given by:

$$K_{A} = \frac{[D_{n}P]}{[D]^{n}[P]} \qquad \qquad \text{Equation(2)}$$

$$\log K_{A} = \log \frac{[D_{n}P]}{[P]} - n\log[D]$$
 Equation(3)

$$\frac{[P]}{[P] + [D_n P]} = \frac{I}{I_0}$$
 Equation(4)

So the quenching equation is:

$$\log K_{A} = \log \frac{\Delta I}{I} - n\log[D]$$
 Equation(5)

Where [D] is the concentration of flavonoid,  $I_0$  is the fluorescence intensity in the absence of flavonoid, I is the fluorescence intensity in the presence of flavonoid, and  $\Delta I = I_0 - I$ .  $K_A$  and n can be calculated from equation (5).

#### Data statistical analysis

Triplicate assays were performed for each experiment. The data were expressed as the mean  $\pm$  standard deviation (S.D.). SPSS 13.0 software was used for all statistical analyses. One-way analysis of variance (ANOVA) was used to determine the significant differences in two comparisons. Statistical significance was set at *P*<0.05.

#### **Results and Discussion**

#### Structures of flavonoid phytochemicals

The flavonoids utilized the current work can be cataloged as flavonols (quercetin), and flavones (rutin, isoquercitrin and icariin), which all share the core structure of flavone and flavanones (hesperetin, hesperidin, naringenin and naringin) (Figure 1). On one hand, these flavonoids share a similar structure, but on the other hand they also have their individual intrinsic structures (Figure 1). The structural differences among these flavonoids might result in and explain their functional differences.

## Purity analyses of flavonoid phytochemicals

To ensure that the fluorescence results we obtained were indeed due to the flavonoid, a purity analysis was carried out. The HPLC chromatograms of quercetin, rutin, isoquercitrin, icariin, hesperetin, naringenin, hesperidin and naringin are shown in Figure 2. The peak of each individual flavonoid dominated its chromatogram and the calculated purities of the eight flavonoids were over 95% based on both the peak area and intensity height (Table 1).

## Binding of flavonoid phytochemicals to BSA

Epidemiological evidence indicates that a diet rich in flavonoids reduces the incidence of



Figure 1. The chemical structures of quercetin, rutin, isoquercitrin, icariin, hesperetin, hesperidin, naringenin and naringin.

chronic diseases such as cardiovascular diseases, diabetes, cancers, neuronal diseases and stroke (29, 30). However, whether flavonoids play a significant role in preventing such chronic diseases has been questioned because of their low bioavailability (31). The interactions of flavonoids with SA determine their bioavailability and toxicology. BSA-flavonoid binding has been investigated by analyzing the perturbation of the intrinsic BSA fluorescence due to its Trp residues, which is observed by excitation at a wavelength of 280 nm (5, 14, 15). The fluorescence emission intensity of BSA has been observed to decrease as flavonoid concentration increases until at a 1:1 BSA:flavonoid ratio the complex is not fluorescent. Thus flavonoid molecules quench the Trp fluorescence, indicating that they bind near the Trp position (5, 14, 15). Our results indicate that as the flavonoid concentration increases, the fluorescence intensity of BSA decreases remarkably, implying that flavonoids we have studied also bind with BSA at or near the



**Figure 2.** HPLC chromatograms of the eight flavonoids. Plots A, B, C, D, E, F, G and H are for quercetin, rutin, isoquercitrin, icariin, hesperetin, naringenin, hesperidin and naringin, respectively. HPLC was performed with an Intersil ODS-3 C18 column ( $4.6 \times 250$  mm) at 25 °C. The mobile phase was 60% (v/v) methanol containing 0.2% phosphoric acid. The column was eluted with a flow rate of 1.0 mL/ min. The absorbance wavelength was set at 254 nm for quercetin, rutin, isoquercitrin, hesperetin and hesperidin, 283 nm for naringenin and naringin, and 270 nm for icariin.

Flavonoids	$\mathbf{N}^{\mathbf{a}}$	Wavelength (nm) <sup>b</sup>	Retention time (min)	Purity (area) (%) <sup>c</sup>	Purity (height ) (%) <sup>d</sup>			
Quercetin	3	254	$6.75\pm0.01$	$98.22\pm0.02$	$98.49\pm0.02$			
Isoquercitrin	3	254	$4.51\pm0.01$	$97.12\pm0.02$	$97.20\pm0.02$			
Rutin	3	254	$4.40\pm0.01$	$99.33 \pm 0.03$	$99.33 \pm 0.02$			
Naringenin	3	254	$7.38\pm0.02$	$95.64\pm0.02$	$95.10\pm0.02$			
Naringin	3	254	$4.26\pm0.01$	$95.06\pm0.04$	$95.31\pm0.03$			
Hesperetin	3	283	$7.83\pm0.02$	$95.85\pm0.03$	$95.02\pm0.03$			
Hesperidin	3	283	$4.35\pm0.02$	$96.53 \pm 0.02$	$96.78\pm0.03$			
Icariin	3	270	$9.78\pm0.02$	$96.85\pm0.03$	$96.29\pm0.03$			

 Table 1. HPLC Determinations of the purities of the eight flavonoid phytochemicals.

 $^{a}$  N = the number of measurements per experiment;  $^{b}$  The wavelength used for monitoring the elution of flavonoid phytochemicals by HPLC;  $^{c}$  The purities of eight flavonoid phytochemicals were calculated based on their peak areas;  $^{d}$  The purities of eight flavonoid phytochemicals were calculated based on their peak areas;  $^{d}$  The purities of eight flavonoid phytochemicals were calculated based on their peak areas;  $^{d}$  The purities of eight flavonoid phytochemicals were calculated based on their peak areas;  $^{d}$  The purities of eight flavonoid phytochemicals were calculated based on their peak heights.

Trp residues. Figure 3 shows the plots of  $\log \Delta I/I$  *versus* log [D] for quercetin, rutin, isoquercitrin, icariin, hesperetin, naringenin, hesperidin and naringin for concentrations, [D], between 0.5  $\mu$ M and 5  $\mu$ M.

The K<sub>A</sub> values and the number (n) of binding sites for all eight flavonoids were calculated and are summarized in Table 2. Hesperetin had the greatest K<sub>A</sub> for binding with BSA  $(5.59 \times 10^5)$ , followed by quercetin  $(4.94 \times 10^5)$ , naringenin  $(3.04 \times 10^5)$ , isoquercitrin  $(4.66 \times 10^4)$ , icariin  $(3.60 \times 10^4)$ , rutin  $(1.65 \times 10^4)$ , hesperidin  $(2.50 \times 10^3)$  and naringin  $(8.70 \times 10^2)$ . SPPS 13.0 analysis indicated that the differences of binding activities of all eight flavonoid phytochemicals used in the current work are statistically significant (*P*<0.05). For all eight flavonoids the number of BSA binding sites was close to 1 (Table 2).

The binding capacities of flavonoids with BSA are determined by the specifics of their structures. Rutin and isoquercitrin share the same core flavone structure with quercetin but have no free 3-OH group on the C-ring due to their O-Glc- $\alpha$ -Rha and O- $\beta$ -Glc substituents (Figure 1), which leads to a decrease in the BSA binding association constant, K<sub>A</sub>, from 4.94 × 10<sup>5</sup> to 4.66 × 10<sup>4</sup> and 1.65 × 10<sup>4</sup>, respectively. A similar phenomenon was also observed for hesperetin and hesperidin, and for naringenin and naringin. When the free 7-OH groups in the A-ring of hesperetin and naringenin were replaced by 7-O- $\beta$ -Glc( $6 \rightarrow 1$ )- $\alpha$ -Rha and

7-O- $\beta$ -Glc(2 $\rightarrow$ 1)- $\alpha$ -Rha to become hesperidin and naringin, respectively, their BSA binding association constants decreased from 5.59×105 to  $2.50 \times 10^3$  and from  $3.04 \times 10^5$  to  $8.70 \times 10^2$ . Considering that the only structural differences between hesperidin and hesperetin, and naringin and naringenin are the replacement of the 7-OH group by 7-O- $\beta$ -Glc (6 $\rightarrow$ 1)- $\alpha$ -Rha and 7-O- $\beta$ - $Glc(2\rightarrow 1)$ - $\alpha$ -Rha, it is apparent that the presence of the free 7-OH enhances the binding capacities of flavanones with BSA. The hydroxyl group can interact with amino acid residues of BSA to form hydrogen bonds to improve the binding affinities. When the free -OH is replaced by saccharides, the molecular size increases and the resultant steric hindrances in the binding pocket result in a lower binding affinity to BSA. Furthermore, the additional polar saccharide groups render these molecules less hydrophobic, and could affect their orientation in relation to the hydrophobic environment of BSA (32-38). Thus substitution of the 3-OH and 7-OH groups has marked effects on the binding affinities of flavonoids with BSA.

When the 3-OH in the C-ring of quercetin was replaced by 3-O-Glc- $\alpha$ -Rha in rutin and 3-O- $\beta$ -Glc in isoquercitrin, the binding affinities to BSA decreased by one order of magnitude, but when the 7-OH groups in the A-rings of hesperetin and naringenin were replaced by 7-O- $\beta$ -Glc( $6 \rightarrow 1$ )- $\alpha$ -Rha and 7-O- $\beta$ -Glc( $2 \rightarrow 1$ )- $\alpha$ -Rha in hesperidin and naringin, the binding affinities dropped by two orders of magnitude.



**Figure 3.** Double-logarithm curves of BSA fluorescence for flavonoids. A, B, C, D, E, F, G and H are the plots of log ( $\Delta$ I/I) *versus* log [D] of quercetin, rutin, isoquercitrin, icariin, hesperetin, naringenin, hesperidin and naringin with the concentration ranges of  $5.30 \times 10^{7}$   $^{-}5.30 \times 10^{-6}$  M,  $8.30 \times 10^{-7}$ - $3.73 \times 10^{-6}$  M,  $1.55 \times 10^{-6}$ - $5.18 \times 10^{-6}$  M,  $1.04 \times 10^{-6}$ - $5.18 \times 10^{-6}$  M,  $1.04 \times 10^{-6}$ - $5.18 \times 10^{-6}$  M,  $5.18 \times 10^{-7}$ - $5.18 \times 10^{-6}$  M,  $4.15 \times 10^{-7}$ - $3.73 \times 10^{-6}$  M and  $8.30 \times 10^{-7}$ - $4.15 \times 10^{-6}$  M, respectively. Significant differences exist for all comparisons between any two individual flavonoids (P<0.05) by SPSS 13.0 statistical analysis.

Flavonoids subclass	Name of flavonoids	Molecular weight (MW)	Binding sites number (n)	Apparent association constant (K <sub>A</sub> )
Flavone 7 6 5 0 4' 3' 4' 5' 5' 6' 5'	Rutin (5,7,3 <sup>°</sup> ,4 <sup>°</sup> -OH) Isoquercitrin (5,7, 3 <sup>°</sup> ,4 <sup>°</sup> -OH) Icariin (5-OH, 4 <sup>°</sup> -OMe)	610.52 464.38 676.66	0.86 0.92 0.95	1.65×10 <sup>4</sup> 4.66×10 <sup>4</sup> 3.60×10 <sup>4</sup>
Flavonol 3' 4' 3' 4' 3' 4' 5' 4' 0 3' 4' 0 3' 4' 0 3' 4' 0 3' 4' 0 3' 4' 0 3' 4' 0 3' 4' 0 0 3' 4' 0 0 0 0 0 0 0 0	Quercetin (3,5,7, 3',4'-OH)	302.20	0.97	4.94×10 <sup>s</sup>
Flavanone $2'$ $3'$ $4'$ 7 $0$ $2'$ $5'$ $5'$	Hesperetin(5,7,3'-OH, 4'-OMe) Naringenin (5,7,4-OH)	302.29 272.26	1.08 1.06	5.59×10 <sup>s</sup> 3.04×10 <sup>s</sup>
	Hesperidin (5,3'-OH, 4'-OMe) Naringin (5,4'-OH)	610.58 580.55	0.74 0.63	$2.50 \times 10^{3}$ $8.70 \times 10^{2}$
	•••			

**Table 2.** The number of binding sites (n) and apparent association constants  $(K_{\Lambda})$  of eight flavonoid phytochemicals.

The result indicates that substituents on the hydroxyl group of  $C_7$  in the A ring are more important for flavonoid interaction with BSA than are substituents on the hydroxyl group of  $C_3$  in the C-ring and thus that the A ring might be the more critical structure for binding to the hydrophobic region of BSA.

In comparison, the binding affinity of icariin with BSA was much higher than of hesperidin and naringin. The difference in affinity can be ascribed to the 7-O- $\beta$ -Glc in the A ring of icariin being smaller than the 7-O-Glc- $\alpha$ -Rha of hesperidin and naringin. In addition, the 8-alkyne of icariin enhances its lipid solubility and lipid-soluble drugs are more likely to enter the hydrophobic cavity of BSA and form a drug-BSA complex (33, 34, 36, 37). The two effects combine to contribute to the higher binding affinity of icariin with BSA.

Taken together, the current work indicates

that the binding capacities of flavonoid phytochemicals with BSA are related to their structures. The above results agree with published data for the binding affinities of other kinds of flavonoids with BSA, for which hydrogen bonding, hydrophobic forces and steric hindrance effects have been considered to be the driving forces for BSA-flavonoid association (3, 9, 14, 27, 28, 32-38). The study of the interactions of quercetin, rutin, hyperin and baicalin with HSA, indicated that flavonoids containing the groups C(3)-OH and C(4)=O, or C(5)–OH and C(4)=O, or C(3')–OH and C(4')– OH could form flavonoid-HSA complexes (9, 14, 28, 32). The numbers of hydroxyl groups on the A and B-rings of flavonoids affected the binding affinities and the number of binding sites of flavonoids with BSA significantly. Increasing numbers of hydroxyl groups on the A- and B-rings increased binding affinities of

flavonols with BSA, presumably by forming hydrogen bonds. BSA has hydrophobic groups in the interior of its tertiary structure and polar groups at its surface. Hydrogen bonds may take place between -OH groups of flavonoids and polar groups at the BSA surface (35, 36). Glycosylation of their hydroxyl groups decreased the binding affinities of flavonoids with proteins (37, 38), as confirmed by our results. Glycosylation increases the polarity of the molecule and this lessens the ability of the flavonoid to penetrate into the Trprich hydrophobic interior regions of BSA. Glycosylation also increases the size of the molecule and steric hindrance effects in the binding pocket may also weaken the binding affinity of the flavonoid with BSA. Finally, glycosylation decreases the hydrophobicity of flavonoids, reducing their binding affinity with BSA (37, 38).

## Conclusion

In spite of relatively small differences in the chemical structures of the eight flavonoids studied, large differences were observed in their binding affinities to BSA. The binding affinities of flavonoids are closely associated with their intrinsic structures in that minor modifications/ arrangements of the functional groups affect the binding of flavonoids to BSA. Relative hydrophobicities, the presence of 3-OH and 7-OH groups, and steric hindrance effects seem to be the key factors determining the affinities of flavonoids towards BSA. The current work facilitates understanding of the structural bases of flavonoid transport in plasma and assists the prediction of their physiological behaviors.

#### Acknowledgement

This work was supported by grants from the National Natural Science Foundation of China (81171957), the Key Laboratory of the Department of Education of Liaoning (LS2010050), the Educational Department of Liaoning Province (2008S077), and the Distinguished Young Scholars in Universities of Liaoning Province Development Plan (LJQ2011094).

## References

- (1) Hegde AH, Sandhya B and Seetharamappa J. Evaluation of binding and thermodynamic characteristics of interactions between a citrus flavonoid hesperitin with protein and effects of metal ions on binding. *Mol. Biol. Rep.* (2011) 38: 4921-4929.
- (2) Bolli A, Marino M, Rimbach G, Fanali G, Fasano M and Ascenzi P. Flavonoid binding to human serum albumin. *Biochem. Biophys. Res. Commun.* (2010) 398: 444-449.
- (3) Xie MX, Xu XY and Wang YD. Interaction between hesperetin and human serum albumin revealed by spectroscopic methods. *Biochim. Biophys. Acta.* (2005) 1724: 215-224.
- (4) Zsila F, Bikadi Z and Simonyi M. Probing the binding of the flavonoid, quercetin to human serum albumin by circular dichroism, electronic absorption spectroscopy and molecular modelling methods. *Biochem. Pharmacol.* (2003) 65: 447-456.
- (5) Kanakis CD, Tarantilis PA, Polissiou MG, Diamantoglou S and Tajmir-Riahi HA. Antioxidant flavonoids bind human serum albumin. J. Mol. Struct. (2006) 798: 69-74.
- (6) Knjazeva T and Kaljurand M. Capillary electrophoresis frontal analysis for the study of flavonoid interactions with human serum albumin. *Anal. Bioanal. Chem.* (2010) 397:2211-2219.
- (7) Matei I and Hillebrand M. Interaction of kaempferol with human serum albumin: a fluorescence and circular dichroism study. *J. Pharm. Biomed. Anal.* (2010) 51: 768-773.
- (8) Tian JN, Liu JQ, Tian X, Hu ZD and Chen XG. Study of the interaction of kaempferol with serum albumin. J. Mol. Struct. (2004) 691: 197-202.
- (9) Diniz A, Escuder-Gilabert L, Lopes NP, Villanueva-Camanas RM, Sagrado S and Medina-Hernandez MJ. Characterization of interactions between polyphenolic compounds and human serum proteins by capillary electrophoresis. *Anal. Bioanal. Chem.* (2008) 391: 625-632.
- (10) Pastukhov AV, Levchenko LA and Sadkov AP. Spectroscopic study on binding of rutin to human serum albumin. J. Mol. Struct. (2007) 842: 60-66.
- (11) Xiao JB, Wu MX, Kai GY, Wang FJ, Cao H and Yu XB. ZnO-ZnSQDs interfacial heterostructure for drug and food delivery application: enhancement of the binding affinities of flavonoid aglycones to bovine serum albumin. *Nanomed-Nanotechnol.* (2011) 7: 850-858.
- (12) Xi JQ and Fan L. Study on the binding of puerarin to bovine serum albumin by isothermal titration calorimetry and spectroscopic approaches. J. Pherm. Anal. Calorim. (2010) 102: 217-223.
- (13) Mishra B, Barik A, Priyadarsini KI and Mohan H. Fluorescence spectroscopic studies on binding of a flavonoid antioxidant quercetin to serum albumins. J. *Chem. Sci.* (2005) 117: 641-647.
- (14) Bi SY, Ding L, Tian Y, Song DQ, Zhou X, Liu X and Zhang HQ. Investigation of the interaction between

flavonoids and human serum albumin. J. Mol. Struct. (2004) 703: 37-45

- (15) Fang R, Jing H, Chai Z, Zhao GH, Stoll S, Ren FZ and Liu F. Study of the physicochemical properties of the BSA: flavonoid nanoparticle. *Eur. Food Res. Technol.* (2011) 233: 275-283.
- (16) Cavia-Saiz M, Busto MD, Pilar-Izquierdo MC, Ortega N, Perez-Mateos M and Muniz P. Antioxidant properties, radical scavenging activity and biomolecule protection capacity of flavonoid naringenin and its glycoside naringin: a comparative study. J. Sci. Food Agric. (2010) 90: 1238-1244.
- (17) Macready AL, Kennedy OB, Ellis JA, Williams CM, Spencer JP and Butler LT. Flavonoids and cognitive function: a review of human randomized controlled trial studies and recommendations for future studies. *Genes Nutr.* (2009) 4: 227-242.
- (18) Khoo NK, White CR, Pozzo-Miller L, Zhou F, Constance C, Inoue T, Patel RP and Parks DA. Dietary flavonoid quercetin stimulates vasorelaxation in aortic vessels. *Free Radic. Biol. Med.* (2010) 49: 339-347.
- (19) Rao AS, Reddy SG and Babu PP. The antioxidant and antiproliferative activities of methanolic extracts from *Njavara* rice bran. *BMC. Complement. Altern. Med.* (2010) 10: 1-9.
- (20) Whitman SC, Kurowska EM, Manthey JA and Daugherty A. Nobiletin, a citrus flavonoid isolated from tangerines, selectively inhibits class A scavenger receptor-mediated metabolism of acetylated LDL by mouse macrophages. *Atherosclerosis* (2005) 178: 25-32.
- (21) Kalgaonkar S, Gross HB and Yokoyama W. Effects of a flavonol-rich diet on select cardiovascular parameters in a golden Syrian hamster model. *J. Med. Food* (2010) 13: 108-115.
- (22) Mustafa RA, Hamid AA, Mohamed S and Bakar FA. Total phenolic compounds, flavonoids, and radical scavenging activity of 21 selected tropical plants. J. Food Sci. (2010) 75: 28-35.
- (23) Dufour C and Dangles O. Flavonoid-serum albumin complexation: determination of binding constants and binding sites by fluorescence spectroscopy. *Biochim. Biophys. Acta.* (2005) 1721: 164-173.
- (24) Lorrain B, Dufour C and Dangles O. Influence of serum albumin and the flavonol quercetin on the peroxidase activity of metmyoglobin. *Free Radic. Biol. Med.* (2010) 48: 1162-1172.
- (25) Dufour C, Loonis M and Dangles O. Inhibition of the peroxidation of linoleic acid by the flavonoid quercetin within their complex with human serum albumin. *Free Radic. Biol. Med.* (2007) 43: 241-252.
- (26) Sengupta B and Sengupta PK. The interaction of

quercetin with human serum albumin: a fluorescence spectroscopic study. *Biochem. Biophys. Res. Commun.* (2002) 299: 400-403.

- (27) Benerjee A, Basu K and Sengupta PK. Interaction of 7-hydroxyflavone with human serum albumin: A spectroscopic study. J. Photoch. Photobio. B. (2008) 90: 33-40.
- (28) Zhang Y, Shi S, Sun X, Xiong X and Peng M. The effect of Cu<sup>2+</sup> on interaction between flavonoids with different C-ring substituents and bovine serum albumin: structure-affinity relationship aspect. J. Inorg. Biochem. (2011) 105: 1529-1537.
- (29) Galluzzo P and Marino M. Nutritional flavonoids impact on nuclear and extranuclear estrogen receptor activities. *Genes Nutr.* (2006) 1: 161-176.
- (30) Virgili F and Marino M. Regulation of cellular signals from nutritional molecules: a specific role for phytochemicals, beyond antioxidant activity. *Free Radic. Biol. Med.* (2008) 45: 1205-1216.
- (31) Crozier A, Jaganath IB and Clifford MN. Dietary phenolics: chemistry, bioavailability and effects on health. *Nat. Prod. Rep.* (2009) 26: 1001-1043.
- (32) Papadopoulou A, Green RJ and Frazier RA. Interaction of flavonoids with bovine serum albumin: a fluorescence quenching study. J. Agric. Food Chem. (2005) 53: 158-163.
- (33) Rawel HM, Frey SK, Meidtner K, Kroll J and Schweigert FJ. Determining the binding affinities of phenolic compounds to proteins by quenching of the intrinsic tryptophan fluorescence. *Mol. Nutr. Food Res.* (2006) 50: 705-713.
- (34) Liu EH, Qi LW and Li P. Structural relationship and binding mechanisms of five flavonoids with bovine serum albumin. *Molecules* (2010) 15: 9092-90103.
- (35) Xiao J, Suzuki M, Jiang X, Chen X, Yamamoto K, Ren F and Xu M. Influence of B-ring hydroxylation on interactions of flavonols with bovine serum albumin. J. Agric. Food Chem. (2008) 56: 2350-2356.
- (36) Xiao J, Cao H, Wang Y, Yamamoto K and Wei X. Structure-affinity relationship of flavones on binding to serum albumins: effect of hydroxyl groups on ring A. *Mol. Nutr. Food Res.* (2010) 54: 253-260.
- (37) Xiao J, Cao H, Wang Y, Zhao J and Wei X. Glycosylation of dietary flavonoids decreases the affinities for plasma protein. J. Agric. Food Chem. (2009) 57: 6642-6648.
- (38) Vial Y, Tod M, Hornecker M, Urien S, Conti F, Dauphin A, Calmus Y and Blanchet B. *In-vitro* influence of fatty acids and bilirubin on binding of mycophenolic acid to human serum albumin. *J. Pharm. Biomed. Anal.* (2011) 54: 607-609.

This article is available online at http://www.ijpr.ir