

Progesterone and Cilostazol Protect Mice Pancreatic Islets from Oxidative Stress Induced by Hydrogen Peroxide

Akram Ahangarpour^a, Hamid Heidari^{b*}, Seyyed Ali Mard^c,
Mahmoud Hashemitabar^d and Ali Khodadadi^e

^aHealth Research Institute, Diabetes Research Center, Department of Physiology, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran. ^bCellular and Molecular Research Center (CMRC), Department of Physiology, Faculty of Medical Sciences, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran. ^cPhysiology Research Center (PRC), Research Institute for Infectious Diseases of Digestive System and Department of Physiology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran. ^dCellular and Molecular Research Center (CMRC), Department of Anatomical Science, Ahvaz Jundishapur University of Medical Sciences, Faculty of Medicine, Ahvaz, Iran. ^eCellular and Molecular Research Center (CMRC), Department of Immunology Science, Ahvaz Jundishapur University of Medical Sciences, Faculty of Medicine, Ahvaz, Iran.

Abstract

Reactive oxygen species and oxidative stress impair β -cell function and reduce insulin secretion. It has been shown that progesterone and cilostazol possess antioxidant properties. The present study was aimed to investigate *in-vitro* pretreatment effect of progesterone and cilostazol on insulin secretion as well as their protective effects against hydrogen peroxide-induced oxidative stress in pancreatic isolated islets from mouse.

Pancreatic islets were isolated from 84 male NMRI mice (25-30 g) by collagenase digestion method and pretreated for 48 h with cilostazol (10 μ M), progesterone (0.5 μ M) and glibenclamide (10 μ M) in culture medium. Then islets were exposed to hydrogen peroxide (H_2O_2 500 μ M) for 2 h. Next, culture mediums containing glucose concentration of 2.8 mM or 16.7 mM were added to them and incubated in this status for 1 h. At the end, the rate of insulin output from islets, lipid peroxidation and antioxidant enzymes activities in islet tissues were assayed.

Exposure of islets to H_2O_2 , resulted in a significant decrease in insulin secretion, superoxide dismutase and catalase activities ($P < 0.001$). Also islets malondialdehyde levels were increased by H_2O_2 , after addition of 2.8 mM ($P < 0.05$) and 16.7 mM ($P < 0.001$) glucose. 48 h pretreatment of islets with cilostazol and progesterone, significantly reverted back this changes ($P < 0.05$).

Results of present study showed that cilostazol and progesterone protect mice pancreatic islets against H_2O_2 -induced oxidative stress. At the end, our results suggested that protective effects of progesterone and cilostazol are mediated by augmentation the antioxidant defence system of islets.

Keywords: Cilostazol; H_2O_2 ; Insulin; Islet; Progesterone.

* Corresponding author:

E-mail: physiology@yahoo.com

Introduction

Diabetes is one of the serious endocrine disorders in worldwide and has predicted that its prevalence will increase noticeably by the year 2030 (1).

In-vivo studies have indicated that excessive generation of reactive oxygen species (ROS) occurs in diabetes situation. Also oxidative stress is accompanied by imbalance between oxidant and antioxidant systems, has a critical role in development of this disease and lead to destruction of insulin producing pancreatic β -cells (2, 3).

Since β -cells, contain very low level of antioxidant defense enzymes such as superoxide dismutase (SOD) and catalase (CAT), these cells have extremely great sensitivity to free radical induced damage. On the other hand, elevations of cell antioxidant enzyme activities lead to protection against ROS (4). Also one of the detrimental effects of ROS is lipid peroxidation that may result in β -cells death and loss of insulin secretion through apoptosis process (5). So it seems that exposure of pancreatic islets to exogenous insulinotropic and antioxidant agents is very important in treatment strategies for diabetes. Several studies have shown that progesterone (PRO) possesses antioxidant properties such as scavenging of ROS in cancer cells (6) and increasing SOD activity in human endometrial stromal cells (7). Also Morrissy and *et al.* reported that progesterone can induce antioxidant genes expression in cardiomyocytes cells and exert antioxidant and antiapoptotic effects (8).

Cilostazol (CLZ), a selective phosphodiesterase inhibitor (PDEi), causes increase in intracellular level of cyclic adenosine monophosphate (cAMP) (9). Several investigation in different cells and tissues have indicated to inhibitory effect of CLZ on ROS and superoxide generation as well as its positive effect on hydroxyl radicals scavenging (10, 11). Also there are evidences that CLZ inhibited lipid peroxidation in brain tissue (12), and reduced oxidative stress through decrease the MDA level and improved glutathione level in blood of diabetic patients (13)

It has demonstrated that increase in cAMP

level, can reduce the oxidative stress effect on different cell (14). Cilostazol as a PDEi elevates intracellular level of cAMP (13). Also according to other studies, progesterone through its non-genomic effects increases the cAMP level (7). Since both of these compounds are able to enhance level of cAMP, the present study was performed to evaluate the protective effects of CLZ and PRO alone and in combination on H_2O_2 -induced islet cells damage.

Experimental

Animals

84 Male NMRI mice (25–30 g) (7 mice in each group) were obtained from animal house of Ahvaz Jundishapur University of Medical Science (Ahvaz, Iran) and housed in cages (22 ± 2 °C, under a standard 12 h light: 12 h dark cycle) and allowed *ad libitum* feed access. All experimental protocols were performed according to standards for animal care, established by the ethical committee of Ahvaz Jundishapur University of Medical Sciences (Ahvaz, Iran).

Isolation of mice pancreatic islets

Pancreatic islets were isolated from overnight-fasted male NMRI mice by Lacy and Kostianovsky modified collagenase digestion method (15). In brief, after cervical dislocation, abdomen of animals was opened. The common bile duct (CBD) was occluded at distal end close to the duodenum and 5 mL of Hank's Balanced Salt Solution (HBSS) [all units in mmol/l: 115 NaCl, 10 $NaHCO_3$, 5 KCl, 1.1 $MgCl_2$, 1.2 NaH_2PO_4 , 2.5 $CaCl_2$, 25 HEPES, and 5 D-glucose, pH 7.4 as well as 1% BSA (Merck, Germany)] containing 1.4 mg/mL of collagenase P (Roche, Germany) was injected into the duct (16, 17).

After removal of the pancreas, it was placed into a 50 mL conical tube and incubated for 15 min at 37 °C water bath. Thereafter 15 mL cold Hank's solution was added to tube to dilute collagenase concentration and stop further digestion process. For washing the collagenase from islet tissues, the tube was centrifuged for 2 min at 1200 rpm and supernatant has discarded. The washing procedure of islets, repeated again

and remainders were transferred to a blackened petri dish (17). The islets were separated by handpicking under a stereomicroscope (Euromex, Holland) and were cultured in RPMI-1640 medium (Gibco, USA) which supplemented with 10% fetal calf serum, 100 U/mL penicillin, 100 U/mL streptomycin, 5 mM D-glucose was and gassed with 95% O₂ - 5% CO₂ atmosphere.

Insulin secretion measurement

Insulin secretion was evaluated in a glucose static incubation. The islets were preincubated overnight in RPMI medium. Isolated mouse islets were divided into six groups in separated well, as each group contain 7 islets: (1) control islets cultured for 48 h (18), (2) islets cultured for 48 h and then in same medium were exposed to H₂O₂ 500 μM for 2 h (18), (3) islets cultured for 48 h pretreated with PRO (0.5 μM) (19), and then were exposed to H₂O₂ 500 μM for 2 h, (4) islets cultured for 48 h pretreated with CLZ (10 μM) (20), and then were exposed to H₂O₂ 500 μM for 2 h, (5) islets cultured for 48h pretreated with PRO (0.5 μM)+CLZ (10 μM) together and then were exposed to H₂O₂ 500 μM for 2 h, and (6) islets cultured for 48 h pretreated with Glibenclamide (GLB. 10 μM) (21), and then were exposed to H₂O₂ 500 μM for 2 h (All drugs purchased from Sigma, USA). Then islets in all groups were washed with HBSS and 1 mL culture medium including glucose concentration of 2.8 mM (basal concentration of glucose) or glucose concentration of 16.7 mM (stimulatory concentration of glucose) were added to them and incubated in 37 °C in this status for 1 h (18). Ultimately, the supernatant has taken and assayed by immunoradiometric assay (IRMA) method (DIA source INS-IRMA Kit, Neuve Belgium. C.N: KIP1251) for insulin concentration. The kit has detection limit of 1 μIU/mL, intra assay coefficient of variation (CV) of 2.1% and inter assay CV of 6.5%. Results were expressed as microU/islet/60 min for evaluating insulin secretion.

Preparation of the samples for biochemical analyses

For biochemical analysis, islets were isolated from overnight-fasted Male NMRI mice and

divided to 6 groups similar to those previously mentioned in the insulin secretion measurement protocol in this study, but in this analysis each group contained 50 islets. At the end, islets were incubated 1h in medium containing 2.8 mM or 16.7 mM concentration of glucose (basal and stimulatory), then islets washed three times with ice-cold phosphate-buffered saline (PBS) and lysed through sonication (heilscher - Gmbh, Germany) for 10 s and maintained at 4 °C followed by 10 min centrifugation (18). Supernatants were used immediately for assaying SOD and CAT activities as well as MDA level as follow:

MDA level measurement

Levels of malondialdehyde of islet tissues were measured according to this method (22) by monitoring the thiobarbituric acid reactive substance (TBARS) formation. Five hundred micro liter islets tissue supernatant was added to 1.5 mL trichloroacetic acid (10%). After centrifuging, supernatant (1.5 mL) was blended with 2 mL thiobarbituric acid (0.67%) and placed for 15 min in boiling water bath. After the sample has cooled, it was extracted with 2 mL *n*-butanol followed by centrifuge (at 4000× g for 15 min). The organic phase removed and the absorbance was read at 535 nm (spectrophotometer) using a blank containing all the reagents except the sample. Values were expressed as nmol/mg protein. The MDA standard curve was prepared using various dilution of 1,1,3,3-tetramethoxypropan.

Determination of Catalase activity

For measurement of CAT activity of islet tissues, we used the Claiborne's method (23). Initially, a mixture of 50 mM potassium phosphate (pH 7.0), 19 mM H₂O₂, and 20 mL supernatant of homogenate pancreas islets, was prepared. The H₂O₂ was added to this mixture at the end, for beginning the reaction. The rate of H₂O₂ decomposition was assessed by measuring the absorbance changes at 240 nm for 60 s. One unit of CAT activity is defined as 1 μM of hydrogen peroxide that is consumed in 1 min. ultimately the specific activity of CAT was expressed as unit/mg protein.




Figure 1. 48 h Pretreatment effects of progesterone (PRO. 0.5 μ M), cilostazol (CLZ. 10 μ M), PRO and CLZ together, Glibenclamide (GLB. 10 μ M) on insulin secretion from mice isolated pancreatic islets after 2 h exposure of islets to H_2O_2 (500 μ M) and subsequent 1 h incubation with 2.8 mM (A) or 16.7 mM (B) glucose containing medium. (7 mice in each group). Results are expressed as mean \pm SEM. *** (P < 0.001) vs. control group. ## (P < 0.01) and ### (P < 0.001) vs. H_2O_2 group.

Determination of Superoxide dismutase activity

The SOD activity in pancreatic islet tissues was estimated by applying the Suttle's method (24) with the SOD kit (Randox Labs, Crumlin, UK). The principle of this method is based on SOD ability for inhibiting the reduction of nitroblue tetrazolium (NBT) and formation of red formazan which its concentration is measured by spectrophotometer at 505 nm. 50% inhibition of NBT reduction of the enzyme is defined as 1 unit SOD. The SOD activity was expressed as unit per milligram protein.

Protein measurement

The concentration of protein was assayed according to Bradford method (25). Briefly, a volume of 20 μ L prepared sample was added to 1 mL Bradford reagent. Then the light absorbance was measured after 5 min in 595 nm. Bovine serum albumin was used as standard.

Statistical analysis

Data are expressed by SPSS as mean \pm SEM. One-way Analysis Of Variance (ANOVA) was used for comparison of data of different groups, followed by Tukey's test. p-value < 0.05 was considered as statistically significant differences.

Results and Discussion

Effects of PRO, CLZ on insulin secretion from pancreatic islets

The rate of insulin secretion (microU/islet/60 min) from isolated islets were decreased significantly in the presence of 500 μ M H_2O_2 (1.47 ± 0.15 and 4.09 ± 0.13) in comparison with control group, (2.26 ± 0.16 and 6.84 ± 0.17) in basal (Figure 1A) and stimulatory (Figure 1B) glucose medium, respectively (P < 0.001). Also insulin secretion showed a significant increase in pretreated groups with CLZ (2.16 ± 0.15), (P < 0.01), GLB (3.65 ± 0.16) and PRO+CLZ (3.41 ± 0.14), (P < 0.001) compared to H_2O_2

Table 2. The Effects of progesterone (0.5 μ M), cilostazol (10 μ M), Glibenclamide (10 μ M) pretreatment on the MDA level, SOD and CAT activities of mice isolated pancreatic islets following 2 h exposure of islets to H₂O₂ (500 μ M) and subsequent 1h incubation with 16.7 mM glucose containing medium. (7 mice in each group).

Groups	Enzyme	Catalas activity (U/mg Protein)	SOD activity (U/mg Protein)	MDA (nmol/mg Protein)
Control		4/06 \pm /18	37/96 \pm 4/9	6/22 \pm /08
H ₂ O ₂		1/5 \pm /12 ***	17/66 \pm 3/6 ***	9/8 \pm /17 ***
H ₂ O ₂ + GLB		3/12 \pm /12 *** ###	31/31 \pm 5 * ###	7 \pm /199 ###
H ₂ O ₂ + PRO		2/05 \pm /11 *** #	25/03 \pm 3/9 ***	8/2 \pm /19 *** ###
H ₂ O ₂ + CLZ		2/53 \pm /1 *** ###	27/69 \pm 3/5 *** ##	7/8 \pm /13 *** ###
H ₂ O ₂ + CLZ + PRO		3/54 \pm /15 * ###	36/67 \pm 4/44 ###	4/6 \pm /2 *** ###

H₂O₂: hydrogen peroxide. GLB: Glibenclamide. CLZ: cilostazol. PRO: progesterone

Results are expressed as mean \pm SEM

* (P<0.05) and *** (P<0.001) vs. control group. # (P<0.05) and ## (P<0.01) and ### (P<0.001) vs. H₂O₂ group.

group in basal glucose medium (Figure 1A). Furthermore a remarkable increase in insulin secretion was observed at the stimulatory glucose medium (Figure 1B) in groups which pretreated with PRO and GLB (4.83 \pm 0.16 and 4.75 \pm 0.17 respectively), (P < 0.01), CLZ and PRO+CLZ (5.99 \pm 0.15 and 6.56 \pm 0.16 respectively) (P < 0.001), compared with H₂O₂ alone.

Pancreatic islets MDA levels

MDA levels (nmol/mg protein) following incubation of islets with 500 μ M H₂O₂ increased significantly (P < 0.001) compared to control group in 16.7 mM glucose medium. Also 48 h treatment of islets with PRO+CLZ and GLB decreased (P < 0.001) MDA level as compared to H₂O₂ group in 2.8 mM glucose medium (Table 1). Relative to H₂O₂ group in 16.7 mM glucose medium, significant decrease in MDA level were observed (P < 0.001) in all pretreatment groups. The effect of PRO+CLZ was better than other groups (Table 2).

Pancreatic islets CAT levels

According to acquired results, after addition of high and basal glucose mediums to H₂O₂ group, CAT activity of islet tissues (unit/mg protein) decreased significantly compared to control groups (P < 0.001). Under 2.8 mM glucose, all pretreated drugs were able to increase (P < 0.001) CAT activity relative to H₂O₂ group (Table 1). After addition of 16.7 mM glucose medium, remarkably increase in CAT activity was observed in PRO pretreated

(P < 0.05), and also other pretreated groups (P < 0.001) in comparison with H₂O₂ group (Table 2).

Pancreatic islets SOD levels

After addition of 2.8 mM glucose (Table 1) and also 16.7 mM glucose containing mediums (Table 2), islets SOD activity (unit/mg protein) of negative control group, significantly decreased compared to control group (P < 0.001). Following the addition of basal glucose medium (Table 1), significant improvement of SOD activity was shown in PRO (P < 0.01) and other pretreatment groups (P < 0.001) compared to H₂O₂ group. Pretreatment with CLZ (P < 0.01), PRO+CLZ and GLB (P < 0.001), increased SOD activity in the stimulatory glucose medium (Table 2) compared to H₂O₂ group.

In this study, H₂O₂ is used as a substitute of ROS for evaluation of ROS effects on islet cells function. This model has also been used widely in previous researches for assessing the role of ROS (26). In our investigation, exposure of mice islets to H₂O₂ significantly decreased insulin release from them after addition of 2.8 mM and also 16.7 mM glucose mediums. Our findings are consistent with Xiong and *et al.* report (18) which showed that H₂O₂ damages islet cells and subsequent decreases their insulin secretion. The present study also exhibited that pretreatment of H₂O₂ -induced damaged islets with PRO, increased insulin secretion. In agreement with our study, Shao and *et al.* reported that PRO stimulates insulin secretion in MIN6 β -cells

(27). Also similar excitatory effect on insulin secretion has been achieved by Hollew and *et al.* after 20 h incubation of rat islets with PRO (28). However, Sorenson and *et al.* have shown the inhibitory effect of PRO on insulin secretion from neonatal rat islets (9). Although the accurate cause of this differential response to PRO is not apparent, but it may result from the different kinds of islets that have been studied. In the medium culture, neonatal islets may not be able to respond efficiently to glucose stimulation for insulin secretion and this procedure perhaps is due to immaturity of the metabolic system of neonatal pancreatic beta cells (29).

Also in our study pretreatment with CLZ remarkably increased insulin secretion as compared with H₂O₂ group. The present study showed that combined pretreatment of PRO and CLZ has a stronger effect on insulin secretion from islets.

Because of the vulnerability of pancreatic islets to oxidative damage, exposure of them to ROS can activate several cellular stress-sensitive pathways that have been linked to decreased insulin secretion (30). Our results in this study showed that H₂O₂ as a substitute of ROS, decreased insulin output of pancreatic islets by induction of oxidative stress. Administration of compounds with antioxidant properties can augment the defense capacity of islet cells to deal with oxidative stress (31). Likely pretreatment of islets with CLZ or PRO, to some extent restored secretary function of islets through potentiating of islets antioxidant defense system. On the other hand combination of CLZ and PRO represent higher antioxidant properties and so better preserve islets function.

The rate of lipid peroxidation and destruction of the cell membrane was assayed by evaluation of MDA level (32). At this study, expose to H₂O₂ (as an oxidative stress inducer) significantly increased the MDA level of islet cells of H₂O₂ group after addition of stimulatory concentration of glucose medium in comparison with the control group.

Also several studies which have investigated the toxicity effects of Interleukin1- β , tumor necrosis factor- α and interferon gamma on isolated islets of rats and human, demonstrated that this cytokine mixture induced significant

increases in MDA and decreases in insulin content in islets. In addition, an antioxidant compound (lazaroid U78518E) inhibited this cytokine effects, decreased MDA level and increased insulin of islets (33, 34)

Our findings indicated that the islets MDA level in 16.7 mM glucose medium was higher compared to 2.8 mM glucose medium. It is probably because high concentration of glucose, itself leads to increased ROS production, oxidative damage, as well as cell toxicity. Subsequently this results in more lipid peroxidation and higher MDA generation (35).

Also pretreatment with PRO+CLZ reduced lipid peroxidation and MDA generation in islet cells after addition of 16.7 mM glucose medium and this may partly protect the islets against H₂O₂-induced damage. Pretreatment with PRO+CLZ resulted in more decrease in lipid peroxidation and thus further resistance of islets cells versus free radical damages rather than alone pretreatment of them.

Our results showed that in H₂O₂ group, the antioxidant enzyme activities including SOD and CAT significantly decreased relative to control groups and this pointed to the oxidative effect of H₂O₂ in islet cells. Markus and *et al.* have indicated to similar effect of H₂O₂ on insulin-producing RINm5F cells, and also cited that over expression of these antioxidant enzymes, reduced H₂O₂-mediated toxicity in these cells (36).

In another study, Lortz and *et al.* investigated the combination effect of IL-1 β , TNF- α and IFN- γ on insulin-producing RINm5F cells. They showed that this compound significantly decreased cell viability. Over expression of cytoprotective enzymes such as CAT and SOD protected against toxicity of the cytokine mixture due to inactivation of generated ROS in this cells (37).

Pretreatment with PRO and CLZ elevated CAT activity as compared with H₂O₂ group. Also pretreatment with CLZ alone or with PRO, improved the SOD activity relative to H₂O₂ group after addition of 16.7 mM and 2.8 mM glucose medium. However after addition of basal or stimulatory glucose mediums in our investigation, combined of PRO and CLZ has stronger effect than single pretreatment of them on antioxidant enzymes activities. As observed in this study, the PRO+CLZ effect on lipid peroxidation as well as

their antioxidant effect was more impressive than single pretreatment of them.

It has also been demonstrated that increase in intracellular cyclic nucleotides such as cAMP, could decrease reactive oxygen generation, oxidative stress and subsequent development of cellular dysfunction (14). On the other hand, as mentioned previously, PRO and CLZ probably by elevation of cyclic AMP and also activation of enzymes such as PK-A, could potentiate the cell defense systems against oxidative stress. Apparently, simultaneous pretreatment of PRO and CLZ rather than alone pretreatment of them, create a higher cAMP level, and thereby stronger antioxidant properties as well as more protective effects following oxidative damages in pancreatic islets.

The main limitation in this work is the loss of some intact islets at the end of isolation procedure due to the unavoidable prolonged time necessary for islet isolation.

In conclusion, the present study showed the protective effects of PRO and CLZ against H₂O₂ induced oxidative damage in pancreatic islets. Based on the results of our study, this effect might be mediated through a reduction of lipid peroxidation and elevation of antioxidant enzyme activities. Anyway, further studies seem required to determine the precise molecular mechanism (s) by which combination of these drugs, affect on pancreatic islets.

Acknowledgement

This study is a part of Ph.D. thesis and also financially supported by vice chancellor of research affairs of the Ahvaz Jundishapur University of Medical Sciences (AJUMS) (Grant No. CMRC-61). The authors also thank the Cellular and Molecular Research Center of AJUMS experienced personnel.

Reference

- (1) Baluchnejad mojarada T and Roghani M. Chronic Oral Epigallocatechin-gallate alleviates streptozotocin-induced diabetic neuropathic hyperalgesia in rat: involvement of oxidative stress. *Iran. J. Pharm. Res.* (2012) 11: 1243-1253.
- (2) Rodrigues GR, Porawski M, Marcolin E, Kretzmann AN, Falcao Ferraz A and Richter MF. Treatment with aqueous extract from croton cajucara benth reduces hepatic oxidative stress in streptozotocin-diabetic rats. *J. Biomed. Biotechnol.* (2012) 2012: 1-7.
- (3) Ahmadvand H, Tavafic M and Khalatbary AR. Hepatoprotective and hypolipidemic effects of satreja khuzestanica essential oil in alloxan-induced type 1 diabetic rats. *Iran. J. Pharm. Res.* (2012) 11: 1219-1226.
- (4) Gabriele W, Nicole A, Marta M, Antje B, Jurgen S and James FB. Peroxiredoxin III protects pancreatic beta cells from apoptosis. *J. Endocrinol.* (2010) 207: 163-175.
- (5) Vessby J, Basu S, Mohsen R, Berne C and Vessby B. Oxidative stress and antioxidant status in type 1 diabetes mellitus. *J. Intern. Med.* (2002) 251: 69-76.
- (6) Nguyen H and Syed V. Progesterone inhibits growth and induces apoptosis in cancer cells through modulation of ROS. *Gynecol. Endocrinol.* (2011) 27: 830-836.
- (7) Matsuoka A, Kizuka F, Lee L, Tamura I, Taniguchi K and Asada H. progesterone Increases Manganese Superoxide Dismutase Expression via a cAMP-Dependent Signaling Mediated by Noncanonical Wnt5a Pathway in Human Endometrial Stromal Cells. *J. Clin. Endocrinol. Metab.* (2010) 95: 291-299.
- (8) Morriissy S, Xu B, Aguilar D, Zhang J and Chen QM. Inhibition of apoptosis by progesterone in cardiomyocytes. *Aging Cell* (2010) 9: 799-809.
- (9) Sorenson RL, Brelje TC and Roth C. Effects of steroid and lactogenic hormones on islets of Langerhans: a new hypothesis for the role of pregnancy steroids in the adaptation of islets to pregnancy. *Endocrinol.* (1993) 133: 2227-2234.
- (10) Kim KY, Shin HK, Choi JM and Hong KW. Inhibition of lipopolysaccharide-induced apoptosis by cilostazol in human umbilical vein endothelial cells. *J. Pharmacol. Exp. Ther.* (2002) 300: 709-715.
- (11) Lee WC, Chen HC, Wang CY, Lin PY, Ou TT and Chen CC. Cilostazol ameliorates nephropathy in type 1 diabetic rats involving improvement in oxidative stress and regulation of TGF-Beta and NF-kappaB. *Biosci Biotechnol. Biochem.* (2010) 74: 1355-1361.
- (12) Watanabe T, Zhang N, Liu M, Tanaka R, Mizuno Y and Urabe T. Cilostazol protects against brain white matter damage and cognitive impairment in a rat model of chronic cerebral hypoperfusion. *Stroke* (2006) 37: 1539-1545.
- (13) Agrawal NK, Maiti R, Dash D and Pandey BL. Cilostazol reduces inflammatory burden and oxidative stress in hypertensive type 2 diabetes mellitus patients. *Pharmacol. Res.* (2007) 56: 118-123.
- (14) Milani E, Nikfar S, Khorasani R, Zamani MJ and Abdollahi M. Reduction of diabetes-induced oxidative stress by phosphodiesterase inhibitors in rats. *Comp. Biochem. Physiol. C Toxicol. Pharmacol.* (2005) 140: 251-255.
- (15) Lacy PE and Kostianovsky M. Method for the isolation of intact islets of Langerhans from the rat pancreas. *Diabetes* (1967) 16: 35-39
- (16) Quesada I, Todorova MG, Beltrá M, Carneiro EM,

- Nadal A and Soria B. Glucose induces opposite intracellular Ca²⁺ concentration oscillatory patterns in identified alpha- and beta-cells within intact human islets of Langerhans. *Diabetes* (2006) 55: 2463-2469.
- (17) Jeffrey DC, Stacey BD, Kathryn LC, Runpei Wu and Craig SN. A Practical Guide to Rodent Islet Isolation and Assessment. *Biol. Proced. Online* (2009) 11: 3-29.
- (18) Xiong FL, Sun XH, Gan L, Yang XL and Xu HB. Puerarin protects rat pancreatic islets from damage by hydrogen peroxide. *Eur. J. Pharmacol.* (2006) 529: 1-7.
- (19) Irem N, Mamdouh H, Michelle E and Gladys T. The role of pregnancy hormones in the regulation of Pdx-1 expression. *Mol. Cell Endocrinol.* (2005) 233: 1-13.
- (20) Park SY, Bae JU, Hong KW and Kim CD. HO-1 Induced by Cilostazol Protects Against TNF- α associated Cytotoxicity via a PPAR- γ -dependent Pathway in Human Endothelial Cells. *Korean J. Physiol. Pharmacol.* (2011) 15: 83-88.
- (21) Hargrove DM, Nardone NA, Persson LM, Andrews KM, Shepherd KL, Stevenson RW and Parker JC. Comparison of the glucose dependency of glucagon-like peptide-1 (7-37) and glyburide *in-vitro* and *in-vivo*. *Metabolism* (1996) 45: 404-409.
- (22) Armanna B, Hansona MS, Hatcha E, Steffena A and Fernandez LA. Quantification of Basal and Stimulated ROS Levels as Predictors of Islet Potency and Function. *Am. J. Transplant* (2007) 7: 38-47.
- (23) Clairborne A. *Catalase Activity*. In: Greenwald RA. (ed.) Handbook of Methods for Oxygen Radical Research. Boca Raton, Florida: CRC Press (1985) 283-284.
- (24) Suttle NF. Copper deficiency in ruminants; recent developments. *Vet. Rec.* (1986) 119: 519-522.
- (25) Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* (1976) 72: 248-254.
- (26) Kuchmerovs'ka TM, Donchenko HV, Tykhonenko TM, Huzyk MM, Stavnichuk RV and Klymenko AP. Nicotinamide influence on pancreatic cells viability. *Ukr. Biokhim. Zh.* (2012) 84: 81-88.
- (27) Shao J, Qiao L and Friedman JE. Prolactin, progesterone, and dexamethasone coordinately and adversely regulate glucokinase and cAMP/PDE cascades in MIN6 β -cells. *Am. J. Physiol. Endocrinol. Metab.* (2004) 286: 304-310.
- (28) Howell SL, Tyhurst M and Green IC. Direct effects of progesterone on rat islets of Langerhans *in-vivo* and in tissue culture. *Diabetologia.* (1977) 13: 579-583.
- (29) Jermendy A, Toschi E, Aye T, Koh A, Sharma A and Bonner-Weir S. Rat neonatal beta cells lack the specialized metabolic phenotype of mature beta cells. *Diabetologia.* (2011) 54: 594-604.
- (30) Evans JL, Goldfine ID, Maddux BA and Grodsky GM. Oxidative stress and stress-activated signaling pathways: a unifying hypothesis of type 2 diabetes. *Endocr. Rev.* (2002) 23: 599-622.
- (31) Karunakaran U and Park KG. A Systematic Review of Oxidative Stress and Safety of Antioxidants in Diabetes: Focus on Islets and Their Defense. *Diabetes Metab. J.* (2013) 37: 106-112.
- (32) Zhu HQ, Gao Y, Wang JZ, Pan SH, Jiang HC and Dai WJ. Pretreatment with Bilirubin Protects Islet against Oxidative Injury During Isolation and Purification. *Transplant Proc.* (2011) 43: 1810-1814.
- (33) Suarez-Pinzon WL, Strynadka K and Rabinovitch A. Destruction of rat pancreatic islet beta-cells by cytokines involves the production of cytotoxic aldehydes. *Endocrinol.* (1996): 137: 5290-5296.
- (34) Rabinovitch A, Suarez-Pinzon WL, Strynadka K, Lakey JR and Rajotte RV. Human pancreatic islet beta-cell destruction by cytokines involves oxygen free radicals and aldehyde production. *J. Clin. Endocrinol. Metab.* (1996) 81: 3197-3202.
- (35) Duprez J, Roma LP, Close A-F and Jonas J-C. Protective antioxidant and antiapoptotic effects of zncl2 in rat pancreatic islets cultured in low and high glucose concentrations. *PLoS ONE.* (2012) 7: 46831.
- (36) Markus T, Stephan L, Rex M and Sigurd L. Complementary action of antioxidant enzymes in the protection of bioengineered insulin-producing RINm5F cells against the toxicity of reactive oxygen species. *Diabetes* (1998) 47: 1578-1585.
- (37) Lortz S, Tiedge M, Nachtwey T, Karlsen A.E, Nerup J and Lenzen S. Protection of insulin-producing RINm5F cells against cytokine-mediated toxicity through overexpression of antioxidant enzymes. *Diabetes* (2000) 49: 1123-1130.

This article is available online at <http://www.ijpr.ir>