

## Synthesis and Antiplatelet Aggregation Activity Evaluation of some 2-Aminopyrimidine and 2-Substituted-4,6-diaminopyrimidine Derivatives

Marjan Esfahanizadeh<sup>a,d</sup>, Shohreh Mohebbi<sup>b</sup>, Behnam Dasht Bozorg<sup>a</sup>, Salimeh Amidi<sup>a</sup>,  
Ali Gudarzi<sup>a</sup>, Seyed Abdolmajid Ayatollahi<sup>a,c</sup> and Farzad Kobarfard<sup>a,c,d\*</sup>

<sup>a</sup>Department of Medicinal Chemistry, School of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran. <sup>b</sup>Department of Medicinal Chemistry, School of Pharmacy, Zanjan University of Medical Sciences, Zanjan, Iran. <sup>c</sup>Phytochemistry Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran. <sup>d</sup>Central Research Laboratories, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

### Abstract

A series of novel 2-aminopyrimidine and 2-Substituted-4,6-diaminopyrimidine derivatives have been synthesized and their antiplatelet aggregation activities were assessed against ADP and arachidonic acid-induced platelet aggregation in human plasma using light transmission aggregometry. Among the tested derivatives, compounds Ia, I<sub>b</sub>, I<sub>B</sub> and II<sub>16</sub> exhibited the highest antiplatelet aggregation activity (36.75, 72.4, 62.5 and 80 μM). None of the compounds showed satisfactory activity against the aggregation induced by ADP but acceptable activities were observed against the aggregation induced by arachidonic acid. 2-aminopyrimidines were more active than 4,6-diaminopyrimidines in this respect.

**Keywords:** 2-aminopyrimidines; 2-Substituted-4,6-diaminopyrimidines; Antiplatelet aggregation.

### Introduction

Platelets play an important role in maintaining cardiovascular integrity and in regulating the bleeding process by blood-clot formation (1). However, uncontrolled platelet aggregation is dangerous in arterial blockage and may lead to life threatening disorders (2). Antiplatelet agents are therefore considered as a significant tool in the treatment and/or prevention of cardiovascular thrombotic disease (3-5). Antiplatelet agents such as aspirin (acetylsalicylic acid), clopidogrel or ticlopidine and anticoagulants such as warfarin are currently two predominant groups of orally consumable drugs in standard therapeutic


protocols for prophylaxis and treatment of venous thrombosis and reducing the risk of recurrent myocardial infarction (6-8).

Currently aspirin, which irreversibly inhibits cyclooxygenase I-mediated transformation of arachidonic acid to thromboxane A<sub>2</sub> (TXA<sub>2</sub>), and the P<sub>2</sub>Y<sub>12</sub> antagonists clopidogrel and prasugrel, which selectively and irreversibly bind to the P<sub>2</sub>Y<sub>12</sub> ADP receptor are routinely used as antiplatelet agents (9, 10).

However there are still some serious limitations to these agents which include weak inhibition of platelet function (*eg.* aspirin) (11), slow onset of action (*eg.* clopidogrel) (12), variable response to treatment among the patients (*eg.* clopidogrel and aspirin) (11, 12) and high incidence of bleeding events which occur in both aspirin and clopidogrel drug therapy (13, 14).

\* Corresponding author:

E-mail: farzadkf@yahoo.com



**Figure 1.** Active antiplatelet pyrimidine derivatives. A. 6-alkylamino-2,4-dialkyl(aryl)thiopyrimidines [7], R = Me, *n*-Pr, R1 = *n*-Pr, *i*-Pr, Ph, Bn, R2 = Bn, CH<sub>2</sub>CH<sub>2</sub>Ph; B. Tricyclic pyrimidines [8], R = some amino substituents; c. Substituted 6-amino-2-mercaptopyrimidines [9], R = CH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>NHBOC, CH<sub>2</sub>CH<sub>2</sub>OH, CH<sub>2</sub>CF<sub>3</sub>, 2-Chlorobenzyl, *etc.* R1 = substituted arylsulfonyl moieties.


Considering the current situation, development of novel antiplatelet agents which are safe and effective is an urgent need (15).

Aminopyrimidine derivatives are an interesting group of compounds with various reported biological properties. Pyrimidine ring can be found in the structures of many important drugs such as nucleoside antibiotics, antibacterials and cardiovascular agents (16, 17).


Based on the hypothesis suggested by Cattaneo *et al.* (18), amino pyrimidine ring could be considered as a simplified form of the active metabolites of the thienopyridines and ATP derivatives. The active metabolites of thienopyridines have a simple monocyclic

structure which implies that the presence of a bicyclic structure like that of a purine ring is not an absolute requirement for the affinity for the ADP receptor or so on platelet membrane.

A group of amino pyrimidine derivative with thioether substituents has been synthesized and evaluated by Cattaneo *et al.* The compound showed satisfactory anti platelet aggregation effects when ADP had been used for aggregation induction (18). Based on the mentioned reports and in order to investigate the capability of amino pyrimidine derivatives in inhibition of platelet aggregation pathways, we synthesized two groups of amino pyrimidines with the following structures:



**Figure 2.** Comparing the structures of aminopyrimidines with the structures of purine and the active metabolite of clopidogrel.



**Figure 3.** Chemical structure of the synthesized compounds.

Comparing the activity of these compounds in inhibition of platelet aggregation induced by ADP and arachidonic acid will provide some insights into the structure activity relationship of these compounds.

#### Chemistry

The synthetic procedures for groups I and II are illustrated in Figures 4 and 5.

Group I: Methyl ketones (1a-h) were allowed to react with dimethylformamide-dimethylacetate (DMF-DMA) to produce 3-(dimethylamino)-1-aryl-2-en-1-ones (2a-h). These intermediates could be then condensed with guanidine HCl to obtain the corresponding amino pyrimidine ring systems (19).


Group II: As it is shown in Figure 5 the intermediate II<sub>3</sub> (4,6-diaminopyrimidine-2-thiol) was obtained by the reaction of thiourea and malononitrile in absolute ethanol as the solvent. Subsequent reaction of compound II<sub>3</sub> with various alkylhalides at room temperature afforded compounds II<sub>4-25</sub> in good yields.

Structure confirmation of the synthesized intermediates and the final products was performed using IR, NMR and Mass spectrometry (20).


#### Experimental

##### General

All evaporations were carried out in vacuo with a rotary evaporator. Melting points (°C) were determined by capillary method on an electrothermal melting point apparatus. Infrared spectra were recorded as thin films of KBr plates with  $U_{\max}$  in inverse centimeters. Nuclear magnetic resonance spectra for proton (<sup>1</sup>H NMR) were recorded on a Bruker DRX-Avance (500 MHz) spectrometer. Chemical shift values are expressed in ppm (parts per million) relative to tetramethylsilane (TMS) as internal standard; s: singlet, d: doublet, dd: double doublet, t: triplet, q: quartet, m: multiplet, br s: broad singlet. Thin layer chromatography (TLC) was performed on Whatman SilG/UV<sub>254</sub> silica gel plates with fluorescent indicator and the



**Figure 4.** Compounds (I<sub>a-h</sub>) and (I<sub>A-H</sub>) synthesis scheme. Reagent and conditions: (a) DMF, reflux, 24 h; (b) NaOCH<sub>3</sub>, Isopropanol, reflux, 48 h.



**Figure 5.** Compounds ( $\text{II}_{4-25}$ ) synthesis scheme. Reagents and conditions: (a) EtONa, reflux, 3 h; (b) NaOH 0.1 M,  $\text{CH}_3\text{OH}$ ,  $\text{R-CH}_2\text{-X}$  (X = Cl, Br), rt, 18 h10).

spots were visualized under 254 and 366 nm illumination. Mass analyses were performed on an Agilent 6400 series equipped with an electrospray (ESI) ionization interface (drying gas adjusted at 300 °C, nebulizing gas flow at 12 L/min). All the compounds were analyzed for C, H, N and S on a Costech model 4010 and agreed with the proposed structures within  $\pm 0.4\%$  of the theoretical values.

#### General procedure for the preparation of compounds ( $\text{I}_{a-h}$ )

The detailed description of the methods used for preparation of compounds  $\text{I}_a$  to  $\text{I}_h$  and compounds  $\text{I}_A$  to  $\text{I}_H$  is reported in reference 19. Briefly to a solution of acetophenone (84 mmol) in DMF (16 mL), was added DMF-DMA (84 mmol) and the solution was heated under reflux for 24 h. Brine (25 mL) was added to the reaction mixture after cooling and the reaction mixture was then extracted with  $\text{CH}_2\text{Cl}_2$  ( $3 \times 25$  mL). The combined organic fractions were dried over anhydrous  $\text{Na}_2\text{SO}_4$  and the solvent was evaporated under reduced pressure. The residue was dissolved in EtOAc (12 mL), followed by addition of n-hexane (100 mL). The precipitate thus obtained was filtered and dried to give the pure product  $\text{I}_a$  as


a yellow solid (19).

#### General procedure for the preparation of compounds ( $\text{I}_{A-H}$ )

To a solution of compounds  $\text{I}_a$  (2.68 mmol) in isopropanol (13.5 mL) were added sodium methoxide (10.7 mmol) and guanidine hydrochloride (4.02 mmol) and the solution was heated under reflux for 48h. Distilled water (25 mL) was added to the reaction mixture after cooling and the mixture was then extracted with EtOAc ( $3 \times 15$  mL). The combined organic fractions were dried over anhydrous  $\text{Na}_2\text{SO}_4$  and the solvent was evaporated under reduced pressure. The residue was dissolved in EtOAc (2 mL), followed by addition of n-hexane (25 mL). The precipitate thus obtained was filtered and dried to give  $\text{I}_{A-H}$ .

#### Representative procedure for the preparation of compounds ( $\text{II}_{4-25}$ )

The detailed description of the methods used for preparation of compounds  $\text{II}_4$  to  $\text{II}_{25}$  is reported in reference 20. Briefly, to a solution of 4,6-diaminopyrimidine-2-thiol ( $\text{II}_3$ , 1.4 mmol) in methanol, was added alkyl halide (3.5 mmol) under basic conditions (NaOH 0.1 M, 14 mL). The mixture was then stirred for 18h at

**Table 1.** Antiplatelet activity of group I derivatives.

Compound	R	A.A IC <sub>50</sub> MM	ADP %inhibition
Ia (1.25 mM)	ph	36.75	45.2
I <sub>A</sub> (0.75 mM)	ph	544	29
Ib (1 mM)	4-F- ph	72.4	28.6
I <sub>B</sub> (2.5 mM)	4-F- ph	62.5	53
I <sub>c</sub> (1 mM)	3-thienyl	1000	35.5
I <sub>C</sub> (1.6 mM)	3-thienyl	>1000	57.3
I <sub>d</sub> (1.3 mM)	4-(CF <sub>3</sub> )- ph	>1000	24.3
I <sub>D</sub> (0.5 mM)	4-(CF <sub>3</sub> )- ph	340	42.5
Ie (1.3 mM)	4-( ph)- ph	>1000	24.3
I <sub>E</sub> (1 mM)	4-( ph)- ph	>1000	50
If (1 mM)	4-pyridyl	>1000	1
I <sub>F</sub> (1 mM)	4-pyridyl	ND	ND
Ig (2.5 mM)	3-pyridyl	1000	35
I <sub>G</sub> (0.5 mM)	3-pyridyl	192	39.5
I <sub>h</sub> (1.25 mM)	2-pyridyl	752	27.1
I <sub>H</sub> (1.25 mM)	2-pyridyl	>1000	44
Indomethacin		3.0	
Aspirin		30.3	


room temperature. After removing the solvent under reduced pressure, the residue was washed by water and the precipitate was collected as a solid. All compounds were obtained in acceptable purity and no further purification was needed (20).

### Results and Discussion

All the synthesized compounds were screened for their effects on human platelet aggregation induced by arachidonic acid and ADP using light transmission aggregometry. IC<sub>50</sub> was determined as the concentration of the test compounds that exhibit platelet aggregation

by 50%. The *in-vitro* antiplatelet activity of the synthesized derivatives is listed in Tables 1 and 2.

Comparing the activities of the two aminopyrimidine groups indicates that none of the compounds showed satisfactory activity against the aggregation induced by ADP. Therefore it could be concluded that the compounds do not interfere with ADP receptors on platelet membrane. However, acceptable activities were observed in both groups against the aggregation induced by arachidonic acid. This is not in agreement with the report by Cattaneo *et al.* who introduced a group of aminopyrimidines as active platelet aggregation

**Table 2.** Antiplatelet activity of group II derivatives.

Compound (In 1 mM)	R	A.A IC <sub>50</sub> MM	ADP % inhibition
II <sub>4</sub>	H	>1000	10
II <sub>5</sub>	Methyl	>1000	33.8
II <sub>6</sub>	Ethyl	500	42
II <sub>7</sub>	Propyl	ND	ND
II <sub>8</sub>	Butyl	500	42
II <sub>9</sub>	Pentyl	>1000	53
II <sub>10</sub>	Cyclobutyl	>1000	49
II <sub>11</sub>	Morpholinomethyl	>1000	ND
II <sub>12</sub>	2-methylphenyl	213	38.3
II <sub>13</sub>	3-methylphenyl	220	10
II <sub>14</sub>	2-nitrophenyl	209	20
II <sub>15</sub>	2-chlorophenyl	>1000	14.7
II <sub>16</sub>	2-fluorophenyl	<b>80</b>	23
II <sub>17</sub>	3-fluorophenyl	700	49
II <sub>18</sub>	4-fluorophenyl	214	24.3
II <sub>19</sub>	3-methoxyphenyl	>1000	8
II <sub>20</sub>	4-cyanophenyl	>1000	28
II <sub>21</sub>	3-(trifluoromethyl)phenyl	>1000	50
II <sub>22</sub>	Benzyl	ND	ND
II <sub>23</sub>	2-phenethyl	760	70
II <sub>24</sub>	1-naphthalenyl	1000	51.9
II <sub>25</sub>	1-(1-methylpiperidin-2-yl)methyl	>1000	95
Indomethacin		3.0	
aspirin		30.3	

inhibitors which interfere with ADP receptors.

Comparing the overall results obtained for aminopyrimidines I and II indicates that 2-aminopyrimidines (I) were more active than 4,6-diaminopyrimidines (II). Only compound 16 in group II showed satisfactory IC<sub>50</sub> (80 μM).

Among the 2-aminopyrimidines group (I), on the other hand, a few compounds (I<sub>a</sub>, I<sub>b</sub>, I<sub>B</sub> and I<sub>G</sub>) showed good activities (36.75, 72.4,

62.5 and 192 μM). Interestingly, compounds with fluorine substituent on phenyl ring (I<sub>b</sub>, I<sub>B</sub>) were among the most active compounds.

Global physicochemical properties for compounds I<sub>a-h</sub>, I<sub>A-H</sub> and II<sub>4-25</sub> were calculated using Chemdraw Ultra, Chem3D Ultra version 8.0 and Hyper Chem professional and the results are presented in Tables 3 and 4.

Efforts to find a relationship between these

**Table 3.** Global physicochemical properties for compounds group I.

Compound	ClogP <sup>A</sup>	P <sup>B</sup>	V <sup>C</sup>	SA <sup>D</sup>	D <sup>E</sup>
Ia	2.2974	20.85	606.6	388.66	2.669
I <sub>A</sub>	1.774	20.03	549.5	259.2	1.319
Ib	2.516	24.43	632.5	398.5	3.181
I <sub>B</sub>	1.926	22.06	553.45	357.2	1.122
I <sub>c</sub>	2.0739	20.38	581.23	379.5	2.56
I <sub>C</sub>	1.4697	19.55	521.3	342.54	0.778
I <sub>d</sub>	3.3127	22.42	683.67	430.8	3.844
I <sub>D</sub>	2.6730	21.59	620.86	395.87	2.208
Ie	4.185	30.5	811.04	495.06	2.708
I <sub>E</sub>	3.66	29.69	747.46	458.7	1.548
If	1.1964	20.14	592.2	382.9	3.2
I <sub>F</sub>	0.409	19.32	529.56	344.89	1.194
Ig	1.1964	20.14	591.97	383.02	1.283
I <sub>G</sub>	0.1997	19.32	529.45	345.85	1.944
I <sub>h</sub>	1.5964	22.95	611.69	389.11	1.914
I <sub>H</sub>	0.409	19.32	535.7	351.3	1.3

<sup>A</sup>ClogP were calculated by using Chem Draw Ultra version 8.0. <sup>B</sup>Polarizability values were calculated by using Hyper Chem Professional. <sup>C</sup>Molecular volume values were calculated by using Hyper Chem Professional. <sup>D</sup>Surface area values were calculated by using Hyper Chem Professional. <sup>E</sup>Dipole (debye) values were calculated by using Chem3D Ultra version 8.0.

**Table 4.** Global physicochemical properties for compounds group II.

Compound	ClogP <sup>A</sup>	P <sup>B</sup>	V <sup>C</sup>	SA <sup>D</sup>	D <sup>E</sup>
II <sub>4</sub>	0.866	16.55	487.68	331.52	3.1
II <sub>5</sub>	1.395	18.39	542.6	362.6	2.926
II <sub>6</sub>	1.924	20.22	596.38	394.4	2.933
II <sub>7</sub>	2.453	22.06	650.7	423.37	2.895
II <sub>8</sub>	2.982	23.89	703.83	455.17	2.9047
II <sub>9</sub>	3.511	25.73	749.73	477.11	2.884
II <sub>10</sub>	2.398	23.12	653.14	414.09	2.898
II <sub>11</sub>	0.8381	26.94	745.2	461.24	3.168
II <sub>12</sub>	2.883	28.05	744.66	467.25	3.084
II <sub>13</sub>	2.933	28.05	754.32	472.14	2.902
II <sub>14</sub>	2.097	32.61	767.72	477.94	4.137
II <sub>15</sub>	3.147	28.14	729.07	460.65	41.994
II <sub>16</sub>	2.577	26.12	706.53	448.67	1.818
II <sub>17</sub>	2.577	26.12	711.03	452.9	3.835
II <sub>18</sub>	2.577	26.12	711.7	453.5	4.995
II <sub>19</sub>	2.353	28.68	771.85	485.4	1.774
II <sub>20</sub>	1.867	28.06	760.45	479.2	6.326
II <sub>21</sub>	3.317	27.77	779.18	489.7	4.5
II <sub>22</sub>	2.963	28.05	754.7	477.364	3.067
II <sub>23</sub>	3.342	29.88	809.23	507.6	3.145
II <sub>24</sub>	3.608	33.48	826.6	509.25	2.885
II <sub>25</sub>	2.397	29.47	799.19	477.84	1.924

<sup>A</sup>ClogP were calculated by using Chem Draw Ultra version 8.0. <sup>B</sup>Polarizability values were calculated by using Hyper Chem Professional. <sup>C</sup>Molecular volume values were calculated by using Hyper Chem Professional. <sup>D</sup>Surface area values were calculated by using Hyper Chem Professional. <sup>E</sup>Dipole (debye) values were calculated by using Chem3D Ultra version 8.0.

physicochemical parameters and anti platelet aggregation activity of the compounds did not result in a clear correlation. This could be due to the fact that the compounds are different in their access to their targets in the platelet aggregation pathway induced by arachidonic acid.

Further mechanistic studies are needed to clarify the mechanism of antiplatelet activity of these compounds.

### References

- (1) Zia-Ul-Haq M, Ahmad Shahid S, Ahmed S, Ahmad Sh, Qayum M and Khan I. Antiplatelet activity of methanolic extract of *Grewiaasiatica* L. leaves and *Terminallachebula* Retz Fruits. *J. Med. Plants Res.* (2012) 6: 2029-2032.
- (2) Mathers C, Fat DM and Boerma JT. World Health Organization. *The Global Burden of Diseases: 2004 Update*. Geneva (2008).
- (3) White HD. Oral antiplatelet therapy for atherothrombotic disease: Current evidence and new directions. *Am. Heart J.* (2011) 161: 450-461.
- (4) Latib A, Ielasi A, Ferri L, Chieffo A, Godino C, Carlino M, Montorfano M and Colombo A. Aspirin intolerance and the need for dual antiplatelet therapy after stent implantation: A proposed alternative regimen. *Int. J. Cardiol.* (2013) 165: 444-447.
- (5) Reiter RA and Jilma B. Platelets and new antiplatelet drugs. *Therapy* (2005) 2: 465-502.
- (6) Ilas J, Jakopin Z, Borstnar T, Stegnar M and Kikelj D. 3,4-Dihydro-2H-1,4-benzoxazine derivatives combining thrombin inhibitory and glycoprotein IIb/IIIa receptor antagonistic activity as a novel class of antithrombotic compounds with dual function. *J. Med. Chem.* (2008) 51: 5617-5629.
- (7) De Meyer SF, Vanhoorelbeke K, Broos K, Salles II and Deckmyn H. Antiplatelet drugs. *Br. J. Haematol.* (2008) 142: 515-528.
- (8) Amidi S, Kobarfard F, Moghaddam AB, Tabib K and Soleymani Z. Electrochemical synthesis of novel 1,3-indandione derivatives and evaluation of their antiplatelet aggregation activities. *Iran. J. Pharm. Res.* (2013) 12: 91-103.
- (9) Maree AO and Fitzgerald DJ. Variable platelet response to aspirin and clopidogrel in atherothrombotic disease. *Circulation* (2007) 115: 2196-2207.
- (10) Meadows TA and Bhatt DL. Clinical aspects of platelet inhibitors and thrombus formation. *Circ. Res.* (2007) 100: 1261-1275.
- (11) Patrono C, Collier B, Fitz Gerald GA, Hirsh J and Roth G. Platelet-active drugs: the relationships among dose, effectiveness, and side effects. *Chest.* (2004) 126: 234-264.
- (12) Bassand JP. Unmet needs in antiplatelet therapy. *Eur. Heart J.* (2008) 10: 3-11.
- (13) Guthrie R. Review and management of side effects associated with antiplatelet therapy for prevention of recurrent cerebrovascular events. *Adv. Ther.* (2011) 28: 473-482.
- (14) Chan YC, Valenti D, Mansfield AO and Stansby G. Warfarin induced skin necrosis. *Br. J. Surg.* (2000) 87: 266-272.
- (15) Liu G, Xu J, Chen N, Zhang S, Ding Z and Du H. Synthesis of N6-alkyl(aryl)-2-alkyl(aryl) thioadenosines as antiplatelet agents. *Eur. J. Med. Chem.* (2012) 53: 114-123.
- (16) Tozkoparan B, Ertan M, Kelicen P and Demirdamar R. Synthesis and anti-inflammatory activities of some thiazolo[3,2-a]pyrimidine derivatives. *Farmacol.* (1999) 54: 588-593.
- (17) Clark J, Shahhet MJ, Korakas D and Varvounis G. Synthesis of thieno[2,3-d]pyrimidines from 4,6-dichloropyrimidine-5-carbaldehydes. *J. Heterocyclic Chem.* (1993) 30: 1065-1072.
- (18) Crepaldia P, Cacciari B, Bonache MC, Spalluto G, Varani K, Borea PA, Kugelen IV, Hoffman K, Pugliano M, Razzari C and Cattaneo M. 6-amino-2-mercapto-3H-pyrimine-4-one derivatives as new candidates for the antagonism at the P2Y<sub>12</sub> receptors. *Bioorg. Med. Chem. J.* (2009) 17: 4612-4621.
- (19) Mohebbi Sh, Shirazi FH, Sharifnia SH and Kobarfard F. Introducing synthesis route-based hit identification approach as a tool in medicinal chemistry and its application in investigating the antiproliferative and antimicrobial effects of 2-aminopyrimidine derivatives. *Int. J. Drug Discov.* (2011) 3: 78-87.
- (20) Mohebbi Sh, Falcón-Pérez JM, González E, Millet O, Mato JM and Kobarfard F. Synthesis, dihydrofolate reductase inhibition, anti-proliferative testing and saturation transfer difference <sup>1</sup>H-NMR study of some new 2-substituted-4,6-diaminopyrimidine derivatives. *Chem. Pharm. Bull.* (2012) 60: 70-78.

---

This article is available online at <http://www.ijpr.ir>