

Radioprotective Effect of Thymol Against Salivary Glands Dysfunction Induced by Ionizing Radiation in Rats

Seyed Mohammad Abedi^a, Fateme Yarmand^{b,c}, Mina Motalebnejad^{b,c}, Maryam Seyedmajidi^d, Dariush Moslemi^e, Ali Bijani^f and Seyed Jalal Hosseinimehr^{g*}

^aDepartment of Radiology, Faculty of Medicine, Mazandaran University of Medical Sciences, Sari, Iran. ^bDepartment of Oral Medicine, Faculty of Dentistry, Babol University of Medical Science, Babol, Iran. ^cCellular and Biology Research Center, Babol University of Medical Science, Babol, Iran. ^dDental Materials Research Center, Faculty of Dentistry, Babol University of Medical Science, Babol, Iran. ^eDepartment of Radiotherapy, Faculty of Medicine, Babol University of Medical Science, Babol, Iran. ^fDepartment of Physiology and Cellular and Molecular Biology Research Center, Babol University of Medical Science, Babol, Iran. ^gDepartment of Radiopharmacy, Faculty of Pharmacy, Mazandaran University of Medical Sciences, Sari, Iran.

Abstract

The aim of this study was to investigate the radioprotective effect of thymol as a natural product against salivary glands dysfunction induced by ionizing radiation in rats. The rats were treated with thymol at dose of 50 mg/Kg before exposure to ionizing radiation at dose 15 Gy. Salivary gland function was evaluated with radioisotope scintigraphy and then salivary gland to background counts ratio was calculated. Ionizing radiation caused significant salivary glands dysfunction at the 3th and the 70th days with reduction in radioactivity uptake in salivary glands. Ratios of salivary gland to background radioactivities were 2.0 ± 0.05 , 1.58 ± 0.62 and 1.99 ± 0.07 at 3th days for control, radiation, and thymol plus radiation groups, respectively. Thymol significantly protected acute and chronic salivary gland dysfunction induced by ionizing radiation in the rats. This finding may have been a promising application of thymol for the protection of salivary glands dysfunction induced by ionizing irradiation in patients exposed to radiation in head and neck cancer therapy.

Keywords: Thymol; Radioprotective; Salivary gland dysfunction; Scintigraphy; Radiotherapy.

Introduction

Radiotherapy of head and neck cancer can cause severe adverse effects on oral cavity such as xerostomia and salivary gland dysfunction which impair quality life of patients undergoing radiation therapy (1-3). Ionizing radiation

produces reactive oxygen species (ROS) and other toxic substances, which interact with critical macromolecules such as DNA, and leads to serious cellular dysfunction and death (4, 5). However, ionizing radiation (IR) is focusing on tumor tissue for killing cancerous cells; it may also cause side effects on the normal organs (6). The salivary glands are organs that are unwantedly exposed to radiation in the head and neck region. However, salivary glands are

* Corresponding author:

E-mail: sjhosseinim@yahoo.com

considered to be radio-resistance due to matured and differentiated cells; exposure to high dose of ionizing radiation such as head and neck radiotherapy or radioiodine therapy affects their functions (7-9). IR causes major changes on salivary gland and leads to salivary glands dysfunctions (10, 11). The main complications related to salivary gland dysfunction are including xerostomia, difficulty in swallowing and speaking, taste changes and high dental caries risk, which affect the patient's quality of life (12, 13). Also salivary gland dysfunction is the most common side effect of oral β -emitting ^{131}I for the treatment of well-differentiated thyroid carcinoma (DTC). The severity of salivary gland dysfunction increases with increase dosage of ^{131}I radioiodine administration. Fallahi *et al.* reported that consumption of vitamin E may be associated with a significant protective effect against radiation-induced dysfunction in salivary glands following single-dose ^{131}I therapy in patients with differentiated thyroid cancer (14). It is important to protect salivary glands against radiation-toxicity and also improves quality life of patients after radiotherapy. Antioxidants can scavenge free radicals and toxic substances produced by IR and have a beneficial role in protection of cellular macromolecules against toxicity induced by IR (15). Thymol is a natural phenolic compound that presents in various plants, such as thyme (Lamiaceae) and Zataria (16-18). Thymol has several biological properties such as anti-inflammatory activity (19) and protective effects against toxicity induced by oxidative stress in liver and lymphocytes (20-22). Thymol acts as an antioxidant, free radical scavenging and anti-lipid peroxidation, which can protect cell against free radicals (20, 23). With respect to these protective mechanisms, it is possible thymol protects salivary glands dysfunction caused by IR. Salivary scintigraphy is a useful technique to evaluate objectively the salivary gland function of patients with head and neck irradiated tumors. Salivary gland scintigraphy with $\text{Na}^{99\text{m}}\text{TcO}_4$ is a well-established procedure for the evaluation of human salivary gland function (24, 25). There is a good correlation between the scintigraphic findings and the measured salivary flow rates in evaluation of salivary gland function (26).

The aim of this study was to evaluate

the radioprotective effects of thymol on the radiation-induced damages on salivary glands in the irradiated rats. This protective effect was evaluated by salivary gland scintigraphy with using $\text{Na}^{99\text{m}}\text{TcO}_4$.

Material and methods

Animal treatment

Male Wistar rats between 8 and 10 weeks old were used (Pasteur Institute, Iran). They were kept in polycarbonate cages under an alternating 12 h light-dark cycle in animal house of university. This study was approved by research and education deputy of Heart Center Hospital, Mazandaran University of Medical Sciences, Sari, Iran (5394). Animals were maintained on laboratory chow and water *ad libitum*. The animals were randomly divided into three groups of seven animals as control, radiation and thymol plus radiation. Control group (C) was received three doses of sesame oil given intraperitoneally 48, 24 and 1h; radiation group (IR) was treated with three doses sesame oil given intraperitoneally 48, 24 and 1h before irradiation. Thymol plus radiation group was received thymol at single dose 50 mg/Kg given intraperitoneally 48, 24 and 1h before the gamma irradiation.

Irradiation

Thirty minutes before irradiations, the animals were weighed and anesthetized by an intramuscular injection of ketamine chlorhydrate (0.1 mg/Kg) and xylazine (0.05 mg/Kg). Animals were placed on the table with a 3 mm lead to protect the body, so that only the head and neck regions were exposed. The animals were irradiated with a cobalt-60 gamma radiation source (Theratron 780, AECL, Ontario, Canada) in Babolsar radiotherapy hospital with a single exposure to 15 Gy of radiation.

Salivary glands function assessment

Salivary glands function assessments were performed in Heart Central Hospital (Sari, Iran) Ten minutes after IV injection (via tail's vein) of $\text{Na}^{99\text{m}}\text{TcO}_4$ ($^{99}\text{Mo}/^{99\text{m}}\text{Tc}$ generator, Pars isotope, Tehran, Iran), the rats placed on the gamma camera table for imaging. Salivary gland scintigraphy was acquired by a digital

Table 1. Salivary gland-to-background ratios of treated groups in scintigraphy assessment (n=7).

	Before Radiation Mean ± SD	3 days after radiation Mean ± SD	70 days after radiation Mean ± SD
Control	2.04±0.07 *	2±0.5	1.99±0.11
Radiation	2.03±0.04 *	1.58±0.62 **	1.58±0.08**
Thymol + radiation	2.12±0.1*	1.99±0.07 ***	2.01±0.1***

*Non-significant differences between control, radiation and thymol+radiation groups before radiation

**Significant difference between control and radiation groups

***Significant difference between thymol+radiation and radiation groups

gamma camera (dual head, Siemens e.cam, Germany) with a matrix of 256×256 pixels, and using a pinhole collimator with a zoom factor of 1.78. Imaging started 10 min after injection of $\text{Na}^{99\text{m}}\text{TcO}_4$ and was conducted up to a quantity of 300 kilocounts. Imaging process was taken about five min. The processing ESOFIT software was used on gamma camera. The uptake of the salivary glands was determined by a region of interest (27) technique. All ROIs were defined manually for each single rat and left and right gland region. The background activity was placed in the supraclavicular area. For the evaluation of glandular function, the ratio of the accumulation in the gland ROI to the accumulation in the background ROI, termed gland-to-background ratio was used. Static salivary gland scintigraphy scans were performed in three times; first scintigraphy was performed 7 days before the radiation or sham irradiation. The second and third scintigraphies were performed at 3 and 70 days after radiation or sham irradiation.

Statistical analysis

Experimental data were expressed as the mean ± standard deviation (SD). The results were compared with the control group, and statistical analysis was performed by independent t-test to determine the significance of the difference between groups. The differences were considered significant when the P value was <0.05.

Results

Rats were exposed to gamma ray at a single dose 15 Gy on the oral region. All rats were alive up to end of the study. Static salivary gland radioisotope scintigraphy was performed for assessment of any tissue dysfunction caused

by ionizing radiation in the rats. Scintigraphy scans of salivary glands are shown in Figure 1. Radioisotope uptake was quantified in salivary glands of rats. In this study, the target tissue was salivary glands and salivary gland-to-background ratios (T/B ratio) were determined in the treated groups with irradiation and/or thymol (Figure 1.). The T/B ratio is about two for rats before exposed to radiation. It was not observed any significant difference between the groups before the gamma radiation exposure in T/B ratios (Table 1).

Exposure to irradiation exhibited a significant salivary gland dysfunction that was shown a reducing in T/B ratio. IR reduced the uptake of radioisotope in salivary glands. The T/B ratios were 2 ± 0.5 and 1.58 ± 0.62 on the 3th day and 1.99 ± 0.11 and 1.58 ± 0.08 on the 70th day for control and IR groups, respectively (Table 1). Thymol increased the T/B ratio up to 1.99 ± 0.07 and 2.01 ± 0.4 in the irradiated rats treated with thymol compared to the gamma-irradiated rat alone on the 3th day and the 70th day. It is clear that thymol treatment increased uptake of $\text{Na}^{99\text{m}}\text{TcO}_4$ in salivary glands and it improved salivary gland hypofunction caused by irradiation.

Discussion

In this study, the radioprotective effect of thymol against salivary glands dysfunction induced by ionizing radiation was investigated using radioisotope scintigraphy scans. The salivary gland to background radioactivity counts ratio was calculated. The right and left salivary glands, and background ROI were used for the calculation of the target-to-background ratio (26). In static scintigraphy, an increased




Figure 1. Scintigraphy of salivary glands of rats treated with thymol (50 mg/Kg) and/or radiation on the before and 3th and 70th days after irradiation. Rats were treated with oil or thymol and then irradiated with gamma irradiation at dose of 15 Gy. A significant reduction in salivary glands uptake of radioactive was observed in irradiated rats as compared with control and thymol groups. A: salivary glands (T), B: background (B). T/B ratios were calculated with respect to radioactivity accumulation in salivary glands and background.

uptake of radioisotope was seen in the salivary glands in the Wistar rats. Thymol at three doses of 50 mg/Kg significantly protected the salivary glands hypofunction caused by ionizing radiation in rats. Thymol elevated uptake $\text{Na}^{99\text{m}}\text{TcO}_4$ in salivary glands that reduced by ionizing radiation. In this study, local mouth exposure to radiation caused an acute salivary glands dysfunction on the 3th day after the irradiation and the chronic organ impairment was continued up to 70th days without any improvement. Ionizing radiation caused a severe salivary glands damage, which was not improved at long time. Radiation induces apoptosis in salivary glands cells and causes salivary glands hypofunction (28, 29). Ionizing radiation caused a reduction of saliva secretion, salivary amylase activity, and superoxide dismutase (30). In this study, a high dose of 15 Gy of gamma rays

was used to test the properties of thymol as a radioprotector. It was previously demonstrated that high dose of ionizing radiation about 15 Gy starts to induce salivary gland dysfunction (31).

Our experiments showed that thymol protected acute and chronic side effects of irradiation on the salivary glands, which evaluated at the 3th and the 70th days after irradiation. Rats have a similar distribution of the sodium/iodide symporter with human, which suggests the suitability of this animal for scintigraphic studies of the salivary glands (32). Thymol as a monoterpene phenol compound has free radical scavenging and antioxidant properties. It directly scavenges free radicals produced by oxidative stress process in the cells as well as restored antioxidant capacities of cells through increasing of cell glutathione (33). Thymol suppressed radiation-induced genotoxicity, apoptosis, and necrosis in the

lung fibroblast cell. These protective effects are related to free radical scavenging and modulation of oxidative stress by thymol (34). Thymol pretreatment enhanced reduced intercellular defense enzymes such as glutathione, glutathione-S-transferase, catalase, and superoxide dismutase levels in mouse liver homogenates after exposure to radiation. Thymol treatment before exposure to ionizing radiation resulted in a significant increase in hematological parameters (21). Thymol can protect genotoxicity induced by ionizing radiation on human lymphocyte cells (35). The phenolic structure of thymol is playing a crucial role in absorbing and neutralizing free radicals (23, 36).

In conclusion, ionizing radiation caused salivary glands dysfunction in rats through a reduction of radioactivity uptake in salivary glands. This side effect was severe and continued for a long time on the 70th day after irradiation. Thymol at dose 50 mg/Kg significantly improved salivary gland dysfunction caused by ionizing radiation. Short and late side effects of radiation on the salivary glands were protected by thymol in rats. Several previous studies were reported radioprotective effects of thymol in various tissues in animal and as well as in vitro. Thymol is a promising radioprotective agent for patients who receive radiation in head and neck cancer therapy.

Conflict of interest statement

The authors declared no potential conflict of interest with respect to the authorship, and/or publication of this study.

References

- (1) Satheeshkumar PS, Chamba MS, Balan A, Sreelatha KT, Bhatathiri VN and Bose T. Effectiveness of triclosan in the management of radiation-induced oral mucositis: a randomized clinical trial. *J. Cancer Res. Ther.* (2010) 6: 466-72.
- (2) Simcock R, Fallowfield L, Monson K, Solis-Trapala I, Parlour L, Langridge C and Jenkins V. ARIX: a randomised trial of acupuncture v oral care sessions in patients with chronic xerostomia following treatment of head and neck cancer. *Ann. Oncol.* (2013) 24: 776-83.
- (3) Rodrigues NA, Killion L, Hickey G, Silver B, Martin C, Stevenson MA, Mauch PM and Ng AK. A prospective study of salivary gland function in lymphoma patients receiving head and neck irradiation. *Int. J. Radiat. Oncol. Biol. Phys.* (2009) 75: 1079-83.
- (4) Patni N, Patni S and Bapna A. The optimal use of granulocyte macrophage colony stimulating factor in radiation induced mucositis in head and neck squamous cell carcinoma. *J. Cancer Res. Ther.* (2005) 1: 136-41.
- (5) Hosseinimehr SJ. Flavonoids and genomic instability induced by ionizing radiation. *Drug Discov. Today* (2010) 15: 907-18.
- (6) Liu LN, Shi L, Li SC, Zhang WJ, Zhang Y and Zhang ZP. Protective Role of Rheum Tanguticum Polysaccharide 1 in Radiation- induced Intestinal Mucosal Injury. *Iran. J. Pharm. Res.* (2015) 14: 833-41.
- (7) Coppes RP, Zeilstra LJ, Vissink A and Konings AW. Sialogogue-related radioprotection of salivary gland function: the degranulation concept revisited. *Radiat. Res.* (1997) 148: 240-7.
- (8) Nagler RM, Baum BJ and Fox PC. A 2 week paired study of early X-irradiation effects on rat major salivary gland function. *Arch. Oral Biol.* (1996) 41: 713-7.
- (9) Noaparast Z and Hosseinimehr SJ. Radioprotective agents for the prevention of side effects induced by radioiodine-131 therapy. *Future Oncology* (2013) 9: 1145-59.
- (10) Langendijk JA, Doornaert P, Verdonck-de Leeuw IM, Leemans CR, Aaronson NK and Slotman BJ. Impact of late treatment-related toxicity on quality of life among patients with head and neck cancer treated with radiotherapy. *J. Clin. Oncol.* (2008) 26: 3770-6.
- (11) Graff P, Lapeyre M, Desandes E, Ortholan C, Bensadoun RJ, Alfonsi M, Maingon P, Giraud P, Bourhis J, Marchesi V, Mege A and Peiffert D. Impact of intensity-modulated radiotherapy on health-related quality of life for head and neck cancer patients: matched-pair comparison with conventional radiotherapy. *Int. J. Radiat. Oncol. Biol. Phys.* (2007) 67: 1309-17.
- (12) Takagi K, Yamaguchi K, Sakurai T, Asari T, Hashimoto K and Terakawa S. Secretion of saliva in X-irradiated rat submandibular glands. *Radiat. Res.* (2003) 159: 351-60.
- (13) Franzen L, Funegard U, Ericson T and Henriksson R. Parotid gland function during and following radiotherapy of malignancies in the head and neck. A consecutive study of salivary flow and patient discomfort. *Eur. J. Cancer* (1992) 28: 457-62.
- (14) Fallahi B, Beiki D, Abedi SM, Saghari M, Fard-Esfahani A, Akhzari F, Mokarami B and Eftekhari M. Does vitamin E protect salivary glands from I-131 radiation damage in patients with thyroid cancer? *Nucl. Med. Commun.* (2013) 34: 777-86.
- (15) Hosseinimehr SJ. Trends in the development of radioprotective agents. *Drug Discov. Today* (2007) 12: 794-805.
- (16) Sajed H, Sahebkar A and Iranshahi M. *Zataria multiflora* Boiss. (Shirazi thyme)--an ancient condiment with modern pharmaceutical uses. *J. Ethnopharmacol.* (2013) 145: 686-98.

- (17) Schmidt E, Wanner J, Hiiferl M, Jirovetz L, Buchbauer G, Gochev V, Girova T, Stoyanova A and Geissler M. Chemical composition, olfactory analysis and antibacterial activity of *Thymus vulgaris* chemotypes geraniol, 4-thujanol/terpinen-4-ol, thymol and linalool cultivated in southern France. *Nat. Prod. Commun.* (2012) 7: 1095-8.
- (18) Hosseinimehr SJ, Mahmoudzadeh A, Ahmadi A, Ashrafi SA, Shafaghatahi N and Hedayati N. The Radioprotective effect of *Zataria multiflora* against genotoxicity induced by γ irradiation in human blood lymphocytes. *Cancer Biother. Radiopharm.* (2011) 26: 325-9.
- (19) Azuma Y, Ozasa N, Ueda Y and Takagi N. Pharmacological studies on the anti-inflammatory action of phenolic compounds. *J. Dent. Res.* (1986) 65: 53-6.
- (20) Alam K, Nagi MN, Badary OA, Al-Shabanah OA, Al-Rikabi AC and Al-Bekairi AM. The protective action of thymol against carbon tetrachloride hepatotoxicity in mice. *Pharmacol. Res.* (1999) 40: 159-63.
- (21) P RA, Nageshwar Rao B and Satish Rao BS. In vivo radioprotective potential of thymol, a monoterpene phenol derivative of cymene. *Mutat. Res.* (2011) 726: 136-45.
- (22) Al-Malki AL. Antioxidant Properties of Thymol and Butylated Hydroxytoluene in Carbon Tetrachloride-Induced Mice Liver Injury. *JKAU: Sci.* (2010) 22: 239-48.
- (23) Kruk I, Michalska T, Lichszeld K, Kladna A and Aboul-Enein HY. The effect of thymol and its derivatives on reactions generating reactive oxygen species. *Chemosphere* (2000) 41: 1059-64.
- (24) Mateos JJ, Setoain X, Ferre J, Roviroša A, Navalpotro B, Martin F, Ortega M, Lomena F, Fuster D, Pavia J and Pons F. Salivary scintigraphy for assessing the protective effect of pilocarpine in head and neck irradiated tumours. *Nucl. Med. Commun.* (2001) 22: 651-6.
- (25) van Acker F, Flamen P, Lambin P, Maes A, Kutcher GJ, Weltens C, Hermans R, Baetens J, Dupont P, Rijnders A, van den Bogaert W and Mortelmans L. The utility of SPECT in determining the relationship between radiation dose and salivary gland dysfunction after radiotherapy. *Nucl. Med. Commun.* (2001) 22: 225-31.
- (26) Pfestroff A, Muller F, Librizzi D, Eivazi B, Behe M, Hoefken H, Behr TM and Teymoortash A. Scintigraphic assessment of salivary gland function in a rat model. *In Vivo* (2010) 24: 681-5.
- (27) Radfar L and Sirois DA. Structural and functional injury in minipig salivary glands following fractionated exposure to 70 Gy of ionizing radiation: an animal model for human radiation-induced salivary gland injury. *Oral Surg. Oral Med. Oral Pathol. Oral Radiol. Endod.* (2003) 96: 267-74.
- (28) Limesand KH, Said S and Anderson SM. Suppression of radiation-induced salivary gland dysfunction by IGF-1. *PLoS One* (2009) 4: e4663.
- (29) Bralic M, Muhvic-Urek M, Stemberga V, Golemac M, Jurkovic S, Borcic J, Braut A and Tomac J. Cell death and cell proliferation in mouse submandibular gland during early post-irradiation phase. *Acta. Med. Okayama.* (2005) 59: 153-9.
- (30) Xu L, Yang X, Cai J, Ma J, Cheng H, Zhao K, Yang L, Cao Y, Qin Q, Zhang C, Zhang Q and Sun X. Resveratrol attenuates radiation-induced salivary gland dysfunction in mice. *Laryngoscope* (2013) 123: E23-9.
- (31) Vissink A, s-Gravenmade EJ, Ligeon EE and Konings WT. A functional and chemical study of radiation effects on rat parotid and submandibular/sublingual glands. *Radiat. Res.* (1990) 124: 259-65.
- (32) Abok K, Brunk U, Jung B and Ericsson J. Morphologic and histochemical studies on the differing radiosensitivity of ductular and acinar cells of the rat submandibular gland. *Virchows Arch. B Cell Pathol. Incl. Mol. Pathol.* (1984) 45: 443-60.
- (33) Kim YS, Hwang JW, Kang SH, Kim EH, Jeon YJ, Jeong JH, Kim HR, Moon SH, Jeon BT and Park PJ. Thymol from *Thymus quinquecostatus* Celak. protects against tert-butyl hydroperoxide-induced oxidative stress in Chang cells. *J. Nat. Med.* (2014) 68: 154-62.
- (34) Archana PR, Nageshwar Rao B and Satish Rao BS. Modulation of gamma ray-induced genotoxic effect by thymol, a monoterpene phenol derivative of cymene. *Integr. Cancer Ther.* (2011) 10: 374-83.
- (35) Hosseinimehr SJ, Mahmoudzadeh A, Ahmadi A, Ashrafi SA, Shafaghatahi N and Hedayati N. The radioprotective effect of *Zataria multiflora* against genotoxicity induced by gamma irradiation in human blood lymphocytes. *Cancer Biother. Radiopharm.* (2011) 26: 325-9.
- (36) Braga PC, Dal Sasso M, Culici M, Galastri L, Marceca MT and Guffanti EE. Antioxidant potential of thymol determined by chemiluminescence inhibition in human neutrophils and cell-free systems. *Pharmacology* (2006) 76: 61-8.