Original Article

Cytotoxic Properties of Three Isolated Coumarin-hemiterpene Ether Derivatives from *Artemisia armeniaca* Lam.

Mahdi Mojarrab^a, Seyed Ahmad Emami^b, Abbas Delazar^d and Zahra Tayarani-Najaran^{c*}

^aPharmaceutical Sciences Research Center, School of Pharmacy, Kermanshah University of Medical Sciences, Kermanshah, Iran. ^bDepartment of Pharmacognosy, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran. ^cDepartment of Pharmacognosy, School of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran. ^dDepartment of Pharmacodynamics and Toxicology, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran.

Abstract

Considering multiple reports on cytotoxic activity of the *Artemisia* genus and its phytochemicals, in the current study *A. armeniaca* Lam. and the three components isolated from the plant were subjected to cytotoxic studies. Analytical fractionation of *A. armeniaca* aerial parts for the first time was directed to the isolation of 7-hydroxy-8-(4-hydroxy-3-methylbutoxy) comarin (armenin), 8-hydroxy-7-(4-hydroxy-3-methylbutoxy) comarin (isoarmenin) and deoxylacarol. Cytotoxicity assessed with alamalBlue® assay and apoptosis was detected by PI staining and western blot analysis of Bax and PARP proteins. Extracts and all compounds exhibited cytotoxic activity against apoptosis-proficient HL-60 and apoptosis-resistant K562 cells, with the lowest cytotoxic activity on J774 cell line as non-malignant cell.

Armenin as the most potent component decreased the viability of cell with IC50 of 22.5 and 71.1 μ M for K562 and HL-60 cells respectively and selected for further mechanistic study. Armenin increased the sub-G1 peak in flow cytometry histogram of HL-60 and K562 treated cells and increase in the amount of Bax protein and the cleavage of PARP in comparison with the control after treatment for 48 h in K562 treated cells verified the apoptotic activity of the armenin.

Taken together, according to the finding of this study armenin was introduced as a novel cytotoxic compound with apoptotic activity, which is encouraging for further mechanistic and clinical studies.

Keywords: *Artemisia armeniaca*; Asteraceae leukemic cell lines; apoptosis; coumarinhemiterpene ether derivatives.

Introduction

Cancer chemotherapy is mainly owed to the compounds of natural origin. Naturally derived anticancer compounds have been applied directly or as new leads in the discovery of novel structures with anticancer action. Newer generation of cancer chemotherapeutics are selected according to their efficacy and less toxicity profiles. Demanding high potential compounds with the lowest toxicity is the base for screening compounds with the highest potential in clinical implications (1).

^{*} Corresponding author:

E-mail: tayaraninz@mums.ac.ir.

Afterward elucidation of the molecular signaling events elicited by the compounds would give insight to the mechanistic pathways evoked by phytochemicals. Understanding the fact that most anticancer chemicals interfere with programmed cell death propelled most studies to clarify the complex cascade of events that have occurred in apoptosis. In this regards many plants and phytochemicals are proved to be inducers of apoptosis as cancer preventive or chemotherapeutics (2).

Artemisia armeniaca Lam., locally known as Dermane ye Armanestani, is an evergreen or semi-evergreen sub-shrub of the genus Artemisia L. (Asteraceae), which grows abundantly in the Arasbaran area of Eastern Azarbaijan province, northwest of Iran (3). Former phytochemical studies on the aerial parts of A. armeniaca resulted in the identification of simple and prenylated coumarins (4-7) as well as some flavonoids (7). α -Pinene and 1,8-cineole have been reported to be the major constituents of the obtained hydrodistilled oil from the aerial parts of this plant (8) while non-terpene hydrocarbons were identified as the main components of the essential oil in another study (9). Coumarins include a very large class of secondary metabolites observed throughout the plant kingdom (10). Biological effects of coumarin and related derivatives, such as anti-thrombotic and vasodilatory (11), antibacterial (12), anti-mutagenic (13) and cancer chemoprevention as well as anti-tumour (14-18), seem to originate from the coumarin nucleus (16, 19).

The current study aimed to evaluate the *in-vitro* cytotoxic and apoptogenic potential of MeOH and CH_2Cl_2 extracts besides three previously reported coumarins from *A. armeniaca* on two human apoptosis-proficient HL-60 and apoptosis-resistant K562 cancer cells. Extraction and isolation of the compounds were accomplished using the previously described method for purification of coumarinhemiterpene ether derivatives (7).

Experimental

Reagents and chemicals

AlamarBlue[®] (resazurin) from Sigma (Saint Louis, MO, USA); (RPMI-1640) and FCS

from Gibco; Bax (Cat number: 2772), β -actin (Cat number: 4970) and PARP (Cat number: 9542) antibodies, anti-rabbit IgG and HRP linked antibody from CellSignaling technology (Boston, USA); ECL Western blotting detection reagent from Bio-RaD (USA); the fluorescent probe propidium iodide (PI), protease inhibitor cocktail, phosphatase inhibitor cocktail, sodium citrate, Triton X-100, phenylmethylsulfonyl fluoride and Bio-Rad Protein Assay Kit (Hercules, CA); all the solvents used in the extraction and purification procedures were of gradient grade and purchased from Scharlau (Spain) and Caledon (Canada).

Plant materials

The aerial parts of *A. armeniaca* Lam. were collected from Arasbaran, East Azarbaijan province (Iran) during August 2008. The plant was identified by Mr. AH Talebpour from Tabriz University of Medical Sciences. Voucher specimen (No. TBZfph 528) was deposited in the Herbarium of School of Pharmacy, Tabriz University of Medical Science, Tabriz, Iran.

Instrumentation

Vacuum Liquid Chromatography was performed on Sep-Pak Vac 35 cc (10 g) C18 cartridges (Waters). Reversed- phase semipreparative HPLC separations were carried out on a Wellchrom Knauer system (Herbert Knauer GmbH, Berlin, Germany), consisting of a Knauer K-1800 pump, Dr. Maisch GmbH ODS column (250 mm \times 16 mm, 10 μ M) and a Knauer K-2800 UV-Vis detector. NMR spectra were recorded on a Bruker Avance 400 MHz in CD₃OD as the solvent and residual solvent peaks used as internal standard. ESIMS data were obtained on a Finnigan MAT95 spectrometer.

Extraction and purification

The dried and ground aerial parts of *A*. *armeniaca* (100 g) were extracted in a Soxhlet apparatus with CH_2Cl_2 and MeOH (1.5 L each) for 20 and 24 h, successively.

The extracts were separately dried under reduced pressure at a maximum temperature of 45 °C to yield 10.9 and 7.1 g of CH_2Cl_2 and MeOH extracts, respectively.

A portion of the methanol extract $(2 \times 2 \text{ g})$ was



Figure 1. Partitioning scheme of A. armeniaca.

subjected to a Vacuum liquid chromatography (VLC) system with H_2O containing increasing amounts of MeOH (10%, 20%, 40%, 60%, 80% and 100%) to give six fractions (A, B, C, D, E & F) respectively.

Further purification of the fraction C by semipreparative HPLC (mobile phase: 0-60 min, MeOH from 40 to 60% in H₂O; 60-64 min, 60% MeOH in H₂O; 64-68 min MeOH from 60 to 100% in H₂O; 68-75 min 100% MeOH, flow rate 8 mL/min, detection at 220 nm) yielded three prenylated coumarins including armenin (4.4 mg; $t_R = 43.8$ min), isoarmenin (8.0 mg, $t_R = 46.2$ min) and deoxylacarol (3.4 mg, $t_R = 55.8$ min). Structure elucidation of armenin (7-hydroxy-8-(4-hydroxy-3-methylbutoxy)-2H-chromen-2-one), isoarmenin (8-hydroxy-7-(4-hydroxy-3-methylbutoxy)-2H-chromen-2-one) and deoxylacarol (8-(4-Hydroxy-3-methylbutoxy)-7-methoxy-2H-chromen-2-one) was achieved using spectroscopic techniques including ESIMS and proton NMR experiments (Figure 1 and Table 1). Retention time values as well as the obtained spectroscopic data (as follows) were in good agreement with those given in the literature (5, 7).

Armenin (1). White amorphous powder. ¹H-NMR (CD₃OD, 400 MHz): 0.92 (3H, d, *J* = 6.5 Hz, Me-5'), 1.52 (1H, m, H-2'a), 1.82 (1H, m, H-2'b), 1.88 (1H, m, H-3'), 3.36 (1H, m, H-4'a), 3.45 (1H, m, H-4'b), 4.18 (2H, m, H-1'), 6.12 (1H, d, *J* = 9.5 Hz, H-3), 6.78 (1H, d, *J* = 8.8 Hz, H-6), 7.14 (1H, d, *J* = 8.8 Hz, H-5), 7.77 (1H, d, *J* = 9.5 Hz, H-4).

ESIMS: m/z 265.1 [M+H]⁺, 287.2 [M+Na]⁺. Isoarmenin (2). White amorphous powder.

Table 1.	Chemical	structure of	armenin,	isoarmenin	and d	leoxylacarol.

Compound name	R1	R2
Armenin	ОН	4-Hydroxy-3-methylbutoxy
Isoarmenin	4-Hydroxy-3-methylbutoxy	ОН
Deoxylacarol	OCH ₃	4-Hydroxy-3-methylbutoxy

¹H-NMR (CD₃OD, 400 MHz): 1.00 (3H, d, J = 6.5 Hz, Me-5'), 1.56 (1H, m, H-2'a), 1.96 (1H, m, H-2'b), 2.07 (1H, m, H-3'), 3.36 (1H, m, H-4'a), 3.49 (1H, m, H-4'b), 4.19 (2H, m, H-1'), 6.26 (1H, d, J = 9.5 Hz, H-3), 7.02 (1H, d, J = 8.8 Hz, H-6), 7.09 (1H, d, J = 8.8 Hz, H-5), 7.90 (1H, d, J = 9.5 Hz, H-4).

ESIMS: *m/z* 265.1 [M+H]⁺, 287.2 [M+Na]⁺.

Deoxylacarol (3). White amorphous powder. ¹H-NMR (CD₃OD, 400 MHz): 1.08 (3H, d, J = 6.7 Hz, Me-5'), 1.52 (1H, m, H-2'a), 1.82 (1H, m, H-2'b), 2.03 (1H, m, H-3'), 3.52 (1H, m, H-4'a), 3.56 (1H, m, H-4'b), 3.99 (3H, s, methoxy), 4.17 (2H, m, H-1'), 6.31 (1H, d, J = 9.5 Hz, H-3), 7.10 (1H, d, J = 8.7 Hz, H-6), 7.41 (1H, d, J = 8.7 Hz, H-5), 7.93 (1H, d, J = 9.5 Hz, H-4).

ESIMS: *m/z* 279.2 [M+H]⁺, 301.2 [M+Na]⁺.

Cell cultures and treatment agents

The human leukemic cancer cell lines HL-60 and K562 were obtained from Pasteur Institute (Tehran, Iran) and maintained in RPMI-1640 medium with 10% v/v fetal bovine serum and 100 u/mL penicillin and 100 mg/mL streptomycin at 37 °C in a humidified atmosphere of 5% CO_2 and 95% of air.

In-vitro cell viability

AlamarBlue[®] is an indicator dye which incorporates an oxidation-reduction (REDOX) indicator that both fluoresces and changes colour in response to the chemical reduction of growth medium, resulting from cell growth. The alamarBlue® assay is designed to quantitatively measure the proliferation of various human and animal cell lines, bacteria and fungi. Upon entering cells, the blue and non florescent resazurin convert to the florescent and purple resorufin in viable cells (20). About 5×10^4 K562 and 10⁵ HL-60 cells were seeded in each well of 96-microwell plate and treated with various concentrations of each extracts (0-200 µg/mL) and armenin, isoarmenin and deoxylacarol (0-200 µM). J774 cell line was used as nonmalignant cells. After 48 incubation, 20 µL resazurin (0.01% w/v in PBS) was added to each well and the plates were incubated at 37 °C for 4 h before the absorbance was measured at 570 nm (test wavelength) and 600 nm (reference wavelength) in a Synergy H4 Hybrid MultiMode Microplate Reader (BioTek, Winooski, USA). The cytotoxicity of the pure compounds was expressed as IC_{50} , calculated using Prism 5 Software (GraphPad, La Jolla, CA, USA) and presented as mean \pm SD from three independent experiments (with three replicates for each concentration tested extract). For each study, a control sample remained untreated and received only medium in place of the text materials. Paclitaxel 1 μ M was used as positive control.

PI staining

Apoptotic cells were detected by PI staining of small fragments of DNA in treated cells followed by flow cytometry. It has been reported that following DNA fragmentation the so-called sub-G1 peak can be noticed following incubation of cells in a hypotonic phosphate-citrate buffer containing quantitative DNA-binding dye such as PI. Apoptotic cells that have lost DNA will take up less stain and will show up in the left side of the G1 peak in the histogram. Briefly, 106 K562 and HL-60 cells were seeded in each well of a 24-well plate and treated with armenin in different concentrations (0, 25 and 50 μ M) for 48 h. Floating and adherent cells were then harvested and incubated at 4 °C overnight in the dark with 750 μ L of a hypotonic buffer (50 μ g/ mL PI in 0.1% sodium citrate plus 0.1% Triton X-100) before flow cytometric analysis using a FACScan flow cytometer (Becton Dickinson, San Diego, CA) was performed. A minimum of 10⁴ events was acquired for each sample. All data were then analyzed using WinMDI version 2.8 software.

Western blot analysis

About 10^7 HL-60 and K562 cells were treated with 12.5, 25 and 50 µM of armenin, isoarmenin and deoxylacarol (0-200 µM) for 48 h. The cells rinsed and harvested with cool PBS for 3 times, the cell pellet was resuspended in a lysis buffer containing 50 mM Tris-HCl (PH 7.4), 150 mM NaCl, 1% TritonX-100, 1 mM EDTA, 0.2% SDS, 1% Protease inhibitor cocktail, 1% phosphatase inhibitor cocktailand 1 mM phenylmethylsulfonyl fluorideand left on ice for 30 min. After centrifugation at 10000 rpm for 20 min at 4 °C, the cell lysate was collected and protein concentration was determined according



Figure 2. The concentration-dependent effects of CH2Cl2 and MeOH extracts (0-200 μ g/mL) and 7-hydroxy-8-(4-hydroxy-3-methylbutoxy) comarin (armenin), 8-hydroxy-7-(4-hydroxy-3-methylbutoxy) comarin (isoarmenin) and deoxylacarol on the viability of K562 and HL-60 cells. Among three compounds armenin exhibited high cytotoxic activity against apoptosis-proficient HL-60 and apoptosis-resistant K562 cells, with IC50 values ranging from 22.5 to 71.1 μ M with much less cytotoxic effects on normal human lymphocytes. Paclitaxel 1 μ M was used as positive control. Values were mean±SD of at least three independent experiments, each in triplicates. +P < 0.05, ++P < 0.01 and +++P < 0.001 compared to control.

to the Bio-Rad Protein Assay kit. Equal amount of proteins were subjected to 12% SDS-page (W/V). The proteins were transferred to a polyvinylidene fluoride (PVDF) membrane and subjected to immunoblotting using Bax, β -actin and, PARP antibody as primary antibodies and anti-rabbitIgG and HRP-linked antibody as secondary antibodies, Bax protein band and PARP cleavage in K562 and HL-60 cells were detected by enhanced chemiluminescenceusing the ECL western blotting detectin reagent. Images were quantified using Gel-pro Analyser V.6.0 Gel Analysis software (Media Cybernetics, InC, Bethesda, MD).

Statistical analysis

One way analysis of variance (ANOVA) and Bonferroni posthoc test were used for

data analysis. All the results were expressed as mean \pm SD and *p* values below 0.05 were considered statistically significant.

Results

Cytotoxicity of isolated compounds

Armenin, isoarmenin and deoxylacarol were examined for cytotoxic potential on K562, HL-60 and normal cells (J774). Cells were incubated at 37 °C and 5% CO₂ with various concentrations of the CH₂Cl₂ and MeOH extracts (0-200 μ g/mL) and compounds (0-200 μ M) for 48 h. Results demonstrated that extracts decreased cell viability in a concentration-dependant manner (Figure 2). Among all the samples, armenin demonstrated the most cytotoxic effects on cancer cells, with limited adverse effect on

Table 2. IC_{s_0} values for CH_2Cl_2 and MeOH extracts (μ g/mL) and 7-hydroxy-8-(4-hydroxy-3-methylbutoxy) comarin (armenin), 8-hydroxy-7-(4-hydroxy-3-methylbutoxy) comarin (isoarmenin) and deoxylacarol (μ M) in HL-60 and K562 cell lines.

Cell Line	Extracts		Pure compounds			
	CH2Cl2	МеОН	Armenin	Isoarmenin	Deoxylacarol	
K562	130.7	155.4	22.5	>200	108.9	
HL-60	74.95	>200	71.1	>200	>200	



Figure 3. Percentage of sub-G1 peak after PI staining and flow cytometry analysis of armenin (0.0, 25 and 50 μ M) showing apoptosis induction in K562 and HL-60 cells. All experiments were done in triplicate. *P < 0.05 compared to control.

normal cells. IC_{50} values (μ M) for different extracts of armenin in HL-60 and K562 cells are presented in Table 2. Paclitaxel 1 μ M was used as positive control and decreased the viability of K562, HL-60 to 25.5 ± 0.5 and 96.3 ± 1.5 respectively at this concentration.

Apoptosis induction by isolated compounds

Apoptosis in K562 and HL-60 cell lines was detected with flow cytometry using PI staining test. Cells incubated with various concentrations (0, 25 and 50 μ M) of armenin for 48 h. Sub-G1 peak of treated cells in flow cytometry histograms compared to that (=Sub-G1 peak) of untreated control cells revealed the induction of apoptosis in treated cells (Figure 3).

Effect of armenin on poly (ADP ribose) polymerase cleavage

Since K562 cells were more sensitive to armenin K562 cells were selected for western blot analysis. The increase in the amount of Bax and cleavage of PARP-1 116 kDa to 89 and 24 kDa fragments was used as an indicator of apoptosis. In K562 cells, the amount of Bax was increased and PARP-1 was cleaved clearly to the 89 kDa and 24 kDa fragments after treatment with armenin (0.0, 12.5 and 50 μ M) after 48 h (Figure 4).

Discussion

We have previously demonstrated the potent cytotoxic and apoptotic activity of the CH₂Cl₂ extract of A. armeniaca in K562 and HL-60 cells (21). In this regard, we have chosen the plant for further analytical and biological assays. In this study, we have examined the cytotoxic properties of CH₂Cl₂ and MeOH extracts (0-200 µg/mL) and armenin, isoarmenin and deoxylacarol isolated from A. armeniaca. The results show that the reduced viability of cancer cells treated with armenin is accompanied by apoptosis induction verified by sub G1 peak in the flowcytometery histogram and PARP cleavage of treated cells. We therefore infer that among three pure compounds, armenin is a potent cytotoxic and apotogenic one that could triggers at least certain aspects of the apoptosis signaling pathway, which activates apoptotic mediators that are closely associated with the levels of DNA damage and degradation of active DNA repairing enzymes, PARP (22). PARP cleavage can be seen by not only in K562 cells with IC_{50} values of armenin, but also in lower concentrations of the compound.



Figure 4. Proteolytic cleavage of poly (ADP-ribose) polymerase (PARP) in K562 cells after 48 h exposure to armenin (0.0, 12.5 and 50 μ M). β -Actin was used as a loading control. All Western blots were representative of 3 independent experiments. All experiments were done in triplicate.

Although K562 cells are apoptosis resistant (23), our findings suggest this cell line is more sensitive to armenin than HL-60 cells. This controversy may result from different pathways affected by armenin in K562 and HL-60 cells.

Armenin's cytotoxic mechanism on cancer cells is still not fully undrestood. The difference in the cytotoxic activity of armenin, isoarmenin and deoxylacarol may be linked to other factors such as side chains, chiral structure, lipophilicity, and molecular or electrical features, affecting the affinity of the molecule to the active site of action.

Cytotoxic and apoptogenic properties of different extracts of various species from the genus *Artemisia* have been studied so far (24-27) and have resulted in the isolation of some known coumarins like scopoletin and isoscopoletin from the active extract (25). A previous study (28) has revealed the cytotoxic activity of isoscopoletin against lung and colon cancer cell lines. Cytotoxic and cytostatic effects of scopoletin on tumoural lymphocytes had already been reported as well (29).

There are reports that prenylated coumarins can exhibit cytotoxic activity. Ferulenol, a prenylated 4-hydroxycoumarin from *Ferula communis* with remarkable dose-dependent cytotoxicity against various human tumor cell lines, displayed stimulation of tubulin polymerization *in-vitro* and rearrangement of cellular microtubule network as well as alteration in nuclear morphology (30). Based on another study, ferulenol possessed cytotoxic properties on hepatocytes. Additionally, the prenylated coumarin reduced factor X biosynthesis at noncytotoxic concentrations.

The prenyl residue played a major role in ferulenol activity (31). A study on the antileukemic activities of a series of prenylated coumarins isolated from *Toddalia asiatica* (L.) Lam. indicated that the most potent cytotoxic and anti-proliferative compound displayed a dual effect as a cell differentiating agent and apoptosis inducer in U-937 cells (32). Minutin B, a monoterpene coumarin from *Micromelum minutum* Wight & Arn. leaves exhibited strong cytotoxic activity against four lung adenocarcinoma cell lines including A549, SBC3, K562, and K562/ADM (33).

Conclusion

We have previously demonstrated that CH_2Cl_2 extract of *A. armeniaca* is a highly potent cytotoxic agent against K562 and HL-60 cells with minimal effect on normal cells. Herein we have done additional cytotoxic and apoptogenic assays on three pure compounds isolated from *A. armeniaca* on human leukemia cell lines and human normal cells.

Based on this study the cytotoxic and apoptogenic activity of armenin was confirmed. It is also worthy to emphasize that the high antileukemic activity of armenin was accomplished with low toxicity to human normal cells. Thus, it is imperative to continue the pursuit of new therapeutic aspects of armenin as a novel anticancer agent and it may be beneficial to discover multiple signaling pathways affected by the compound for clinical usage.

Conflict of Interest

The authors declare that there are no conflicts of interest.

Acknowledgments

The authors would like to thank Mr. M. Malaekeh for his assistance in flow cytometry. This work was supported by grants (No. 89681) from Research Affairs of Mashhad University of Medical Sciences as a part of Pharm. D. thesis.

References

- Narang AS and Desai DS. Anticancer Drug Development Unique Aspects of Pharmaceutical Development. In: Lu Y and Mahato RI (eds.) *Pharmaceutical Perspectives of Cancer Therapeutics*. Springer Science+Business Media (2009) 49-93.
- (2) Kong AN, Yu R, Chen C, Mandlekar S and Primiano T. Signal transduction events elicited by natural products: role of MAPK and caspase pathways in homeostatic response and induction of apoptosis. *Arch. Pharm. Res.* (2000) 23: 1-16.
- (3) Mozaffarian V. A Dictionary of Iranian Plant Names. Farhang Moaser Publishers, Tehran (1998) 56-8.
- (4) Rybalko KS, Konovalova OA, Sheichenko VI and Zakharov PI. Armenin a new coumarin from *Artemisia armeniaca. Khim. Prir. Soed.* (1976) 3: 294-8.
- (5) Szabo G, Greger H and Hofer O. Coumarin-hemiterpene ethers from *Artemisia* species. *Phytochemistry* (1985) 24: 537-41.

- (6) Mojarrab M, Delazar A, Hamburger M and Potterat O. New coumarin-hemiterpene ether glucosides and a structurally related phenylpropanoic acid derivative from *Artemisia armeniaca*. *Nat. Prod. Commun.* (2010) 4: 1619-22.
- (7) Mojarrab M, Delazar A, Moghadam SB, Nazemiyeh H, Nahar L, Kumarasamy Y, Asnaashari S, Hadjiakhoondi A and Sarker SD. Armenin and isoarmenin - two prenylated coumarins from the aerial parts of *Artemisia armeniaca. Chem. Biodivers.* (2011) 8: 2097-103.
- (8) Kazemi M, Zand MR, Roshanaei K, Mehrzad M and Rustaiyan A. Composition of the volatile oils of *Artemisia armenica* Lam. and *Artemisia splendens* Willd. From Iran. J. Essent. Oil Res. (2010) 22: 126-8.
- (9) Mojarrab M, Delazar A, Esnaashari S and Heshmati Afshar F. Chemical composition and general toxicity of essential oils extracted from the aerial parts of *Artemisia armefniaca* Lam. and *A. incana* (L.) Druce growing in Iran. *Iran. Res. Pharm. Sci.* (2013) 8: 65-9.
- (10) Egan D, O'Kennedy R, Moran E, Cox D, Prosser E and Thornes RD. The pharmacology metabolism analysis and applications of coumarin and coumarin related compounds. *Drug Metab. Rev.* (1990) 22: 503-29.
- (11) Hoult JRS and Paya M. Pharmacological and biochemical actions of simple coumarins: natural products with therapeutic potential. *Gen. Pharmacol.* (1996) 27: 713-22.
- (12) Laurin P, Klich M, Dupis-Hamelin C, Mauvais P, Lassaigne P, Bonnefoy A and Musicki B. Synthesis and *in-vitro* evaluation of novel highly potent coumarin inhibitors of gyrase B. *Bioorg. Med. Chem. Lett.* (1999) 9: 2079-84.
- (13) Pillai SP, Menon SR, Mitscher LA, Pillai CA and Shankel DA. Umbelliferone analogues and their potential to inhibit benzo(a)pyrene and hydrogen peroxide induced mutations. J. Nat. Prod. (1999) 62: 1358-62.
- (14) Chen Y, Liu HR, Liu HS, Cheng M, Xia P, Qian K, Wu PC, Lai CY, Xia Y, Yang ZY, Morris-Natschke SL and Lee KH. Antitumor agents 292. Design synthesis and pharmacological study of S- and O-substituted 7-mercapto- or hydroxy-coumarins and chromones as potent cytotoxic agents. *Eur. J. Med. Chem.* (2012) 49: 74-85.
- (15) Egan D, James P, Cooke D and O'Kennedy R. Studies on the cytostatic and cytotoxic effects and mode of action of 8-nitro- 7-hydroxycoumarin. *Cancer Lett.* (1997) 118: 201-11.
- (16) Finn GJ, Creaven B and Egan D. Study of the *in-vitro* potential of natural and synthetic coumarin derivatives using human normal and neoplastic skin cell lines. *Melanoma Res.* (2001) 11: 461-47.
- (17) Kostova I. Synthetic and natural coumarins as cytotoxic agents. *Curr. Med. Chem. Anticancer Agents* (2005) 4: 29-46.
- (18) Maucher A, Kager M and von Angerer E. Evaluation of the antitumor activity of coumarin in prostate cancer models. J. Clin. Res. Clin. Oncol. (1993) 119: 150-4.

- (19) Jimenez-Orozco FA, Molina-guarneros JA, Mendozapatino N, Leon-cedeno F, Flores-perez B, Santos-santos E and Mandoki JJ. Cytostatic activity of coumarin metabolites and derivatives in the B-16-F10 murine melanoma cell line. *Melanoma Res.* (1999) 9: 243-7.
- (20) O'Brien J, Wilso I, Orton T and Pognan F. Investigation of the alamar blue (resazurin) fluorescent dye for the assessment of mammalian cell cytotoxicity. *Eur. J. Biochem.* (2000) 267: 5421-6.
- (21) Mojarrab M, Lagzian MS, Emami SA, Asili J and Tayarani-Najaran Z. *In-vitro* anti-proliferative and apoptotic activity of different fractions of *Artemisia armeniaca* Lam. *Rev. Bras. Farmacogn.* (2013) 23: 783-8.
- (22) Bartek J and Lukas J. DNA damage checkpoints: from initiation to recovery or adaptation. *Curr. Opin. Cell Biol.* (2007) 19: 238–45.
- (23) Munker R and Andreeff M. Induction of death (CD95/ FAS) activation and adhesion (CD54) molecules on blast cells of acute myelogenous leukemias by TNFalpha and IFN-gamma. *Cytokines Cell Mol. Ther.* (1996) 2: 147-59.
- (24) Akrout A, Gonzalez LA, El Jani H and Madrid PC. Antioxidant and antitumor activities of *Artemisia campestris* and *Thymelaea hirsuta* from southern Tunisia. *Food Chem. Toxicol.* (2011) 49: 342-7.
- (25) Piao GC, Li YX, Yuan HD and Jin GZ. Cytotoxic fraction from *Artemisia sacrorum* Ledeb. against three human cancer cell lines and separation and identification of its compounds. *Nat. Prod. Res.* (2012) 26: 1483-91.
- (26) Tayarani-Najaran Z, Sareban M, Gholami A, Emami SA and Mojarrab M. Cytotoxic and apoptotic effects of different extracts of *Artemisia armeniaca* Krasch. on K562 and HL-60 cell lines. *Scientific World Journal* (2013) ID 628073.
- (27) Tayarani-Najaran Z, Hajian Z, Mojarrab M and Emami SA. Cytotoxic and Apoptotic Effects of Extracts of *Artemisia ciniformis* Krasch. & Popov ex Poljakov on K562 and HL-60 Cell Lines. *Asian Pac. J. Cancer Prev.* (2014) 15: 7055-9.
- (28) McGovern PE, Christofidou-Solomidou M, Wang W, Dukes F, Davidson T and El-Deiry WS. Anticancer activity of botanical compounds in ancient fermented beverages. *Int. J. Oncol.* (2010) 37: 5-14.
- (29) Manuele MG, Ferraro G, Barreiro Arcos ML, López P, Cremaschi G and Anesini C. Comparative immunomodulatory effect of scopoletin on tumoral and normal lymphocytes. *Life Sci.* (2006) 79: 2043-8.
- (30) Bocca C, Gabriel L, Bozzo F and Miglietta A. Microtubule-interacting activity and cytotoxicity of the prenylated coumarin ferulenol. *Planta Med.* (2002) 68: 1135-7.
- (31) Monti M, Pinotti M, Appendino G, Dallocchio F, Bellini T, Antognoni F, Poli F and Bernardi F. Characterization of anti-coagulant properties of prenylated coumarin ferulenol. *BBA-Gen. Subjects* (2007) 1770: 1437-40.
- (32) Vázquez R, Riveiro ME, Vermeulen M, Mondillo

C, Coombes PH, Crouch NR, Ismail F, Mulholland DA, Baldi A, Shayo C and Davio C. Toddaculin a natural coumarin from *Toddalia asiatica* induces differentiation and apoptosis in U-937 leukemic cells. *Phytomedicine* (2012) 19: 737-46.

(33) Sakunpak A, Matsunami K, Otsuka H and

Panichayupakaranant P. Isolation of new monoterpene coumarins from *Micromelum minutum* leaves and their cytotoxic activity against *Leishmania major* and cancer cells. *Food Chem*. (2013) 139: 458-63.

This article is available online at http://www.ijpr.ir