

Synthesis and Evaluation of the Antimicrobial Activity of Spiro-4H-pyran Derivatives on Some Gram Positive and Gram Negative Bacteria

Pardis Nazari^a, Aliyeh Bazi^a, Seyed AbdulMajid Ayatollahi^b, Hadi Dolati^b, Seyedeh Mahbobeh Mahdavi^b, Laleh Rafighdoost^a and Marzieh Amirmostofian^{c*}

^aDepartment of Pharmaceutical Biotechnology, School of Pharmacy, Zabol University of Medical Sciences, Zabol, Iran. ^bPhytochemistry Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran. ^cDepartment of Medicinal Chemistry, School of Pharmacy, Zabol University of Medical Sciences, Zabol, Iran.

Abstract

Infections are one of the most important causes of death, disability and inappropriate conditions for millions of people around the world. Therefore, the development in prognosis, prevention and treatment of infectious diseases made a considerable progress in designing and synthesis of new antimicrobial drugs. Nowadays, due to the increase in microbial resistance, discovery of new compounds with broad spectrum effects is granted. 4H-pyran derivatives and spiro compounds are the most important fragments in some effective drugs with antimicrobial activity. Therefore, in this study, 6 compounds with spiro-4H-pyran core were synthesized and evaluated for their antimicrobial activity against four different bacterial species using microbroth dilution and disk diffusion methods. Minimum inhibitory concentration (MIC) has been measured for each compound and also for the standard antibiotic, gentamicin, and they were all compared together. According to our result, one of the spiro-pyran derivative (5d) containing both the indole and the cytosine ring, has been shown good antibacterial effects against standard and clinical isolates of *Staphylococcus aureus* and *Streptococcus pyogenes*.

Keywords: Spiro-4H-pyran; one-pot reaction; three-component reaction; *Staphylococcus aureus*; *Streptococcus pyogenes*; Microbroth dilution; Disk diffusion.


Introduction

Despite a century of often successful struggles for prevention and treatment of infectious diseases, the issue remains an important worldwide problem in public health which brings about the loss of more than 13 million individuals each year (1). In the last decades, antibiotic resistance, especially in the developing countries, elevated with an accelerating steep. The noticeable factors which can cause the

rise of antibiotic-resistant strains are the free access of patients to antibiotics and the use of subtherapeutic doses. In addition, the lack of an effective, forethought, and judicious antibiotic policy, and also the insufficient intervention of the medical professions to supervisory programs, if any exist, spread the antibiotic-resistance (2). Increasing transportation facilities and thus international travels, provides great opportunities for antibiotic resistant bacteria dissemination to other countries (3,4). In recent years, pharmaceutical companies spent less budget in the field of antimicrobial discovery research and development and as a result the

* Corresponding author:

E-mail: *mamirmostofian@gmail.com



Scheme 1. One-pot three-component synthesis of spiroopyran compounds(1-6)d.

number of new antibiotics reached approval for human use was too low. In addition, there has been less innovation in the structural and pharmaceutical characteristics of designed and synthesized antimicrobials, so the outcome is a shortage of newer, safer and more effective antimicrobial agents. The consequences of these events are the emergence and spread of antibiotic resistant bacteria which followed by the increasing incidence of potentially serious MDR bacterial infections (5-7).

In fact, amino-4H-pyrans and constitutes are biologically active compounds with a wide spectrum of activities such as antimicrobial, antifungal, antitumor (8), anticoagulant (9), diuretic (10), spasmolytic, and antianaphylactic activities (11). Furthermore, amino-4H-pyrans represent the building blocks of a series of considered bioactive products (5). On the other hand, spiroheterocycles have recently attracted much attention as an important class of heterocyclics which have useful biological and pharmacological properties, such as anticonvulsants (12), antimicrobials and antibacterials (13, 14), anti-tubercular agents (15), anticancer agents (16), antidermatitis agents (17), pesticides (18), antioxidants (19), and enzyme inhibitors (20).

Increase in microbial resistance worldwide against current antimicrobial agents results in the necessity of comprehensive evaluation of prepared new innovative agents. In this regard, 6 chemical compounds with spiroaminopyran nucleus (Table 1) were synthesized. The synthesis was carried out by one-pot, three-component reaction of satin or ninhydrin with malononitrile and beta-dicarbonyl substances in the presence of 4-dimethylaminopyridine (DMAP) as catalyst

(Scheme 1). The compounds were examined by using microbroth dilution and disk diffusion methods on a selected group of gram positive and gram negative bacteria.

Experimental

Chemicals


All chemicals, reagents and solvents used in this study were purchased from Merck AG, Fluka and Aldrich Chemical Company and were used without further purifications. Infrared spectra were acquired using a Perkin Elmer Model 1420 spectrometer. A Bruker AMX-300 MHz instrument (Bruker Biosciences, Germany) was used to acquire ¹H NMR spectra with TMS as internal standard. Chloroform-D and DMSO-D₆ were used as solvents. Coupling constants (J) values are calculated in hertz (Hz) and spin multiplicities are given as s (singlet), d (double), t (triplet), and m (multi). The mass spectra are available in references (21-24).

Chemistry

General procedure for the preparation of compounds (1-6)d

In a general procedure, a mixture of ninhydrin (1 mmol) or isatin (1 mmol) and malononitrile (1 mmol) in 2 mL of DMF were stirred in a 5-mL round-bottomed flask in the presence of 5 mol% of 4-dimethylaminopyridine (DMAP) at room temperature. Then beta-dicarbonyl compounds **c** (1 mmol) was added to the mixture and the contents were stirred at room temperature for appropriate time (Scheme 1, Table 1). After the completion of the reaction as followed by TLC, water (5 mL) was added to the mixture and the solid product was collected

Table 1. The structures and yields of compounds (1-6)d.



Entry	a,a'	c	Compounds	Time (min)	Yield (%)	Mp (oC)		Ref
						Found	Ref	
1			(1d) 	6	88	291-293	290-292	21
2			(2d) 	20	80	274-277	275-277	21
3			(3d) 	35	75	225-227	220-225	21
4			(4d) 	25	90	265-268	260-262	22
5			(5d) 	10	85	>300	298-299	23
6			(6d) 	8	65	286-288	284-286	24

by filtration and washed with almost 20 mL of water and the purified compounds (1-6) d were obtained (Table 1).

2-amino-7,7-dimethyl-1',3',5-trioxo-1',3',5,6,7,8-hexahydrospiro[chromene-4,2'-indene]-3-carbonitrile (1d): Yellow powder,

IR (KBr) (λ_{\max} , cm^{-1}): 3376, 3311, 3143, 2192, 1722, 1682, 1655. ^1H NMR (DMSO- d_6 , 300 MHz): 8.04–7.97 (m, 4H, Ar-H), 7.66 (s, 2H, NH_2), 2.60 (s, 2H, CH_2), 2.18 (s, 2H, CH_2), 1.02 (s, 6H, 2CH_3). ^{13}C NMR (DMSO- d_6): 199.7, 196.0, 166.5, 159.8, 140.5, 136.6, 123.1, 116.8, 110.0, 53.0, 51.7, 48.9, 32.4, 27.1.

7'-amino-1,2',3,4'-tetraoxo-1,1',2',3,3',4'-hexahydrospiro[indene-2,5'-pyrano[2,3-d]pyrimidine]-6'-carbonitrile (2d): Yellow powder, IR (KBr) (λ_{\max} , cm^{-1}): 3361, 3312, 3198, 2196, 1745, 1715, 1673, 1591, 1341, 1263. ^1H NMR (DMSO- d_6 , 300 MHz): 11.13 (s, 1H, NH), 9.33 (s, 1H, NH), 8.06–8.00 (m, 4H, Ar-H), 7.71 (s, 2H, NH_2). ^{13}C NMR (DMSO- d_6): 200.3, 194.1, 163.3, 162.6, 160.0, 156.3, 151.5, 150.9, 141.5, 140.4, 136.7, 135.1, 123.1, 122.9, 116.7, 113.0, 84.6, 57.8, 53.1, 44.0, 22.7.

Ethyl-2'-amino-3'-cyano-6'-methyl-1,3-dioxo-1,3-dihydrospiro[indene-2,4'-pyran]-5'-carboxylate(3d): Yellow powder, IR (KBr) (λ_{\max} , cm^{-1}): 3334, 3176, 2926, 2192, 1659, 1429, 1275, 1166, 1089, 1013. ^1H NMR (DMSO- d_6 , 300 MHz): 7.99–7.84 (m, 3H, Ar-H), 7.81–7.76 (m, H, Ar-H), 6.90 (s, 2H, NH_2), 4.31–4.15 (m, 2H), 2.25 (s, 3H), 1.35–1.42 (m, 3H). ^{13}C NMR (DMSO- d_6): 193.9, 168.7, 167.5, 164.2, 143.9, 137.4, 137.3, 133.2, 125.7, 125.2, 116.7, 107.3, 60.6, 56.7, 14.5, 14.0.

Ethyl-2'-amino-3'-cyano-6'-methyl-2-oxospiro[indoline-3,4'-pyran]-5'-carboxylate(4d): White powder, IR (KBr) (λ_{\max} , cm^{-1}): 3482, 3283, 3158, 3078, 2978, 2191, 1723, 1701, 1676, 1618. ^1H NMR (DMSO- d_6 , 300 MHz): 10.40 (s, 1H, NH), 7.18 (t, 1H, $J=7.6$ Hz, Ar-H), 7.16 (s, 2H, NH_2), 7.05 (d, 1H, $J=7.1$ Hz, Ar-H), 6.94 (t, 1H, $J=7.3$ Hz, Ar-H), 6.79 (d, 1H, $J=7.6$ Hz, Ar-H), 3.70–3.79 (m, 2H, CH_2), 2.29 (s, 3H, CH_3), 0.74 (t, 3H, $J=7.0$ Hz, CH_3). ^{13}C NMR (DMSO- d_6): 178.6, 164.5, 158.9, 158.5, 142.1, 134.5, 128.5, 123.4, 121.8, 117.5, 109.3, 104.6, 60.2, 56.4, 48.9, 18.5, 13.0.

7'-amino-2,2',4'-trioxo-1',2',3',4'-tetrahydrospiro[indoline-3,5'-pyrano[2,3-d]pyrimidine]-6'-carbonitrile (5d): White powder, IR (KBr) (λ_{\max} , cm^{-1}): 3559, 3286, 3215, 2199,

1717, 1642, 1324. ^1H NMR (DMSO- d_6 , 300 MHz): 12.09 (s, 1H, NH), 11.01 (s, 1H, NH), 10.45 (s, 1H, NH), 7.33 (s, 2H, NH_2), 7.09–7.28 (m, 4H, Ar-H). ^{13}C NMR (DMSO- d_6): 177.7, 161.5, 158.3, 153.8, 149.6, 142.5, 133.6, 128.3, 123.7, 121.7, 117.0, 109.2, 86.5, 57.7, 46.6.

2'-amino-2,5'-dioxo-5'H-spiro[indoline-3,4'-pyranof[3,2-c]chromene]-3'-carbonitrile (6d): White powder, IR (KBr) (λ_{\max} , cm^{-1}): 3324, 3178, 2201, 1721, 1673, 1610, 1471, 1358. ^1H NMR (DMSO- d_6 , 300 MHz): 10.67 (s, 1H, NH), 7.92 (d, 1H, $J=7.9$ Hz, Ar-H), 7.73 (t, 1H, $J=8.1$ Hz, Ar-H), 7.66 (s, 2H, NH_2), 7.50 (t, 1H, $J=7.6$ Hz, Ar-H), 7.46 (d, 1H, $J=8.4$ Hz, Ar-H), 7.19 (t, 2H, $J=7.5$ Hz, Ar-H), 6.90 (t, 1H, $J=7.4$ Hz, Ar-H), 6.83 (d, 1H, $J=8.0$ Hz, Ar-H). ^{13}C NMR (DMSO- d_6): 177.2, 158.4, 155.0, 152.0, 142.0, 133.6, 133.0, 128.9, 125.0, 124.1, 122.6, 122.0, 116.9, 116.6, 112.4, 109.5, 101.4, 57.0, 47.5.

Microbial strains

From June 2014 to January 2015, five clinical isolates of each microbial strain, including *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, and *Streptococcus pyogenes* were recovered from human infections. The clinical specimens isolation was carried out based on standard bacteriological procedures. Then the isolates were identified by use of gram staining technique, differential cultures, and routine biochemical tests. Reference strains which were used in this study consist of *S. aureus* (ATCC25923), *E. coli* (ATCC8739), *P. aeruginosa* (PTCC1599), *S. pyogenes* (PTCC1447).

Antimicrobial susceptibility testing

A stock solution (1024 $\mu\text{g}/\text{mL}$) of each compound (1-6)d was prepared in dimethyl sulfoxide 20% to be used in antimicrobial susceptibility testing. Evaluation of spiroaminopyran compounds antibacterial activities was performed by broth microdilution and disk diffusion methods. For this purpose, 100 μL of the Mueller Hinton broth medium was added to each of the 96 wells in sterile microtiter plates. A series of doubling dilutions for each compound were made in microtiter plate wells, in the concentration range of 1 to 512 $\mu\text{g}/\text{mL}$ for

each of six different compounds. Fresh culture of each clinical and standard strain was harvested and microbial colonies were suspended in 0.9% NaCl solution. The microbial concentration was adjusted to 0.5 McFarland standard by turbidimetric method. Each well was inoculated with the prepared suspension in the way that the final 5×10^5 CFU/ml bacterial concentration in the well was obtained. The plates were incubated at 37 °C for 24 h. All the mentioned procedures were done also for gentamicin as the standard antimicrobial agent by the difference that a stock solution of 1,024 mg/liter gentamicin powder in sterile water was prepared. The lowest compound concentration that visibly inhibits microbial growth was reported as the minimum inhibitory concentration (MIC). It should be noted that on each plate, positive control containing broth medium with bacterial inoculum and negative control which was prepared by adding DMSO (20%) to broth medium were noticed and all the procedure was according to the standard protocol of NCCLS broth microdilution MIC determining method.

For the semi quantitative disk diffusion procedure, sterile 6 mm paper disks which were impregnated by spiroaminopyran compounds were used. Mueller Hinton agar plates were inoculated by a swab from 1.5×10^8 CFU/ml bacterial suspension and the antibiotic contained disks were placed on the plate surface. The amount of each compound was loaded on paper disks chosen based on microbroth dilution results in the range of 16, 32, 64, and 128 µg. The plates were incubated for 24 h at 37 °C. By the way, a 10 µg gentamicin standard disk also was included. Following incubation, the inhibition zone diameter around each disk was measured in millimeters. All the procedure was done according to the standard protocol of NCCLS disk diffusion susceptibility method (25). All the above tests for each compound were done in duplicate for three times.

Results and Discussion

In this article, we have described an efficient one-pot, three-component reaction for the synthesis of spiroamino-pyran compounds. Several methods have been reported for the

synthesis of spiropyran-fused heterocycles. Although each of the known methods for the synthesis of spiropyran systems has its advantages, still there is further scope to develop a new methodology using a less expensive catalyst under mild reaction conditions and applicable to a wide range of substrates in great demand. As mentioned, in this paper, 4-dimethylaminopyridine (DMAP) (26) was used as an efficient important and useful catalyst in this one-pot three-component reaction for the synthesis of spiroamino-pyran(1-6)d compounds.

At first, we tested the reaction of ninhydrin (a) malononitrile (b) and dimedon (c) as a simple model substrate to find the optimal conditions and amount of catalyst required for the synthesis of spiropyran derivatives in different solvent and molar ratio of DMAP as catalyst. The results are summarized in Table 2.

We started our investigation by reacting ninhydrin (a) with malononitrile (b) and dimedon (c) in the presence of (DMAP) 10 mol% in various solvents (Scheme 1-Table 2). It was found that when the reaction was carried out with Et₂O as solvent, only trace product was detected (Table 2, entry4). As shown in Table 2, THF, H₂O, DMSO, CHCl₃ and CH₂Cl₂, gave low yield and long reaction time of product (Table 2, entries 5-9). The yield of the reaction in DMF is greater and the experiments obviously showed an increased reaction rate in DMF as compared to other solvents such as C₂H₅OH, CH₃OH, n-hexan (Table 2, entries 1,2,10). The optimized amount of the catalyst was determined using different mole percent of DMAP such as (2 mol %) (5 mol %). (15 mol %) and (20 mol%) (Table 2). Accordingly, the best yield in the shortest reaction time was obtained using the 5 mol% of DMAP (Table 2). The subsequent increase in the catalyst load did not improve the product yield of the reaction and 2 mol% of DMAP catalyzed the reaction but needs a longer reaction time.

Spiropyran (Ld) of ninhydrin, malononitrile and dimedonin in the presence of DMAP (5 mol%) and using DMF as a solvent at ambient temperature was obtained in 88 % yield. The scope and efficiency of this approach was explored for the synthesis of a variety

Table 2. Evaluation of molar ratio of DMPA and solvents in the synthesis of compound 1d.

Entry	Solvent	Catalist DMPA (mol %)	Time (min)	Yield (%)
1	C ₂ H ₅ OH	(10 mol %)	15	69
2	CH ₃ OH	(10 mol %)	12	65
3	DMF	(10 mol %)	5	75
4	Et ₂ O	(10 mol %)	160	trace
5	THF	(10 mol %)	30	42
6	H ₂ O	(10 mol %)	10	62
7	DMSO	(10 mol %)	20	60
8	CHCl ₃	(10 mol %)	30	65
9	CH ₂ Cl ₂	(10 mol %)	10	57
10	n-hexan	(10 mol %)	120	72
11	DMF	(2mol %)	10	76
12	DMF	(5mol %)	6	88
13	DMF	(15mol %)	5	63
14	DMF	(20mol %)	3	58

of spiroopyrans by isatin or ninhydrin with malononitrile and beta-dicarbonyl compounds and results are summarized in Table 1. Fortunately, it was found that nearly all of the corresponding products were obtained in good yields.

A detailed proposed mechanism of the reaction is shown in Figure 1. As shown in Figure 1, using the DAMP as a catalyst, the Knoevenagel reaction is initiated by formation of malononitrile anion (II). In this mechanism, the anion (II) is attacked to ninhydrin (a) or isatin (a') to yield the compound III which undergoes the Knoevenagel condensation to afford the compound V by elimination a molecule of water. The beta-dicarbonyl compounds (c) activated by DAMP (VI), is attacked to the compound V. Also, the new intermediate compound VII undergoes the other spontaneously reactions to obtain the final products the spiroaminopyran compounds (1-6) d.

The antimicrobial activity of the spiroamino-4H-pyran compounds (1-6) d was evaluated against a standard and 5 isolated clinical bacterial strains of *Staphylococcus aureus*, *Streptococcus pyogenes*, *Escherichia coli*, and *Pseudomonas*

aeruginosa, which were identified by use of gram staining (Figure 2) and biochemical tests, by the disk diffusion and broth microdilution methods (25). Preliminary screening and MIC determination of the spiroaminopyran derivatives was performed in different concentrations encompassed the range of 1, 2, 4, 8, 256 and 512 µg/mL.

The MIC (µg/mL) of mentioned compounds (1-6) d against tested strains are presented in Table 3. Amongst all of these compounds, the cytosine derivative of spiroaminopyran, compound 5d, was the most effective one with the most activity against the clinical isolates of both *Staphylococcus aureus* and *Streptococcus pyogenes*. Compound 5d, especially, exhibited good activity against clinical isolates of *S. aureus* and *S. pyogenes* with MICs of 32 and 64 µg/mL, respectively (Table 3). The compound showed moderate antibacterial activity against the standard *S. aureus* and *S. pyogenes*. It was inactive against standard and clinical isolates of gram negative bacteria (MIC ≥ 512) (Table 3).

Growth inhibition was recorded by measuring the inhibition zone diameter at the end of 18-24 h. incubation and the results were shown in




Figure 1. A detailed mechanism of preparation of spiroaminopyrans with DMAP.

Table 4. Gentamicin was used as standard drug in this assay.

Since other derivatives of spiroaminopyran than no 5 d did not show any significant effects on investigated bacteria, the results of inhibition zone diameter measurements were not included.

By comparing the inhibiting effects of the spiroaminopyran derivatives (1-6)d we can deduce that the presence of indole ring and cytosine-like moiety conferred the highest antibacterial activity in these substances. In some researches, the compounds containing only cyanide and amino substituents on aminopyran derivatives have not shown appropriate effects against bacteria (27-29) which is also shown in our investigation. Compounds with just spiro-pyran and indole-pyran core showed a potent antibacterial activity (28, 30) and some others, containing spiroaminopyran-indole moiety exhibited very high antimicrobial effects (28, 31, 32). Also in some studies, compounds with cytosine like moiety has good antibacterial

activities (33, 34).

It is fascinating to notice that between all spiroaminopyran compounds (1-6) d in this study, compound 5d with spiro[aminopyran-indole] containing cytosine-like ring in its structure, has significant antimicrobial activities which is in accordance with previous published surveys.

Conclusion

In the present study, a series of spiroaminopyran derivatives were synthesized as a novel and efficient catalytic method through a one-pot, three-component reaction catalyzed by DMAP. Mainly, the significant advantages of this protocol are the excellent product yields, broader substrate scope, short reaction time, high reaction rates, easy work-up (no chromatographic separation) and benign reaction conditions (room temperature and one-pot manner).

They were evaluated for their antimicrobial activities against some gram positive and

Table 3. Spiroaminopyrans MIC ($\mu\text{g/mL}$) against clinical and standard isolates by microbroth dilution.

Compounds	MIC ($\mu\text{g/mL}$)							
	Clinical isolates				Standard strains			
	<i>S. aureus</i>	<i>S. pyogenes</i>	<i>E. coli</i>	<i>P. aeruginosa</i>	<i>S. aureus</i>	<i>S. pyogenes</i>	<i>E. coli</i>	<i>P. aeruginosa</i>
Gentamicin	4	8	16	16	4	8	32	16
1d	512 \leq	512 \leq	512 \leq	512 \leq	512 \leq	512 \leq	512 \leq	512 \leq
2d	512 \leq	512 \leq	512 \leq	512 \leq	512 \leq	512 \leq	512 \leq	512 \leq
3d	512 \leq	512 \leq	512 \leq	512 \leq	512 \leq	512 \leq	512 \leq	512 \leq
4d	512 \leq	512 \leq	512 \leq	512 \leq	512 \leq	512 \leq	512 \leq	512 \leq
5d	32	64	512	512	128	128	512 \leq	512 \leq
6d	512 \leq	512 \leq	512 \leq	512 \leq	512 \leq	512 \leq	512 \leq	512 \leq




Figure 2 . Gram staining of isolated clinical strains, a) *S. aureus*, b) *S. pyogenes*, c) *E. coli*, and d) *P. aeruginosa*.

Table 4. Inhibition zone diameter of compound 5d against *S. aureus* and *S. pyogenes*.

Species	5d disk (μg)	5d disk (μg)				Gentamicin disk (10 μg)
		16	32	64	128	
<i>S. aureus</i>	clinical isolates	10.2 \pm 1.44	12.6 \pm 1.12	15.6 \pm 1.71	19.6 \pm 1.52	21.5 \pm 1.45
	standard species	0	0	0	10.5 \pm 1.45	25.5
<i>S. pyogenes</i>	clinical isolates	0	0	10.4 \pm 0.88	12.6 \pm 0.77	14.5 \pm 0.22
	standard species	0	0	0	9.5 \pm 0.56	17

gram negative bacteria. The results showed that compound 5d has the greatest inhibitory activity against tested microorganisms, whereas the rest of the compounds have no considerable effects on microbial growth. Also, the most susceptible species were *S. aureus*. However, these compounds, aside from compound 5d, did not show a remarkable growth inhibitory activity against the gram negative bacteria. Whereas, based on the gentamicin results, it should be mentioned that the new compounds are so less potent in comparison with regular antimicrobial agents. It could be concluded that spiro[aminopyrans-indole] core with cytosine-like moiety, based on their antimicrobial activity, can be the next antibiotics generation.

- (1) Cohen ML. Changing patterns of infectious disease. *Nature* (2000) 406: 762-7.
- (2) Sabir R, Danish Alvi SF and Fawwad A. Antimicrobial susceptibility pattern of aerobic microbial isolates in a clinical laboratory in Karachi-Pakistan. *Pak. J. Med. Sc.* (2013) 29: 851-5.
- (3) Nys S, Okeke IN, Kariuki S, Dinant GJ, Driessen C and Stobberingh EE. Antibiotic resistance of faecal *Escherichia coli* from healthy volunteers from eight developing countries. *J. Antimicrob. Chemother.* (2004) 54: 952-5.
- (4) Mana M, Abdissa A and Sewunet T. Antimicrobial susceptibility pattern of bacterial isolates from wound infection and their sensitivity to alternative topical agents at Jimma University Specialized Hospital, South-West Ethiopia. *Ann. Clin. Microbiol. Antimicrob.* (2014) 13: 14-24.
- (5) Alanis AJ. Resistance to Antibiotics: Are We in the Post-Antibiotic Era. *Arch. Med. Res.* (2005) 36: 697-705.
- (6) Sistanizad M, Kouчек M, Miri M, Goharani R, Solouki M, Ayazkhoo L, Foroumand M and Mokhtari M. Carbapenem Resistance and its Effect on Bacterial Resistance in an Intensive Care unit of a Teaching Hospital. *Iran. J. Pharm. Res.* (2013) 12: 503-9.
- (7) Davies J and Davies D. Origins and evolution of antibiotic resistance. *Microbiol. Mol. Biol. Rev.* (2010) 47: 417-33.
- (8) Akbari A, Azami-Sardooei Z and Hosseini-Nia A. Synthesis and biological evaluation of 2-Amino-4H-pyran-3,4,5-tricarboxylate salt derivatives. *J. Korean Chem. Soc.* (2013) 57: 455-60.
- (9) Litvinov YM, Mortikov VY and Shestopalov AM. Versatile three-component procedure for combinatorial synthesis of 2-aminospiro[(3'H)-indol-3',4-(4H)-pyrans]. *ACS. Comb. Sci.* (2008) 10: 741-5.
- (10) Bonsignore L, Loy G, Secci D and Calignano A. Synthesis and pharmacological activity of 2-oxo-(2H)-1-benzopyran-3-carboxamide derivatives. *Eur. J. Med. Chem.* (1993) 28: 517-20.
- (11) Romdhane A and Jannet H.B. Synthesis of new pyran and pyranoquinoline derivatives. *Arab. J. Chem.* (2017) 10: S3128-S3134.
- (12) Kaminski K, Obniska J and Dybala M. Synthesis, physicochemical and anticonvulsant properties of new N-phenylamino derivatives of 2-azaspiro [4.4] nonane- and [4.5] decane-1, 3-diones: Part V. *Eur. J. Med. Chem.* (2008). 43: 53-61.
- (13) Thadhaney B, Sain D, Pemawat G and Talesara G. Synthesis and antimicrobial evaluation of ethoxyphthalimide derivatized spiro [indole-3,5'-(1,3) thiazolo(4,5-c)isoxazol]-2 (1H)-ones via ring closure metathesis. *Indian J. Chem. Section B, Organic Including Medicinal.* (2010) 49: 368-73.
- (14) Van der Sar SA, Blunt JW and Munro MH. Spiro-Mamakone A: a unique relative of the spirobisanthralene class of compounds. *Org. Lett.* (2006) 8: 2059-61.
- (15) Chande MS, Verma RS, Barve PA, Khanwelkar RR, Vaidya R and Ajaikumar K. Facile synthesis of active antitubercular, cytotoxic and antibacterial agents: a Michael addition approach. *Eur. J. Med. Chem.* (2005) 40: 1143-8.
- (16) Su BN, Mi Q, Chai HB, Riswan S, Kardono L, Ruskandi A, Farnsworth NR, Swanson SM and Kinghorn AD. Potential Anticancer Activity of Naturally Occurring and Semisynthetic Derivatives of Aculeatins A and B from *Amomum aculeatum*. *J. Nat. Prod.* (2008) 71: 390-5.
- (17) Nakao K, Ikeda K, Kurokawa T, Togashi Y, Umeuchi H, Honda T, Okano K and Mochizuki H. Effect of TRK-820, a selective kappa opioid receptor agonist, on scratching behavior in an animal model of atopic dermatitis. *Jpn. J. Pharmacol.* (2008) 28: 75-83.
- (18) Lindell S, Sanft U and Thonessen MT. Heterocyclic spiro compounds as pesticides. Int. Pat. WO/2001/011968 (App. PCT/EP2000/007851). (2001).
- (19) Sarma BK, Manna D, Minoura M and Mughesh G. Synthesis, structure, spirocyclization mechanism and glutathione peroxidase-like antioxidant activity of stable spirodiazaselenurane and spirodiazatellurane. *J. Am. Chem. Soc.* (2010) 132: 5364-74
- (20) Lee JH, Lee S, Park MY and Myung H. Characterization of thiobarbituric acid derivatives as inhibitors of hepatitis C virus NS5B polymerase. *Virol. J.* (2011) 8: 1-4.
- (21) Khurana JM and Yadav S. Highly monodispersed PEG-stabilized Ni nanoparticles: proficient catalyst for the synthesis of biologically important spiropyranes. *Aust. J. Chem.* (2012) 65: 314-9.
- (22) Chai SJ, Lai YF, Xu JC, Zheng H, Zhu Q and Zhang PF. One-Pot synthesis of spirooxindole derivatives catalyzed by lipase in the presence of water. *Adv. Synth. Catal.* (2011) 353: 371-5.
- (23) Guo RY, An ZM, Mo LP, Wang RZ, Liu HX, Wang SX and Zhang ZH. Meglumine: a novel and efficient catalyst for one-pot, three-component combinatorial

- synthesis of functionalized 2-amino-4 H-pyrans. *ACS Comb. Sci.* (2013) 15: 557-63.
- (24) Kidwa M, Jahan A and Mishra NK. Gold (III) chloride (HAuCl₄·3H₂O) in PEG: A new and efficient catalytic system for the synthesis of functionalized spirochromenes. *General. Appl. Catal.* (2012) 425: 35-43.
- (25) Ashafa A and Olajuyigbe O. Chemical Composition and Antibacterial Activity of Essential Oil of *Cosmos bipinnatus* Cav. Leaves from South Africa. *Iran. J. Pharm. Res.* (2014) 13: 1417-23.
- (26) Murugan R and Scriven E.F.V. Applications of dialkylaminopyridine (DMAP) catalysts in organic synthesis. *Aldrichimica Acta* (2003) 36:21-7.
- (27) Kidwai M, Saxena S, Khan MK and Thukral SS. Aqua mediated synthesis of substituted 2-amino-4H-chromenes and in vitro study as antibacterial agents. *Bioorg. Med. Chem. Lett.* (2005) 15: 4295-8.
- (28) Lakshmi NV, Thirumurugan P, Noorulla KM and Perumal PT. InCl₃ mediated one-pot multicomponent synthesis, anti-microbial, antioxidant and anticancer evaluation of 3-pyranyl indole derivatives. *Bioorg. Med. Chem. Lett.* (2010) 20: 5054-61
- (29) Kumar D, Reddy VB, Sharad S, Dube U and Kapur S. A facile one-pot green synthesis and antibacterial activity of 2-amino-4H-pyrans and 2-amino-5-oxo-5,6,7,8-tetrahydro-4H-chromenes. *Eur. J. Med. Chem.* (2009) 44: 3805-9.
- (30) Nandakumar A, Thirumurugan P, Perumal PT, Vembu P, Ponnuswamy MN and Ramesh P. One-pot multicomponent synthesis and anti-microbial evaluation of 2'-(indol-3-yl)-2-oxospiro(indoline-3,4'-pyran) derivatives. *Bioorg. Med. Chem. Lett.* (2010) 20: 4252-8.
- (31) Sakhuja R, Panda SS, Khanna L, Khurana S and Jain SC. Design and synthesis of spiro[indole-thiazolidine] spiro[indole-pyrans] as antimicrobial agents. *Bioorg. Med. Chem. Lett.* (2011) 21: 5465-9.
- (32) Refat HM. Synthesis and antimicrobial evaluation of some new spiroindolinone derivatives. *J. Heterocyclic Chem.* (2014) doi: 10.1002/jhet.2253.
- (33) Haveliwala DD, Kamdar NR, Mistry PT and Patel SK. Chromone-fused cytosine analogues: synthesis, biological activity and structure-activity relationship. *Nucleosides Nucleotides Nucleic Acids* (2014) 33: 80-91.
- (34) Jagessar RC and Gomathinayagam S. Antimicrobial activity of amide, N-benzoylcytosine. *J. Phys. Chem. Solids.* (2011) 1: 12-9.

This article is available online at <http://www.ijpr.ir>
