

Comparison of Kinetic Study and Protective Effects of Biological Dipeptide and Two Porphyrin Derivatives on Metal Cytotoxicity Toward Human Lymphocytes

Maryam Khosravi^a, Rahmatollah Rahimi^{a*}, Jalal pourahmad^{b*}, Mohammad Hadi Zarei^b and Mahboubeh Rabbani^a

^aBionorganic Chemistry Research Laboratory, Chemistry Department, Iran University of Science and Technology, Narmak, Tehran, Iran. ^bSchool of Pharmacy, Pharmaceutical Sciences Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Abstract

In this research, dipeptide (his- β -alanine) and porphyrin derivatives were chosen for comparing chelating ability of toxic metals such as Al³⁺, Cu²⁺, Hg²⁺ and Pb²⁺ *in-vitro*. The reason for choosing these two compounds is that both of them are naturally present in biological systems and comparison of chelating ability of these two compounds has not yet been done. Synthesis and comparison of kinetic study of dipeptide (his- β -alanine), meso-tetrakis(4-trimethylanilinium) porphyrin (TAPP) and Tetrakis(4-sulfonatophenyl)porphyrin (TPPS₄) were carried out by our team. In addition, cytotoxicity assays of metals and chelators were also performed using methylthiazole tetrazolium (MTT) test. Furthermore we investigated the protective effect of chelators against cytotoxicity, induced by different toxic metals such as Al³⁺, Cu²⁺, Hg²⁺ and Pb²⁺ on human lymphocytes. EC₅₀ values on human lymphocytes obtained after 12 h. incubation for Al³⁺, Cu²⁺ and Hg²⁺ were 30, 51, 3 μ M respectively and for Pb²⁺ no cytotoxicity was observed on human lymphocyte up to 1000 μ M concentration. EC₅₀ obtained for chelators dipeptide, TPPS₄ and TAPP were 948, 472 and 175 μ M respectively. Pretreatment of human lymphocyte with subtoxic concentrations of chelators reduced toxicity of the metals against human blood lymphocytes.

Keywords: Dipeptide; Porphyrin; Chelator; Toxic metals; Cytotoxicity; EC₅₀; lymphocytes.

Introduction

In this research, peptide and porphyrin derivatives were chosen for comparison, because all of them possess important roles in biological systems (1-9). Peptides are one of the best options for drug development due to their high specificity and low toxicity. Peptides are mostly synthesised by biological technology

or chemical methods. The chemical method, mainly solid-phase peptide synthesis (SPPS) is usually used for high production because of its simplified reaction and ordinary purification/isolation steps for the target products (10-13). Porphyrins are a group of tetrapyrrole pigments. Physical and chemical properties of porphyrins are often related to their compositions and structures. The biological role of porphyrins has been known since discovery of significantly biological macromolecules like hemoglobin and chlorophyll. The porphyrins have been applied at different fields such as photo converter, photo

* Corresponding author:

E-mail: j.pourahmadjaktaji@utoronto.ca
Rahimi_rah@iust.ac.ir

catalyst, photo sensitizer in therapeutics, diagnosis of diseases and also photo dynamic therapy and anti-Aids (HIV) drug (14-18). Complexation of peptides and porphyrins with numerous cations, their cytotoxicity and EC_{50} concentrations have so far been studied *in-vitro* and *vivo* (19-31). In this paper, we reported synthesis of dipeptide (his- β -alanine) meso-tetrakis (4-trimethylanilinium) porphyrin (TAPP) and Tetrakis (4-sulfonatophenyl) porphyrin (TPPS₄) as chelating agents for chelating of toxic metals such as Al^{3+} , Cu^{2+} , Hg^{2+} and Pb^{2+} and compared their kinetics by UV-Visible spectrophotometric methods. We also determined molecular structure of the dipeptide and porphyrin compounds using different methods such as UV-Visible spectrophotometry, FT-IR, ¹H NMR and LC-Mass spectrometry only for dipeptide structure. Furthermore, *in-vitro* cytotoxicity assay was done using MTT test and EC_{50} values for metals and chelators on human lymphocyte were obtained after 12 h. incubation. lymphocyte has fundamental roles in the immune system because they are the cells that determine the specificity of the immune response to infectious microorganisms and other foreign substances. Finally we studied protective effect of our synthesized chelators against cytotoxicity of metals on human lymphocyte and results clearly indicated that the chelators reduced toxicity of applied metals in this cellular model.

Experimental

Materials

Chemicals for dipeptide synthesis:

Wang resin, N-Fmoc-N-trityl-L-histidine were purchased from Bachem chemical Company. Boc- β -Alanine-OH was obtained from Aldrich Company, Hydroxybenzotriazole (HOBt) purchased from Fulka Company. Scavengers (anisole and phenol) solvents (trifluoroacetic acid (TFA) piperazine, N,N'-Diisopropylcarbodiimide (DIC) diethyl ether, dichloromethane, N,N-dimethyl formamide and methanol, Acetic anhydride, pyridine were obtained from Merck chemical Company.

Chemicals for porphyrin synthesis:

Propionic acid, dimethylaminobenzaldehyde, benzaldehyde, pyrrole, sulfuric acid, acetone,

sodium carbonate, methyl iodide and amberlite resin were obtained from Merck chemical Company.

Chemicals for investigation of cytotoxicity


3-(4,5-Dimethyl-2-thiazolyl) 2,5-diphenyl-2H-tetrazolium bromide (MTT) was purchased from Sigma-Aldrich Company, Ficoll-paque PLUS was obtained from Ge Healthcare Bio-Science Company. RPMI1640 and FBS Fetal Bovine serum) were purchased from Gibco, Life Technologies, Grand Island, NY. The metal salts $AlCl_3$, $CuCl_2$, $HgCl_2$ and $Pb(NO_3)_2$ were obtained from Merck chemical Company.

Sample characterization

PG Instruments T80 Double Beam UV-Visible spectrophotometer was used for UV-Visible measurements. The infrared spectra were recorded on a Shimadzu FT-IR-8400S spectrophotometer in the range of 400–4000 cm^{-1} . Proton NMR spectra were recorded on a Bruker DRX250 (300 MHz) spectrometer in water. LC-MS analyses were performed on Agilent 6410 Triple Quadruple LC-MS (USA).

Dipeptide synthesis

Dipeptide (his- β -alanine) was manually synthesized on solid phase using standard Fmoc and Boc Strategy (Scheme 1) (32, 33). Briefly, the peptide sequence (his- β -alanine) was assembled on Wang resin (1). N-Fmoc-N-trityl-L-histidine (2 equivalents) (2) and two equivalents (HOBt) and (DIC) as a coupling reagent were added to the reaction vessel. The mixture was shaken for 2 h. After, first amino acid is loaded to the resin the un-reacted sites must be end-capped with 2 equivalents acetic anhydride and 2 equivalents pyridine to ensure that future reactions do not react at those unloaded sites. Removal of the Fmoc group (3) was done by the addition of 10% Piperazine, shaken for 30 min under nitrogen atmosphere. The monitoring of the completion of the Fmoc cleavage, performed with color detector (acetaldehyde/chloranil) for detection of free terminal amino groups (4) (34). Sec amino acid Boc- β -alanine-OH (2 equivalents) (5) treated with coupling reagents (HOBt) and (DIC), which added to the resin and shook for 1.5 h at room temperature. In the last



Scheme 1. Steps of dipeptide(his- β -alanine) synthesis.

step of the synthesis, peptide was cleaved from the resin, the solvents evaporated and peptide was precipitated with diethyl ether the cleavage of peptide from resin was done using cocktail TFA/TIS/H₂O/ (82.5:5.5:2.5 v/v) for 45 min. The identity of peptide was confirmed by LC-MS.

Synthesise of meso-tetrakis(4-trimethylanilinium) porphyrin (TAPP)

Tetra-(p-dimethylaminophenyl) porphine (a) was synthesized by the method of krishnamurti (Scheme 2). A slurry of tetra-(p-dimethylaminophenyl) porphine (0.87 g) and methyl iodide (5 mL) was heated to reflux for 30 min. Then it was cooled and washed with acetone and dried in vacuume. For preparing perchlorate salt of (TAPP) (b) passing a solution of (TAPP) iodide through a column of anion exchange resin (35).


Synthesise of Tetrakis(4-sulfonatophenyl) porphyrin(TPPS₄)

TPPS₄ was prepared according to the Adler method (Scheme 3). First tetraphenylporphine (36) was synthesized, then sulfonated (35).

Optimization of concentration of metal salts for kinetic study

Different concentrations of metal salts: AlCl₃, CuCl₂, HgCl₂ and Pb(NO₃)₂, (10⁻¹, 10⁻², 10⁻³, 10⁻⁴ and 10⁻⁵M) were prepared to study the reaction kinetics with chelators [dipeptide (histidine- β -alanine), TAPP and TPPS₄]. The changes of absorption in wavelength of 214 nm (maximum wavelength of dipeptide), 416 nm (maximum wavelength of TAPP) and 412 nm (maximum wavelength of TPPS₄) were investigated by UV-Visible spectrophotometer while concentration of chelators were kept constant at $\sim 10^{-5}$ M vs. metal ions. The kinetic study was done using kinetic mode of spectrophotometer with maximum wavelength of chelators.

Extraction of lymphocyte from human blood



Scheme 2. Steps of meso-tetrakis (4-trimethylanilinium) porphyrin (TAPP) synthesis.

for cytotoxicity study in-vitro

Blood was obtained from healthy, non-smoking volunteers between 20-30 year old. Isolation of lymphocytes from human blood was done by centrifugation according to standard procedures. And lymphocytes were cultured at concentration of 1×10^4 cells per well in 96-well microplates.

Cytotoxicity study of metals and chelators on human lymphocytes in-vitro

Lymphocytes were cultured in 96-well microplates and treated with different concentrations of metals and chelators. Cells viability measured after 12 h. incubation by MTT assay. The EC_{50} values for each metals and chelators were determined.

Effect of dipeptide and porphyrin on metal cytotoxicity toward lymphocyte

Lymphocytes were treated with EC_{50} values of chelators and kept for 30 min at room temperature. After passing of this time, cells were treated by EC_{50} and $2EC_{50}$ values of metals. The Cells viability was measured after 12 h. incubation by MTT assay.

Statistical analysis

Data were analyzed using one way analysis of variance followed by post hoc Tukey test with GraphPad Prism 5 (Graphpad Software, La Jolla, CA). Results are shown as the mean \pm standard error of the mean and $P < 0.05$ was statistically significant report. For analysis of data at least

three independent experiments were used.


Results and Discussion

Characterization of synthesised dipeptide (Histidine- β -Alanine)

Dipeptide was synthesised via the standard BOC method (33). The synthesis of dipeptide (histidine- β -alanine) was structurally confirmed by UV-Visible, FT-IR, 1H NMR and LC-Mass techniques. The UV-Visible absorbance spectra of Histidine- β -Alanine was obtained in water at 25 °C, the UV absorptions appeared at 214 and 268 nm which can be related to electronic transitions of $\pi \rightarrow \pi^*$ and $n \rightarrow \pi^*$ respectively. The following spectral data for dipeptide was obtained, from the FT-IR spectra (KBr, cm^{-1}) with ν_{max} : 3238 (NH_2), 2613-3300 (OH) 1643 ($N-C=O$), 1564 ($-C=N$), and from 1H NMR spectra (300 MHz, D_2O , δ): 7.48 (imidazole ring); 4.69 (2H, CH_2N) 4.24 (d, CO_2H). The LC-MS analysis revealed a single mass peak in $[M+H]^+$ and $[M]^-$ which corresponds to the calculated molecular weight of dipeptide, $C_9H_{14}N_4O_3$, calculated: 226.23, found: $m/z(M+H)^+$: 227.000 and $m/z(M)$ 224.800. For all above data, please refer to the supplementary information.

Characterization of synthesised meso-tetrakis(4-trimethylanilinium)porphyrin (TAPP)

The synthesis of TAPP approved using UV-Visible spectra, $\lambda_{max} = 416$ nm (Soret band), 518, 556, 598, and 636 nm (Q bands); FT-IR spectra (KBr, cm^{-1}), ν_{max} : 1118 (C-N), 1342 (C=C



Scheme 3. Steps of Tetrakis(4-sulfonatophenyl)porphyrin (TPPS₄) synthesis.

exocyclic pyrrol) 1471 (C=C endocyclic pyrrol), 1606 (C=N), 2800-2900 (C-H methyl, C-H aliphatic & aromatic), 3000-3600 (N-H pyrrol, N-anilinium); ¹H-NMR spectra (300 MHz, D₂O, δ) 8.68 (s, N-H pyrrol) 8.07, 8.11 (d, phenyl) 3.80 (s, CH₃). For all above data, please refer to the supplementary information

Characterization of synthesised Tetrakis(4-sulfonatophenyl)porphyrin (TPPS₄)

The synthesis of (TPPS₄) approved using UV-Visible spectra, λ_{max} = 412 nm (Soret band) 512, 550, 578 and 632 nm (Q bands). FT-IR spectra (KBr, cm⁻¹) ν_{max}: 1000-1380 (=C-N, S=O) 1336 (C=C exocyclic pyrrol) 1382 (C=C endocyclic pyrrol) 1560 (-C=N), 2800-3010 (C-H aliphatic & aromatic) 3000-3600 (N-H pyrrol, O-H). ¹H-NMR spectra (300 MHz, D₂O, δ): 8.12 (tetraphenyl), 7.63 (tetrapyrrol) For all above data, please refer to the supplementary information.

Kinetic study of chelating reactions of dipeptide, TAPP and TPPS₄

The UV-Visible absorbance spectra was investigated to study the kinetic reactions of synthesized chelators [Histidine-β-Alanine (~10⁻⁵M), TAPP (~10⁻⁵M) and TPPS₄ (~10⁻⁵M)] with different concentrations (~10⁻¹ to ~10⁻⁵M) of metal ions (Al³⁺, Cu²⁺, Hg²⁺ and Pb²⁺). Interactions of dipeptide, TAPP and TPPS₄ with various concentrations of metal ions show that, the maximum absorption band of dipeptide (214 nm) TAPP (416 nm) and TPPS₄ (412) reduced and

shifted to short and longer wavelength in respect to the size of metal ions. we observed significant changes on the four Q bands in the visible region in such that they reduced to two bands due to the structural symmetry increase from C_{2v} to D_{4h} point groups (38). The results show at initial times (about 5-10 minutes) chelating with metals, which have same concentration with chelators (10⁻⁵M) were faster, but these reactions of metals with concentrations higher than chelators were slower. To compare the chelating ability between dipeptide and porphyrins the optimal concentration of metals was found to be ~10⁻³ M for all chelators. In terms of rate of reaction, the experimental process of chelating shows that the reactions are completed at the first ten minutes, in this regard based on the rate law the reactions was found to be first order. using the integrated first order rate law of Ln [A] = kt + Ln[A₀] the rate constant (k) was determined from the plot of -Ln [A/A₀] vs. time which gives a straight line with a slope of k. In Figure 1, it is shown that the order of rate constant increase for (Al³⁺) is such k_{dipeptide} > k_{TAPP} > k_{TPPS4}, which indicates that the reaction of dipeptide with Al³⁺ is faster than TAPP and TPPS₄. The results of rate order constant increase for other metal ions, Cu²⁺, Pb²⁺, Hg²⁺ are: (Cu²⁺) k_{TPPS4} > k_{TAPP} > k_{dipeptide}, (Pb²⁺) k_{dipeptide} > k_{TPPS4} > k_{TAPP} and (Hg²⁺) k_{TAPP} > k_{TPPS4} > k_{dipeptide}, these observations and calculated data represent that all of the chelators have high chelating potential with these metal ions. Histidine-β-Alanine is a multidentate ligand with five potential metal-coordinating sites (two N of imidazole ring,




Figure 1. Kinetics study of chelating reactions of dipeptide (histidine-β-Alanine), TAPP and TPPS₄ (10^{-5} M) with AlCl₃, CuCl₂, Pb(NO₃)₂ and HgCl₂ (10^{-3} M).

one carboxylate group, an amide linkage and a terminal amino group). From the structural study survey (39) two types of structure was introduced, the tetrahedral and octahedral complexes. About the porphyrin chelators the direct coordination between metal ions and tetrapyrrole-core also somewhat extends the conjugation from porphyrin to metal ions. According to quantum theory, an electronic excitation involved in a larger conjugated system requires lower energy absorption, corresponding to lower radiation frequency or longer wavelength (40). but the accurate configuration and chelating ability of chelators can depend on size of the metal cation, ligand-to-metal ratios, the ionic strength of the supporting solution, structure of chelator and charge density of a metal ion(39)). The Results indicate that for Al³⁺and Pb²⁺, rate constants of dipeptide is relatively higher than porphyrin derivets and for Cu²⁺ and Hg²⁺, porphyrins show maximum chelating rate.

In-Vitro assay of cytotoxicity induced by

metals, chelators and metal-chelatorcomplexes

The effects of different concentrations of metals on human lymphocyte viability were shown in Figure 2. EC₅₀ values calculated using prism software were 30, 51 and 3 μM for Al³⁺, Cu²⁺ and Hg²⁺ respectively and for Pb²⁺ no cytotoxicity was observed on lymphocyte cells up to 1000 μM concentration. The order of the cytotoxicity of metals on lymphocyte was: Hg²⁺>Al³⁺>Cu²⁺>Pb²⁺.

Evaluation of chelator's effects on cytotoxicity was shown in Figure 3. EC₅₀ measured for dipeptide, TPPS₄ and TAPP were equal to 948, 472 and 175 μM, respectively. The order of the cytotoxicity of chelators was: TAPP>TPPS₄>dipeptide.

Preventing effects of chelators against cytotoxicity of metals were shown in Figure 4 Our results show that all of chelators reduced cytotoxicity of metals on lymphocytes. Comparison of the chelators at preventing toxic metal induced cytotoxicity on human lymphocytes were: For Al³⁺(30 μM) dipeptide>TAPP>TPPS₄




Figure 2. Cytotoxicity of different concentrations of Al³⁺, Cu²⁺, Hg²⁺ and Pb²⁺ on human lymphocytes. Viability of lymphocytes after treatment with Al³⁺(A), Cu²⁺(B), Hg²⁺(C), Pb²⁺(D) for 12 h. **P<0.01 and ***P<0.001

and For Al³⁺ (60 µM) dipeptide>TAPP>TPPS₄. Based on the literature survey, the toxic preventing effect of dipeptide relates to the antioxidant properties of this compound (41). For Cu²⁺ (51 µM) TPPS₄>dipeptide~TAPP and For Cu²⁺(102 µM) TPPS₄>TAPP>dipeptide. For Hg²⁺ (3 µM) dipeptide~TAPP>TPPS₄ and For Hg²⁺(6 µM) dipeptide>TAPP~TPPS₄.

The data obtained from the kinetic studies represent that order of rate constant of Al³⁺ with chelators are: dipeptide>porphyrins, because Al³⁺ has the smallest atomic radius and the highest charge among other 3 cations, makes it capable for accepting pair electrons from N and O atoms of dipeptide (39). For Cu²⁺ metal ion, the order of rate constants is: TPPS₄>TAPP>dipeptide Which can be related to coordination of metal ion to tetrapyrrole-core in TPPS₄ that typically accept sp³d² hybrid with four orbitals in porphyrin plane and two orbitals in vertical ± z direction, coordinated with two water molecule giving rise to octahedral geometry. However, Cu²⁺ cation

may experience the so-called “Jahn-Teller effect” because of its asymmetrical d-electron configuration, which results in geometry distortion and extra binding strength (40). The generated complex between Cu²⁺ and dipeptide is such that two molecules of dipeptide with one molecule of water makes it to be five coordination complex. The four nearest ligand atoms are the terminal amino nitrogen, the amide nitrogen, a carboxylate oxygen of one of the dipeptide molecules, and the nitrogen of the imidazole from the sec molecule. this shows that each dipeptide molecule has potential of bounding to two different Cu²⁺ cation centers (20). For Hg²⁺ cation coordination with chelators, rate order constants are TAPP>TPPS₄>dipeptide by this order the monomeric structure of Hg²⁺ complexes with dipeptide shows the lower stability, we assume that the complex can possess a polymeric structure [HgLH]_n²⁺. In this regard the mercury complexation with dipeptide reverses the tautomeric preference between protonation of




Figure 3. Cytotoxicity of different concentrations of TAPP, TPPS₄ and dipeptide on human lymphocytes. Viability of lymphocytes after treatment with TAPP (A) TPPS₄ (B) and dipeptide (C) for 12 h. *P<0.05, **P<0.01 and ***P<0.001

the N-atoms of the imidazole ring (20). The rate constants order for Pb²⁺ cation complexation with ligands are: dipeptide > porphyrins, because Pb²⁺ has lower ratio of charge to radius, therefore can be easily coordinated to dipeptide potential coordinating sites. The strength of porphyrin-metal ion interaction may depend on different factors, including the porphyrin substitutions, charge state (such as H₂P or P²⁻-H₄P²⁺) and type of metal ion, the factors may be more than these but in this work, some of them considered as an effective factor. In supporting of this discussion reference (4) has useful information regarding to the binding of N-alkylated porphyrin HN-Me-TPPS with cations Cd²⁺ and Zn²⁺ shows the following rate constants: 1.3 × 10⁻² and 3.3 × 10⁻¹, respectively, this is exactly what we observed during the experimental procedure. The binding trends revealed in these quantitative data are consistent with our qualitative predictions, although in comparing with some literature surveys, there might be some differences, this is

unavoidable because they might use basically different materials and experimental methods. The findings *in-vitro* on human lymphocytes suggested that greater concentrations than EC₅₀ of metal ions, Pb²⁺ (1000 µM), Cu²⁺ (51 µM) Al³⁺ (30 µM) and Hg²⁺ (3 µM) can decrease Viability% of lymphocytes. For comparing chelators effects against cytotoxicity of metals, we treated lymphocytes with EC₅₀ and 2EC₅₀ values of metals. The results indicated that chelators reduced toxicity of metals against human lymphocytes. Regarding the Pb²⁺ no cytotoxicity was determined on lymphocytes up to 1000 µM concentration. The lymphocyte cytotoxicity with Pb²⁺ chelator complexes were even lower than Pb²⁺ (data not shown). The difference between chelators effects against cytotoxicity of metals can correspond with reasons that have been described about kinetic study. Positive results about protective effects of dipeptide (his-β-alanine) and two porphyrin derivatives on metal cytotoxicity toward human




Figure 4. Comparison of protective effects of chelators on cytotoxicity of toxic metals. Protective effects of different chelators on cytotoxicity induced by EC₅₀ and 2EC₅₀ of Al³⁺ (A), Cu²⁺ (B) and Hg²⁺ (C) on human lymphocytes. *P<0.05, **P<0.01 and ***P<0.001

lymphocytes and numerous studies that have been done by many researchers, acknowledge the importance of these compounds in drug developments. his-β-alanine, a naturally occurring di-peptide, is present in large amounts in long-lived human tissues. Numerous evidence have indicated the multi-functionality of this dipeptide in the human body (4,34) such as physiological buffering agent and a metal ion (e.g., zinc and copper) chelator, regulator of the amount of transition metal ions in biological fluids and tissues. Besides, it has ability to form complexes with metal ions (42-45). his-β-alanine has been used as an eye-drop component used for the treatment and inhibition of senile cataract (46). In addition, his-β-alanine has been shown to suppress the accumulation of amyloid beta peptide in the central nervous system of transgenic mouse model for Alzheimer’s disease (47). and attenuate the *in-vitro* fibrillogenesis of amyloid beta peptide (1–44) (48). It is a

potent scavenger of both reactive oxygen species (ROS) and reactive nitrogen species (RNS) which excite peroxidation of unsaturated lipids present in membranes as well as of toxic reactive α, β-unsaturated aldehydes deriving from this oxidation (49,50). Possessing anti-aging functions (51,52), his-β-alanine illustrated a well-documented anti-glycating activity in proteins, including low-density lipoproteins, glucose decline products, esterase, and histones (53-54). his-β-alanine was reported to increase the thermal unfolding and water availability of glycated protein species (55). It also mitigates and/or prevents the variation in electrophoretic dynamism operated by glyceraldehyde 3-phosphate (56). A study in 2006 by Mahmood *et al.* showed that Zn and the antioxidant his-β-alanine can stabilize the entirety of the small bowel and motivate repair processes in the gut, both *in-vitro* and *vivo* models (57). It has been reported that Zn and his-β-alanine may induce

anti-oxidative stress enzymes *in-vitro* and *in-vivo* model (58). One of the long-standing goals of both researchers and oncologists is to create a framework for the complete cure of cancer with less toxic effect and make better Quality of life (QOL). Experiments to test bioactivation of neutral reagents by light led to modern photodynamic therapy. The priority of photodynamic therapy has no offense during treatment and the selective agglomeration of photosensitizers in tumor tissue. Porphyrin and its analogues are the most effective photosensitizers for PDT. These photosensitizers have a strong absorption band ($\epsilon 5 \times 10^5 \text{ M}^{-1} \text{ cm}^{-1}$) around 400 nm called the Soret band and weak absorption bands between 500 and 800 nm named Q bands. Despite the large molar absorption coefficient, the Soret band is not adequate for PDT of deeper tumor tissues. The Q1 band (600–800 nm) is generally used for PDT (59-60). Anti-HIV-1 activity of the porphyrins was determined by multiple nuclear activation of galactosidase indicator cell (MAGI) assay. Cytotoxicity of different kind of porphyrins are determined. Interestingly, the cytotoxicity of the iron-(III) complexes was lower than that of the corresponding porphyrin free bases. A similar effect on the cytotoxicity was reported by Song *et al.* The introduction of the iron atom might improve the selectivity of the binding of the porphyrin to the V3 loop region. The low cytotoxicity and high anti-HIV-1 activity of these iron(III) porphyrins are a large benefit to the usage of these metal complexes (62)). Our results also showed that pretreatment of lymphocytes with chelators clearly indicate that all of the chelators reduce toxicity of the metals on lymphocytes. Viability % of human lymphocytes following addition of dipeptide was higher than those of porphyrin derivatives.

Conclusions

In this work, synthesis of dipeptide (his- β -alanine) was done by solution phase peptide synthesis (SPS) and solid phase peptide synthesis (SPPS) on Wang resin. Study of chelating properties of dipeptide (his- β -alanine) and porphyrin derivatives by UV-Visible absorbance spectra revealed that chelating activity depends on concentration of metal, physicochemical

properties of metal and finally molar ratio of metal and chelator. The order of the chelator strength at adsorbing Al^{3+} (10^{-3} M) and Pb^{2+} (10^{-3} M) was: dipeptide > porphyrin compounds. It is probably due to $k_{\text{dipeptide}} > k_{\text{porphyrin}}$. For Cu^{2+} (10^{-3} M) and Hg^{2+} (10^{-3} M): $k_{\text{porphyrin}} > k_{\text{dipeptide}}$. Results of pretreatment of lymphocytes with chelators clearly indicate that all of the chelators reduce toxicity of the metals on lymphocytes. Viability % of human lymphocytes following addition of dipeptide was higher than those of porphyrin derivatives.

Acknowledgements

This study was financially supported by Iran University of Science and Technology. We also extend our sincere gratitude for the assistance of Pharmaceutical Sciences Research Center, Shahid Beheshti University of Medical Sciences.

References

- (1) Makhro AV, Mashkina AP, Solenaya OA, Trunova OA, Tyulina OV, Bulygina ER and Boldeyev AA. Carnosine protects cells from oxidative stress induced by hyperhomocysteinemia. *Neurochemical. J.* (2008) 2: 202-8.
- (2) Guiotto A, Calderan A, Ruzza P and Borin G. Carnosine and Carnosine-Related Antioxidants. *A review. Cur. Med. Chem.* (2005) 12: 2293–315.
- (3) Janssen B, Hohenadel D, Brinkkoetter P, Peters V, Rind N, Fischer C, Rychlik I, Cerna M, Romzova M, de Heer E, Baelde H, Bakker SJ, Ziric M, Rondeau E, Mathieson P, Saleem MA, Meyer J, Köppel H, Sauerhoefer S, Bartram CR, Nawroth P, Hammes HP, Yard BA, Zschocke J and Van der Woude FJ. Carnosine as a Protective Factor in Diabetic Nephropathy. *Diabetes* (2005) 54: 2320-7.
- (4) Hipkiss AR. Carnosine, a protective, anti-ageing peptide. *J. Biochem. Cell Biol.* (1998) 8: 863–8.
- (5) Schroder L, Schmitz CH and Bachert P. Carnosine as molecular probe for sensitive detection of Cu(II) ions using localized ^1H NMR spectroscopy. *J. Inorg. Biochem.* (2008) 102: 174-83.
- (6) Jia H, Qi X, Fang S, Jin Y, Han X, Wang Y, Wang A and Zhou H. Carnosine inhibits high glucose-induced mesangial cell proliferation through mediating cell cycle progression. *Regul. Pept.* (2009) 154: 69-76.
- (7) Wu W, Liu N, How SC, Chen WA, Lai CM, Liu HS, Hu CJ and Wang SS. Carnosine's effect on amyloid fibril formation and induced cytotoxicity of lysozyme. *Plos. One.* (2013) 8: e81982.
- (8) Jie L, Shuo S, Liang-Nian J and Wen-Jie M. Investigation on DNA Binding and Photo-Cleavage Properties of

- Water-Soluble Porphyrin and Metalloporphyrins. *Transition Metal Chemistry*.(2005) 30: 1684-90.
- (9) Kim JO, Lee YA, Jin B, Park T, Song R and Kim SK. Binding mode of cationic monomer and dimer porphyrin with native and synthetic polynucleotides studied by polarized light spectroscopy. *Biophys. Chem.* (2004) 111: 63-71.
- (10) Muresan AZ and Lindsey JS. Design and synthesis of water-soluble bioconjugatable trans-AB-porphyrins. *Tetrahedron* (2008) 64: 11440-48.
- (11) Tae-Kyung L, Sun-Jong R and Yoon-Sik L. A new method for the preparation of 2-chlorotriyl resin and its application to solid-phase peptide synthesis. *Tetrahedron Lett.* (2007) 48: 389-91.
- (12) Waqar A. Solid-phase total synthesis of cyclic pentapeptide Longicalycinin A, by using 2-chlorotriyl chloride resin. *J. Cancer. Res. And Exp. Oncol.* (2013) 5: 8-19.
- (13) Bacsa B, Desai B, Dibo G and Kappe CO. Rapid solid-phase peptide synthesis using thermal and controlled microwave irradiation. *J. Pept. Sci.* (2006) 12: 633-8.
- (14) Rahimi R, Jeddi M and Ravanfar SMR. Characterization of Nano Tetrakis (4-sulfonatophenyl) porphyrin Aggregation Through Atomic Force Microscopy and UV-Visible Spectroscopy Methods. *Asian J. Chem.* (2009) 21: 3988-94.
- (15) Rodriguez L, De Bruijn HS, Di Venosa G, Mamone L, Robinson DJ, Juarranz A, Batlle A and Casas A. Porphyrin synthesis from amino levulinic acid esters in endothelial cells and its role in photodynamic therapy. *J. Photochem. Photobiol. B.* (2009) 96: 249-54.
- (16) Van Duijnhoven FH, Aalbers RI, Rovers JP, Terpstra OT and Kuppen PJ. The immunological consequences of photodynamic treatment of cancer, a literature review. *Immunobiology* (2003) 207: 105-13.
- (17) Wang HM, Jiang JQ, Xiao JH, Gao RL, Lin FY and Liu XY. Porphyrin with amino acid moieties a tumor photosensitizer. *Chem. Biol. Interact.* (2008) 172: 154-8.
- (18) Shigenobu Y, Shiho H, Makoto O, Yuichiro H, Shunichiro O, Atsushi I, Hiromi K, Mamoru T and Takashi J. Current states and future views in photodynamic therapy. *J. Photochem. Photobiol. C. Photochem. Rev.* (2011) 12: 46-67.
- (19) Baran E and Diez RP. A density functional study of some physical properties of Carnosine (N- β -alanyl-L-histidine). *J. Mol. Struct.* (2003) 621: 245-51.
- (20) Baran EJ. Metal complexes of Carnosine. *Biochemistry (Mosc.)*. (2000) 65: 789-97.
- (21) Nyarko E, Hara T, Grab DJ, Tabata M and Fukuma T. Toxic effects of mercury(II), cadmium(II) and lead(II) porphyrins on *Trypanosoma brucei* growth. *Chemico-Bio Intera.* (2002) 139: 177-85.
- (22) Basu N, Scheuhammer AM, Rouvinen-Watt K, Evans RD, Trudeau V L and Chan LHM. *In-vitro* and whole animal evidence that methylmercury disrupts GABAergic systems in discrete brain regions in captive mink. *Comparative Biochemistry and Physiology, Part C.* (2010) 151: 379-85.
- (23) Varasteh Z, Aberg O, Velikyan I, Lindeberg G, Sörensen J, Larhed M, Antoni G, Sandström M, Tolmache V and Orlova A. *In-Vitro* and *In-Vivo* Evaluation of a ¹⁸F-Labeled High Affinity NOTA Conjugated Bombesin Antagonist as a PET Ligand for GRPR-Targeted Tumor Imaging. *LOS ONE.* (2013) 8: e81932.
- (24) Baranauskienė D, Stumylaitė L, Kregždytė IR, Ryselis S, Abdrakhmanovas O and Stepaniukas A. Effects of aluminum on delta aminolevulinic acid dehydratase *in vivo* and *in vitro*. *Vetinarija Ir Zootechnika (Vet Med Zoot).* (2009) 48: 3-8.
- (25) Vrouenraets MB, Visser GW, Stigter M, Oopelaar H, Snow GB and Dongen GA. Comparison of aluminium(III) phthalocyanine tetrasulfonate- and meta-tetrahydroxy phenylchlorin-monoclonal antibody conjugates for their efficacy in photodynamic therapy *in vitro*. *Int. J. Cancer* (2002) 98: 793-8.
- (26) Shen Y, Yang J, Li J, Shi X, Ouyang L, Tian Y and Lu J. Carnosine inhibits the proliferation of human gastric cancer SGC-7901 cells through both of the mitochondrial respiration and glycolysis pathways. *PLOS ONE.* (2014) 9: e104632.
- (27) Ooi TC, Mohammad NH and Sharif R. Zinc Carnosine Protects Against Hydrogen Peroxide-Induced DNA Damage in WIL2-NS Lymphoblastoid Cell Line Independent of Poly (ADP-Ribose) Polymerase Expression. *Biol. Trace. Elem. Res.* (2014) 162: 8-17.
- (28) Buchtik R, Travnicek Z, Vanko J, Herchel R and Dvorak Z. Synthesis, characterization, DNA interaction and cleavage, and *in-vitro* cytotoxicity of copper (II) mixed-ligand complexes with 2-phenyl-3-hydroxy-4(1H)-quinolinone. *Dalton Trans.* (2011) 40: 9404-12.
- (29) Fadda AA, El-Mekawy RE, El-Shafei A, Freeman HS, Hinks D and El-Fedawy M. Design, Synthesis, and Pharmacological Screening of Novel Porphyrin Derivatives. *J. Chem.* (2013): 1-11.
- (30) Pourahmad J, Hosseini MJ, Eskandari MR and Rahmani F. Involvement of four different intracellular sites in chloroacetaldehyde- induced oxidative stress cytotoxicity. *Iran. J. Pharm. Res.* (2011) 11: 265-76.
- (31) Pourahmad J and Hosseini MJ. Application of isolated mitochondria in toxicological studies. *Iran. J. Pharm. Res.* (2012) 11: 703-4.
- (32) Ahmad W, Shaheen F, Zhou Y and Zhang L. Solid-phase total synthesis of cyclic pentapeptide Longicalycinin A, by using 2-chlorotriyl chloride resin. *J. Cancer Res. Exp. Oncol.* (2013) 5: 8-19.
- (33) Khoshbakht S, Kobarfard F, Beiki D, Sabzevari O, Amini M, Mehrnejad F, Tabib K and Shahhosseini S. HYNIC a bifunctional prosthetic group for the labelling of peptides with ^{99m}Tc and ¹⁸FDG. *J. Radioanal Nucl Chem. J. Radioanal. Nucl. Chem.* (2016) 307: 1125-3419.
- (34) Hipkiss AR. Carnosine, a protective, anti-ageing peptide. *J. Biochem. Cell Biol.* (1998) 30: 863-8.
- (35) Krishnamurthy M. Synthesis and characterization of a new water-soluble porphyrin. *Ind. J. Chem.* (1977) 10: 964-66.

- (36) Longo A, Adler FR, Finarelli JD, Goldenmacher J and Assour J. A simplified synthesis for meso-tetraphenylporphine. *J. Org. Chem.* (1967) 32: 476-80.
- (37) Hokland P and Heron I. Analysis of the lymphocyte distribution during Isopaque-Ficoll isolation of mononuclear cells from human peripheral blood. *J. Immunol. Methods* (1980) 32: 31-9.
- (38) Huang X, Nakanisha K and Berova N. Porphyrins and Metalloporphyrins- Versatile Circular Dichroic Reporter Groups for Structural Studies. *Chirality* (2000) 12: 237-55.
- (39) Branham ML, Singh P, Bisetty K, Sabela M and Govender T. Preparation, Spectrochemical, and Computational Analysis of L-Carnosine (2-[(3-Aminopropanoyl)amino]-3-(1H-imidazol-5-yl) propanoic Acid) and Its Ruthenium(II) Coordination Complexes in Aqueous Solution. *Molecules* (2011) 16: 10269-91.
- (40) Song Z, Adeyemo AO, Baker J, Traylor SM and Lightfoot ML. Structure Of Porphyrin TPPS4 and its interaction with metal ions as elucidated by 1H NMR and UV-Visible spectra. *Ga. J. Sci.* (2011) 69: 89-101
- (41) Cacciatore I, Cocco A, Costa M, Fontana M, Lucente G, Pecci L and Pinnen F. Biochemical properties of new synthetic Carnosine analogues containing the residue of 2, 3-diaminopropionic acid: the effect of N-acetylation. *Amino Acids* (2005) 28: 77-83.
- (42) Boldyrev AA. Carnosine and Oxidative Stress in Cells and Tissues. *Nova Science Publisher*, New York. (2006).
- (43) Stvolinsky SL and Dobrota D. Anti-ischemic Activity of Carnosine. *Biochemistry (Moscow)* (2000) 65: 998-1005.
- (44) Singh SR, Carreiro ST, Chu J, Prasanna G, Niesman MR, Collette Iii WW, Younis HS, Sartnurak S and Gukasyan HJ. L-carnosine: multifunctional dipeptide buffer for sustained-duration topical ophthalmic formulations. *J. Pharmacy Pharmacol.* (2009) 61: 733-42.
- (45) Torreggiani A, Fini G. and Bottura G. Effect of transition metal binding on the tautomeric equilibrium of the carnosineimidazolic ring. *J. Mol. Structure* (2001) 565: 341-6.
- (46) Babizhayev MA. Designation of imidazole-containing dipeptides as pharmacological chaperones. *Hum. Exp. Toxicol.* (2011) 30: 736-61.
- (47) Corona C, Frazzini V, Silvestri E, Lattanzio R, La Sorda R, Piantelli M, Canzoniero LM, Ciavardelli D and Rizzarelli E. Effects of Dietary Supplementation of Carnosine on Mitochondrial Dysfunction, Amyloid Pathology, and Cognitive Deficits in 3xTg-AD Mice. *Plos. One* (2011) 6: e17971.
- (48) Attanasio F, Convertino M, Magno A, Caffisch A, Corazza A, Haridas H, Esposito G, Cataldo S, Pignataro B, Milardi D and Rizzarelli E. Carnosine Inhibits A beta 42 Aggregation by Perturbing the H-Bond Network in and around the Central Hydrophobic Cluster. *Chembiochem.* (2013) 14: 583-92.
- (49) Fu HY, Katsumura Y, Lin MZ, Muroya Y, Hata K, Fujii K, Yokoya A and Hatano Y. Free radical scavenging and radioprotective effects of carnosine and anserine. *Radiat. Phys. Chem.* (2009) 78: 1192-97.
- (50) Guiotto A, Calderan A, Ruzza P and Borin G. Carnosine and Carnosine-Related Antioxidants: A review. *Curr. Med. Chem.* (2005) 12: 2293-315.
- (51) Hipkiss AR. Would Carnosine or a Carnivorous Diet Help Suppress Aging and Associated Pathologies? *Ann. N.Y. Acad. Sci.* (2006) 1067: 369-74.
- (52) Stuerenburg HJ. The roles of carnosine in aging of skeletal muscle and in neuromuscular diseases. *Biochemistry-Moscow* (2000) 65: 862-5.
- (53) Rashid I, Van Reyk DM and Davies MJ. Carnosine and its constituents inhibit glycation of low-density lipoproteins that promotes foam cell formation *in-vitro*. *Febs. Lett.* (2007) 581: 1067-70.
- (54) Alhamedani MSS, Al-Azzawie HF and Abbas FKH. Decreased formation of advanced glycation end-products in peritoneal fluid by carnosine and related peptides. *Perit. Dial. Int.* (2007) 27: 86-9.
- (55) Yeagans GS and Seidler NW. Carnosine promotes the heat denaturation of glycated protein. *Biochem. Biophys Res. Commun.* (2003) 300: 75-80.
- (56) Seidler NW. Carnosine prevents the glycation-induced changes in electrophoretic mobility of aspartate aminotransferase. *J. Biochem. Mol. Toxicol.* (2000) 14: 215-20.
- (57) Mahmood A, Fitz Gerald AJ, Marchbank T, Ntatsaki E, Murray D, Ghosh S and Playford RJ. Zinc carnosine, a health food supplement that stabilises small bowel integrity and stimulates gut repair processes. *Gut.* (2007) 56: 168-75.
- (58) Ueda K, Ueyama T, Oka M, Ito T, Tsuruo Y and Ichinose M. Polaprezinc (zinc L-carnosine) is a potent inducer of anti-oxidative stress enzyme, hemeoxygenase (HO)-1—a new mechanism of gastric mucosal protection. *J. Pharmacol. Sci.* (2009) 110: 285-94.
- (59) Yanoa Sh, Hiroharab Sh, Obatac M, Hagiyaad Y, OguradSh-ic, Ikedab A, Kataokae H, Tanakae M and Johe T. Current states and future views in photodynamic therapy. *J. Photochem. Photobiol. C: Photochem. Rev.* (2011) 12: 46- 67.
- (60) Malina L, Tomankova KB, Malohlava J, Jiravova J, Manisova B, Zapletalova J and Kolarova H. The *in-vitro* cytotoxicity of metal-complexes of porphyrinsensitizer intended for photodynamic therapy. *Toxicol. in-Vitro* (2016) 34: 246-56.
- (61) Sobczynski J, Kristensena S and Berga K. The influence of Pluronicnanovehicles on darkcytotoxicity, photocytotoxicity and localization of four model photosensitizers in cancer cells. *Photochem. Photobiol. Sci.* (2014) 13: 8-22.
- (62) Watanabe K, NegiSh, Sugiura Y, Kiriyama A, Honbo A, Iga K, Kodama EN, Naitoh T, Matsuoka M and Kano K. Binding of Multivalent Anionic Porphyrins to V3 Loop Fragments of an HIV-1 Envelope and Their Antiviral Activity. *Chem. Asian. J.* (2010) 5: 825 -34.

This article is available online at <http://www.ijpr.ir>