

Investigation of Antioxidative, Cytotoxic, Membrane-Damaging and Membrane-Protective Effects of The Essential Oil of *Origanum majorana* and its Oxygenated Monoterpene Component Linalool in Human-Derived Hep G2 Cell Line

Ayşe Erdoğan^{a*} and Aysun Özkan^a

^aDepartment of Biology, Faculty of Science, University of Akdeniz, 07058, Antalya, Turkey.

Introduction

In all aerobic organisms, including human beings, production of reactive oxygen species (ROS) is balanced by antioxidant defense system. A serious imbalance between the production of ROS and antioxidant defense system is responsible for oxidative stress. Thus, ROS plays an important role in the etiology of many diseases and aging. Antioxidant defense system which prevents oxidative damages of ROS consist of flavanoids, carotenoids, phenolic compounds, vitamins, and antioxidant enzymes (1). Although there are some synthetic antioxidants such as butylated hydroxytoluene (BHT) and butylated hydroxyanisole (BHA) which is commonly used in processed food, it has been reported that these compounds have some side effects (2). Recently, there has been increasing interest in finding plants with high antioxidant capacities since they can protect human body from free radicals and retard the progression of many chronic diseases (3-5).

Various medicinal properties have been ascribed to natural herbs. Medicinal plants constitute the main source of new pharmaceuticals and healthcare products. Plant products are also known to possess potential for food preservation.

Sweet marjoram [*Origanum majorana* L. syn. *Majorana hortensis* Moench], a member of

the Lamiaceae family, is of great economic and industrial importance (6). Lamiaceae consists of more than 150 species occurring mainly in the Mediterranean area and are used as herbal teas for their (folk) medicinal properties. *Origanum* is represented in Turkey by 22 species or 32 taxa, 21 being endemic to Turkey. Out of 52 known taxa of *Origanum*, 60% are recorded to grow in Turkey (7). Lamiaceae plants were widely studied as natural antioxidant sources because of their high contents of polyphenols (8).

O. majorana is one of the most important aromatic plants that contain major antioxidants such as flavonoids and triterpenoids (9). Phenolic acids and flavonoids have been reported to play a role in the prevention of human pathologies (10). Food and Drug Administration regard *O. majorana* to be generally safe.

Genus *Origanum* is used as anti-diabetic, carminative, tonic, digestive, stimulant, expectorant, menstrual regulator, diuretic, and for respiratory problems such as asthma. Aqueous and ethanolic extracts from marjoram played a role in preventing carcinogenesis and oncogenic mutations (11). These extracts caused alterations in methylguanine-DNA methyltransferase activity which is highly expressed in human cancers. *O. majorana*'s crude drug extracts exerted a low cytotoxicity on five human liver-cancer cell lines studied with an average of 39.1% (12). Commercial *O. majorana* oil is used as a spice and condiment. The oil is used in perfumery for its spicy herbaceous notes (13) and as fungicides or insecticides in pharmaceutical

* Corresponding author:

E-mail: ayseerdogan@akdeniz.edu.tr

and industrial products (14).

Linalool is monoterpene compound reported to be major volatile components of the essential oils of several aromatic species. Linalool (3,7-dimethyl-1,6-octadien-3-ol) is one of the commonly used chemicals in cosmetic products and the fragrance and perfume industry. Safety evaluation studies revealed that linalool is not irritating, phototoxic or sensitizing but has a low order of acute toxicity (15).

Hepatoma G2 cells (Hep G2) are considered a good model to study *in vitro* xenobiotic metabolism and toxicity to the liver, since they retain many of the specialized functions which characterize normal human hepatocytes (16). Also, Hepatoma G2 cells are a valuable model to study hepatocellular carcinoma and the liver, where drugs are metabolized (17).

In another experiment, we studied phenolic terpenoid carvacrol's (main component of *O. majorana* essential oil) cytotoxicity, cytoprotective and membrane protective effects on Hep G2 cells (18). Therefore, we used linalool (second main component of *O. majorana* essential oil) for this experiment.

The aim of our work was to compare cytotoxic and membrane-damaging effects of *O. majorana* essential oil and its oxygenated monoterpene component linalool on the human hepatocellular carcinoma cell line Hep G2 and to investigate their possible protective (antioxidant) effects against H₂O₂ induced membrane damage.

Experimental

Collection of plant material

O. majorana was collected from Mahmutseydi Village, Alanya (730 m) in Turkey (Voucher no: TR 102 AL). The taxonomic identification of plant materials were confirmed by a plant taxonomist, Prof. Dr. R. Süleyman Göktürk, in Department of Biology, Akdeniz University, Antalya, Turkey.

Isolation of the essential oil

The air-dried aerial parts of plants (100 g) were powdered and subjected to hydrodistillation for 3 h using the Clevenger-type apparatus (ILDAM Ltd., Ankara, Turkey) at Molecular Biology Department in Biology in Akdeniz University.

The obtained essential oil was dried over anhydrous sodium sulphate and after filtration, stored at +4°C until tested and analysed. Linalool (purity 97%) was purchased from Aldrich.

Gas chromatography mass spectrometry (GC-MS)

The composition of the volatile constituents was established by GC-MS/quadruple detector analyses using a Shimadzu QP 5050 system, equipped with a chemically bonded fused silica column Cp WAX 52 CB (50 m × 0.32 mm internal diameter, 1.2 µm film thickness). GC analysis was employed under the following conditions: Injector temperature was 240°C. Detector temperature was 250 °C. The temperature program for the Cp WAX 52 CB column was from 60 °C to 220 °C at a rate of 2 °C/min and then held at 220 °C for 20 min. Helium was used as a carrier gas at a flow 10 psi and injection volume of each sample was 1 mL, split ratio (1:20). The MS conditions were: ionization energy, 70 eV; mass range, 40–400 amu; scan mode, EI. The percentage composition was computed from the GC peak areas according to the 100% method without using any correction factors. The identification of the components was based on the comparison between their mass spectra and those of Wiley and Nist, Tutore Libraries.

The components were identified based on the comparison between their relative retention time as well as mass spectra and those of standards and literature data (19, 20). The results were also confirmed by the comparison between the compounds elution order and their relative retention indices on non-polar phases reported in the literature (19, 20). All experiments were carried out at least in triplicate. The results were reported as mean values.

Antioxidant Activity

DPPH assay

The hydrogen atoms or electrons donation ability of the corresponding oils and some pure compounds were measured from the bleaching of purple coloured methanol solution of 2,2-diphenyl-1-picrylhydrazyl (DPPH). This spectrophotometric assay uses stable radical DPPH as a reagent (21, 22). Various

concentrations (50 μL) of the essential oil in methanol were added to 5 mL of a 0.004% (w/v) methanol solution of DPPH. After a 30 min incubation period at room temperature the absorbance was read against a blank at 517 nm. Inhibition free radical DPPH in percent (I%) was calculated in following way:

$$I\% = (A_{\text{control}} - A_{\text{sample}} / A_{\text{control}}) \times 100$$

Where A_{blank} is the absorbance of the control reaction (containing all reagents except the test compound), and A_{sample} is the absorbance of the test compound. Essential oil and positive controls concentrations providing 50% inhibition (EC_{50}) were calculated from the graphs. For the calculation of these values, Microsoft Excel software was used. Tests were carried out in triplicate.

β -carotene–linoleic acid assay

In this assay antioxidant capacity is determined by measuring the inhibition of the volatile organic compounds and the conjugated diene hydroperoxides arising from linoleic acid oxidation (23). A stock solution of β -carotene–linoleic acid mixture was prepared as following: 0.5 mg β -carotene was dissolved in 1 mL of chloroform (HPLC grade), 25 μL linoleic acid and 200 mg Tween 40 was added. Chloroform was completely evaporated using a vacuum evaporator. Then 100 mL distilled water saturated with oxygen (30 min, 100 mL/min) was added with a vigorous shaking.

This reaction mixture (2500 μL) was dispersed to test tubes and 350 μL portions of the essential oil prepared at 2 g/L concentrations added and also the emulsion system was incubated up to 24 h at room temperature. The same procedure was repeated with synthetic antioxidant, butylated hydroxytoluene (BHT), ascorbic acid and α -tocopherol as positive controls, and a blank. After this incubation period, the absorbance of the mixtures was measured at 490 nm. Antioxidative capacity of the essential oil was compared with those BHT, ascorbic acid, -tocopherol and blank. Tests were carried out in triplicate.

Cell lines and cell culture

The hepatoma G2 cells (Hep G2) line (purchased from American Type Culture

Collection) was used in this study. Cells were grown in Minimum Essential Medium (MEM) supplemented with 10% (v/v) fetal bovine serum, 1% (v/v) antibiotic-antimycotic solution in a humidified atmosphere containing 5% CO_2 at 37°C. For subculturing, cells were harvested after trypsin/ethylenediaminetetraacetic acid treatment at 37°C. Cells were used when monolayer confluence had reached 75%.

Cytotoxicity assay

The cells were seeded into 96-well microplates (1×10^4 cells well⁻¹) for 24 h and then treated with different concentrations of the essential oil and linalool (5-500 $\mu\text{g}/\text{mL}$) for 24, 48 and 72 h. Cytotoxicity of the essential oil and linalool was assayed by CellTiter-Blue® Cell Viability Assay. The assay is based on the ability of living cells to convert a redox dye (resazurin) into a fluorescent end product (resofurin). Nonviable cells rapidly lose metabolic capacity and thus do not generate a fluorescent signal (24). Following cellular reduction, fluorescence is recorded at 560 nm excitation/590 nm emissions. The data were expressed as average values obtained from eight wells for each concentration. The IC_{50} value was calculated from equation of graph. H_2O_2 cytotoxicity on cancer cells was measured in the same way. For measuring antioxidant effect of the essential oil and its oxygenated monoterpene component linalool against H_2O_2 cytotoxicity, different concentrations of the cells were preincubated (IC_{10} , IC_{20} and IC_{30}) for 1 h, before H_2O_2 treatment (IC_{10} , IC_{50} and IC_{70}) for 24 hour. Essential oil and its oxygenated monoterpene component linalool were dissolved in 0.5% (v/v) dimethyl sulphoxide (DMSO).

Determination of malondialdehyde level

Malondialdehyde (MDA) levels were determined after Hep G2 cells were exposed to different concentration of essential oil and its oxygenated monoterpene component linalool for 24 h (IC_{10} , IC_{50} and IC_{70}). For measuring membrane protective effects of the essential oil and linalool against H_2O_2 , the cells were preincubated with different concentrations (IC_{10} , IC_{20} and IC_{30}) of essential oil and linalool (IC_{10} , IC_{20} and IC_{30}) for 1 h, before H_2O_2 treatment (IC_{10} , IC_{50} and IC_{70}) for 24 h. Essential oil and

Table 1. Chemical composition of the essential oil from *O. majorana*.

Components	RT	Composition (%)
γ -Terpinene	17.4	0.9
Cymene	18.4	0.8
Thymol	65.8	0.3
Carvacrol	67.3	52.5
Linalool	39.3	45.4
Total		99.9

%, relative compound percentage.

RT, retention time (min).

its oxygenated monoterpene component linalool were dissolved in 0.5% (v/v) DMSO.

Hep G2 cells were plated at a density 15×10^4 cell/100 mm dishes. The cells were scraped off culture plates with culture medium and were centrifuged $600 \times g$ for 10 min. The cell pellets were washed with PBS and then sonicated (3×15 sec) in 50 mM potassium phosphate, pH 7.2, containing 1 mM PMSF and 1 μ g/mL of leupeptin and centrifuged at $150,000 \times g$ for 45 min. The supernatant was used for the determination of malondialdehyde level.

Malondialdehyde levels in Hep G2 cells were assayed as described in the previous method (25). This fluorometric method for measuring thiobarbituric acid-reactive substances (TBARS) in supernatant is based on the reaction between malondialdehyde and thiobarbituric acid. The product of this reaction was extracted into butanol and measured at 525 nm (excitation) and 547 nm (emission) spectrofluorometrically.

Protein was determined by the Bradford method (26) with bovine serum as a standard.

Data analysis

The results of the replicates were reported as mean \pm standard error. Comparison of treatments against controls was made using a one-way ANOVA; the significance level chosen for all statistical analysis was $p < 0.05$.

Results and Discussion

Chemical composition of the essential oil

GC/MS analysis of the plant essential oil

led to the identification and quantification of 5 components, which accounted for 99.9% of the total oil (Table 1). The major components of the oil were carvacrol (52.5%) and linalool (45.4%). The essential oil consisted mainly of oxygenated monoterpenes (98.2%), whereas monoterpene hydrocarbons were weakly represented (1.7%). A portion (0.1%) of the total composition was not identified. Terpinen-4-ol was found as a main component in *Origanum majorana* essential oil which was from Kalocsa, Hungary. Linalool was also found as a component (12.1%) but carvacrol was not found (27).

As far as we know, there are many researchers on the investigation of chemical composition of the oils isolated from *Origanum* species (13, 28, 29). It was postulated that the oil exists in two forms. One with terpinen-4-ol and sabinene hydrate as major components (30) and the other with thymol and /or carvacrol (28, 30, 31) as predominant compounds. The marjoram essential oil has shown that the volatile aroma composition varies according to geographic differences and by the climatic features such as temperature (32) thereby altering the biological activities studied (33).

The antioxidant activity

Radical-scavenging activity of the essential oil from aerial parts of *O. majorana* was evaluated by the 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical assay. The concentration that led to 50% inhibition (EC_{50}) is 170 μ g/mL. Essential oil's radical-scavenging activity was found 9.2, 52.96 and 23.45 times lower than BHT, ascorbic

Table 2. Antiradical activities and linoleic acid oxidation inhibitions of *O. majorana* essential oil and positive controls.

Samples	DPPH assay (EC ₅₀ , µg/mL)	β-Carotene/linoleic acid assay (I%)
<i>O. majorana</i>	170 ± 0.3	40.0 ± 0.2
BHT	18.5 ± 0.2	94.8 ± 0.1
Ascorbic acid	3.2 ± 0.4	92.0 ± 0.3
α-Tocopherol	7.2 ± 0.6	96.4 ± 0.5

BHT, butylated hydroxytoluene; DPPH, 2,2-diphenyl-1-picrylhydrazyl; I%, inhibition in percentage.

acid and α-tocopherol respectively (Table 2). 4-terpineol (29.97%), gamma-terpinene (15.40%), trans-sabinene hydrate (10.93%), alpha-terpinene (6.86%), 3-cyclohexene-I-1 methanal, a, a4-trimethyl-,(S)-(CAS) (6.54%), and sabinene (3.91%) were found as main constituents of *Origanum majorana* collected from local market of Cairo, Egypt, and its EC₅₀ concentration was found to be 58.67 mg/mL (34).

Moreover, compared to other *Origanum* species, the DPPH radical scavenging activity of essential oil from *O. majorana* was found higher than *O. vulgare* ssp. *vulgare* which consisted of caryophyllene (14.4%) and spathulenol (11.6%) the main constituents, followed by germacrene-D (8.1%) and α-terpineol (7.5%). but lower than *O. onites* which had linalool (50.53%), carvacrol (24.52%) and thymol (15.66%) as three major constituents (18, 35).

The potential of the essential oil to inhibit lipid peroxidation was evaluated using the β-carotene/linoleic acid bleaching test. The essential oil showed 40% inhibition. The essential oil's linoleic acid oxidation inhibition capacity was found 2.37, 2.3 and 2.41 times lower than BHT, ascorbic acid and α-tocopherol respectively (Table 2). *O. vulgare* ssp. *vulgare* was more effective than *O. majorana* in linoleic acid oxidation (35). Inhibition of linoleic acid oxidation capacity of *O. majorana* was equal to *O. onites* capacity (18).

Linalool's antioxidant activity was studied in the previous research. Linalool's EC₅₀ value was calculated as 16.4 µg/mL in DPPH radical assay and linalool showed antioxidant activity in the β-carotene/linoleic acid bleaching test (36).

The antioxidant effects of the essential oil and linalool on Hep G2 cells

We examined the protective effect (antioxidant) of the essential oil and its oxygenated monoterpene component linalool against strong oxidant H₂O₂ on Hep G2 cells (Figure 1). Preincubation of cells with the essential oil and linalool increased cell viability against H₂O₂ cytotoxicity. These results indicate that the essential oil and linalool are capable of reducing H₂O₂-induced cytotoxicity. Essential oils from wild (camphor, sabinol/sabinyl acetate, thujene and eucalyptol (1,8 cineole) as the major compounds) and cultivated (camphor, eucalyptol (1,8 cineole), sabinol/sabinyl acetate, 3-oxo-beta ionone/isomethyl beta ionone) as the major compounds) form of *Salvia pisidica* reduced the cytotoxicity induced by strong oxidant on H1299 and Hep G2 cells (37). Carvacrol and thymol were the components of *O. majorana*'s essential oil, protected parental, and epirubicin-resistant H1299 cells against H₂O₂-induced cytotoxicity, membrane, and DNA damage when the cells were preincubated with these two compounds at lower concentration (<IC₅₀) before H₂O₂ incubation (38). Aqueous extract from *Morinda officinalis* showed protective effect to H₂O₂-induced cytotoxicity. Viabilities of cells exposed to 100 µmol H₂O₂ decreased below 50% and increased to a statistically significant extent up to 64.0% in the *Morinda officinalis*-treated group at 250 µg/mL (39).

The essential oil and linalool significantly decreased membrane-damaging on H₂O₂ treated Hep G2 cells (Table 3). The membrane-protective effect of linalool was found higher than essential oil. Preincubation with the IC₁₀ and IC₂₀ essential oil concentrations did not

Table 3. Amount of malondialdehyde in Hep G2 cells preincubated with *O. majorana* essential oil and its oxygenated monoterpene component linalool before H₂O₂ treatment.

Groups	MDA levels ^f (nmol/mg protein ± S.E.)
IC ₁₀ E.O + IC ₁₀ H ₂ O ₂	0.56 ± 0.04 ab
IC ₁₀ E.O + IC ₅₀ H ₂ O ₂	0.92 ± 0.07 ab
IC ₁₀ E.O + IC ₇₀ H ₂ O ₂	1.60 ± 0.14 cd
IC ₂₀ E.O + IC ₁₀ H ₂ O ₂	0.42 ± 0.03 a
IC ₂₀ E.O + IC ₅₀ H ₂ O ₂	0.75 ± 0.07 ab
IC ₂₀ E.O + IC ₇₀ H ₂ O ₂	1.52 ± 0.66 cd
IC ₃₀ E.O + IC ₁₀ H ₂ O ₂	0.38 ± 0.06 a
IC ₃₀ E.O + IC ₅₀ H ₂ O ₂	0.68 ± 0.07 ab
IC ₃₀ E.O + IC ₇₀ H ₂ O ₂	1.38 ± 0.03 bc
IC ₁₀ Linalool + IC ₁₀ H ₂ O ₂	0.42 ± 0.08 a
IC ₁₀ Linalool + IC ₅₀ H ₂ O ₂	0.78 ± 0.07 ab
IC ₁₀ Linalool + IC ₇₀ H ₂ O ₂	1.43 ± 0.34 bc
IC ₂₀ Linalool + IC ₁₀ H ₂ O ₂	0.35 ± 0.06 a
IC ₂₀ Linalool + IC ₅₀ H ₂ O ₂	0.72 ± 0.07 ab
IC ₂₀ Linalool + IC ₇₀ H ₂ O ₂	1.30 ± 0.03 bc
IC ₃₀ Linalool + IC ₁₀ H ₂ O ₂	0.29 ± 0.03 a
IC ₃₀ Linalool + IC ₅₀ H ₂ O ₂	0.65 ± 0.07 ab
IC ₃₀ Linalool + IC ₇₀ H ₂ O ₂	1.17 ± 0.55 bc
IC ₁₀ H ₂ O ₂	0.65 ± 0.07 ab
IC ₅₀ H ₂ O ₂	1.1 ± 0.08 b
IC ₇₀ H ₂ O ₂	1.70 ± 0.34 cd
Control	0.23 ± 0.11 a
0.5 % DMSO	0.22 ± 0.20 a

^fMeans (*n*=5) followed by different letters within column are significantly different (*p*≤ 0.05).

MDA; malondialdehyde, E.O; essential oil, S.E.; Standart Error.

change the MDA amounts statistically according to incubation with IC₇₀ H₂O₂. The similar results indicated that the essential oil from *O. onites*, carvacrol and, thymol showed membrane protective (antioxidant) effects at lower than IC₅₀ concentrations on Hep G2 cells (18). In the future *O. majorana* and linalool's protective effects against H₂O₂ will be good sources for hepatocellular carcinoma treatment which need more studies about understanding of the mechanism of the protective effects of essential

oil and linalool against H₂O₂ on Hep G2.

Cytotoxicity of essential oil and linalool on Hep G2 cells

The effects of *O. majorana* essential oil and linalool on Hep G2 cells as assessed by CellTiter-Blue® Cell Viability Assay with different concentrations (5-500 µg/mL) are shown in Figure 2. Dose and time dependent inhibition by the essential oil and linalool were observed with IC₅₀ values of 100, 80 and 63 µg/mL for essential




Figure 1. Protective effect of the essential oil from *O. majorana* and linalool against H₂O₂-induced cytotoxicity on Hep G2 cancer cells. E.O, Essential oil. Values represent mean ± S.E. from three independent experiments.




Figure 2. Cytotoxic activities of *O. majorana* essential oil and linalool for 24, 48 and 72 h on Hep G2 cells. Cell viability was assessed by CellTiter-Blue® Cell Viability Assay.




Figure 3. Dose-dependent membrane-damaging effects of essential oil from *O. majorana* and linalool on Hep G2 cells. E.O., Essential oil.

Values represent mean \pm S.E. from three independent experiments.

* Significantly different from control (untreated cells) and 0.5% (v/v) DMSO control; § Significantly different from IC₅₀ essential oil and linalool treatment in Hep G2 cells cells.

oil and 81.5, 72.7 and 64.7 $\mu\text{g/mL}$ at 24, 48 and 72 h, respectively. The IC₁₀, IC₅₀ and IC₇₀ values of H₂O₂ incubations were 79, 387 and 541 μM respectively. DMSO (0.5%, v/v) did not affect the cell growth when treated for the same time periods. Linalool had more effective cytotoxic activity than essential oil on Hep G2 cells for 24 and 48 h incubations while essential oil had more effective cytotoxic activity than linalool for 72 h incubation. Those results showed that incubation time affected the essential oil's and linalool's cytotoxic effects on Hep G2 cells. *O. onites* essential oil and *O. majorana* essential oil's second and main component respectively. In addition, carvacrol has been reported to be cytotoxic on Hep G2 cells and the essential oil was found to be less toxic than carvacrol and thymol for Hep G2 cells (18). The oils from the rhizome and the aerial part of *A. mollissima* showed cytotoxicity on Hep G2 cells (40). *T. revolutus* Célak essential oil and its two main components (cymene and γ -terpinene) were found cytotoxic in concentration- and time-

dependent manners in Hep G2 cells (41). Plant extracts from *O. majorana* showed cytotoxicity and anti-proliferative effect on Jurkat cells (42). Methanolic extracts of some medicinal plants showed cytotoxic effects on A549, MCF-7, Hep G2 and HT-29 cells (

O. majorana essential oil and linalool induced membrane damage on Hep G2 cells are shown in Figure 3. Membrane-damaging effects of the essential oil and linalool increased with accelerating concentrations. Linalool membrane-damaging effect was found stronger than essential oil membrane-damaging effect. In one of our previous study linalool had more effective membrane damaging effect than *O. majorana* essential oil on epirubicin-resistant H1299 cells but on parental cells, the essential oil had more effective membrane damaging effect than linalool (43). Biochemical changes like membrane structure in drug resistant cells will affect the essential oil's and linalool's cytotoxicity. Differences between the essential oil's and linalool's chemical structures (as a

result of they will produce different ROS and amounts) will be the reason for having different cytotoxic effects on Hep G2 cells of the essential oil and linalool.

Synergistic functions of the various molecules contained in an essential oil, in comparison to the action of one or two main components of the oil, seems questionable. However, it is possible that the activity of the main components is modulated by other minor molecules (44-46). Moreover, it is likely that several components of the essential oil play a role in defining the fragrance, density, texture, colour and above all, cell penetration (47), lipophilic or hydrophilic attraction as well as fixation on cell walls membranes, and cellular distribution. The last feature is very important because the distribution of the oil in the cell determines the different types of radical reactions produced, depending on their compartmentation in the cell.

O. majorana essential oil and its oxygenated monoterpene component linalool showed antioxidant and cytotoxic activity depending on concentrations and time manner. Those results indicate that concentrations are important in their usage. *O. majorana*'s and linalool's antitumoral properties suggest that they could be the potential source of hepatocellular carcinoma treatment.

Acknowledgments

The authors wish to thanks Akdeniz University Scientific Research Projects Unit (2008.01.0105.011) for financial support of this work.

References

- (1) Ozturk-Urek R, Bozkaya LA and Tarhan L. The effects of some antioxidant vitamins and trace elements supplemented diets on activities of SOD, CAT, GSH-Px and LPO levels in chicken tissues. *Cell Biochem. Funct.* (2001) 19: 125-132.
- (2) Ito N, Fukushima S, Hassegawa A, Shibata M and Ogiso T. Carcinogenicity of butylated hydroxyanisole in F344 rats. *J. Natl. Cancer Institute* (1983) 41: 215-217.
- (3) Moeller SM, Jacques P and FBlumberg JB. The potential role of dietary xanthophylls in cataract and age-related macular degeneration. *J. Am. Coll. Nutr.* (2000) 19: 522-527.
- (4) Miliauskas G, Venskutonis PR and Van Beek TA. Screening of radical scavenging activity of some medicinal and aromatic plant extracts. *Food Chem.* (2004) 85: 231-237.
- (5) Ozkan A and Erdogan A. Antioxidant and anticancer activity of fresh corm extract from *Romulea tempskyana* (Iridaceae). *Nat. Prod. Res.* (2012) 26: 2126-2128.
- (6) Werker E, Putievsky E, Ravid U, Dudai N and Katzir I. Glandular hairs and essential oil in developing leaves of *Ocimum basilicum* L. (Lamiaceae). *Ann. Bot. London* (1993) 71: 43-50.
- (7) Akgul A and Kivanc M. Inhibitory effects of selected Turkish species and Oregano components on some foodborne fungi. *Int. J. Food Microbiol.* (1988) 6: 263-268.
- (8) Dapkevicius A, Van Beek TA, Lelyveld GP, Van Veldhuizen A, Groot A, deLinsse JPH and Venskutonis R. Isolation and structure elucidation of radical scavengers from *Thymus vulgaris* leaves. *J. Nat. Prod.* (2002) 65: 892-896.
- (9) Cipak L, Grausova L, Miadokova E, Novotny L and Rauko P. Dual activity of triterpenoids: apoptotic versus antidifferentiation effects. *Arch. Toxicol.* (2006) 80: 429-435.
- (10) Tapiero H, Tew KD, Nguyen Ba G and Mathe G. Polyphenols: do they play a role in the prevention of human pathologies? *Biomed. Pharmacother.* (2002) 56: 200-207.
- (11) Niture SK, Rao US and Srivenugopal KS. Chemopreventative strategies targeting the MGMT repair protein: augmented expression in human lymphocytes and tumor cells by ethanolic and aqueous extracts of several Indian medicinal plants. *Int. J. Oncol.* (2006) 29: 1269-1278.
- (12) Lin LT, Liu LT, Chiang LC and Lin CC. *In-vitro* anti-hepatoma activity of fifteen natural medicines from Canada. *Phytother. Res.* (2002) 16: 440-444.
- (13) Vera RR and Chane-Ming J. Chemical composition of the essential oil of marjoram (*Origanum majorana* L.) from Reunion Island. *Food Chem.* (1999) 66: 143-145.
- (14) Deans SG and Svoboda KP. The antimicrobial properties of marjoram (*Origanum majorana* L.) volatile oil. *Flav. Frag. J.* (1990) 5: 187-190.
- (15) Bickers D, Calow P, Greim H, Hanifin JM, Rogers AE, Saurat JH, Sipes IG, Smith RL, and Tagami H. A toxicological and dermatological assessment of linalool and related esters when used as fragrance ingredients. *Food Chem. Toxicol.* (2003) 41: 919-942.
- (16) Knasmuller S, Parzefall W, Sanyal R, Ecker S, Schwab C, Uhl M, Mersch-Sundermann V, Williamson G, Hietsch G, Langer T, Darroudi F and Natarajan AT. Use of metabolically competent human hepatoma cells for the detection of mutagens and antimutagens. *Mutat. Res.* (1998) 402: 185-202.
- (17) Ozkan A and Fışkın K. Epirubicin HCl toxicity in human-liver derived hepatoma G2 cells. *Pol. J. Pharmacol.* (2004) 56: 435-444.
- (18) Ozkan A and Erdogan A. A Comparative evaluation

- of antioxidant and anticancer activity of essential oil from *Origanum Onites* (Lamiaceae) and its two major phenolic components. *Türk. J. Biol.* (2011) 35: 735-742.
- (19) Baydar H and Baydar Göktürk N. The effects of harvest date, fermentation duration and Tween 20 treatment on essential oil content and composition of industrial oil rose (*Rosa damascena* Mill.). *Ind. Crop Prod.* (2005) 21: 251-255.
- (20) Eikani MH, Golmohammad F, Rowshanzamir S and Mirza M. Recovery of water-soluble constituents of rose oil using simultaneous distillation-extraction. *Flav. Frag. J.* (2005) 20: 555-58.
- (21) Cuendet M, Hostettmann K and Potterat O. Iridoid glucosides with free radical scavenging properties from *Fagraea blumei*. *Helv. Chim. Acta* (1997) 80: 1144-1152.
- (22) Burits M and Bucar F. Antioxidant activity of *Nigella sativa* essential oil. *Phytother. Res.* (2000) 14: 323-328.
- (23) Dapkevicius A, Venskutonis R, Van Beek TA, Linszen PH. Antioxidant activity of extracts obtained by different isolation procedures from some aromatic herbs grown in Lithuania. *J. Sci. Food Agr.* (1998) 77: 140-146.
- (24) Gloeckner H, Jonuleit T and Lemke HD. Monitoring of cell viability and cell growth in a hollow-fiber bioreactor by use of the dye Alamar Blue (TM). *J. Immunol. Methods* (2001) 252: 131-138.
- (25) Wasowicz W, Neve J and Peretz A. Optimized steps in fluorometric determination of thiobarbituric acid-reactive substances in serum; importance of extraction pH and influence of sample preservation and storage. *Clin. Chem.* (1993) 39: 2522-2526.
- (26) Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein dye binding. *Anal. Biochem.* (1976) 72: 248-254.
- (27) Va'gi E, Sima'ndi B, Suhajda A' and He'thelyi E'. Essential oil composition and antimicrobial activity of *Origanum majorana* L. extracts obtained with ethyl alcohol and supercritical carbon dioxide. *J. Agric. Food Chem.* (2005) 38: 51-57.
- (28) Baser KHC, Kirimer N and Tümen G. Composition of the essential oil of *Origanum majorana* L. from Turkey. *J. Essent. Oil Res.* (1993) 5: 577-579.
- (29) Busatta C, Vidal RS, Popiolski AS, Mossi AJ, Dariva C, Rodrigues MRA, Corazza FC, Corazza ML, Vladimir Oliveira J and Cansian RL. Application of *Origanum majorana* L. essential oil as an antimicrobial agent in sausage. *Food Microbiol.* (2008) 25: 207-211.
- (30) Sarer E, Scheffer JJC and Svedsen AB. Monoterpenes in the essential oil of *Origanum majorana*. *J. Med. Plants Res. Planta Med.* (1982) 46: 236-239.
- (31) Jolivet J, Rey P and Boussarie MF. Differentiation of some essential oils with similar structure (especially by examining their ultraviolet, infrared, and Raman spectrum). II. Essence of *Origanum majorana* L., and essence of *Origanum vulgare* L. *Plantes Med. Phytother.* (1971) 5: 199-208.
- (32) Vági E, Rapavi E, Hadolin M, Vásárhelyiné Perédi K, Balázs A, Blázovics A and Simándi B. Phenolic and triterpenoid antioxidants from *Origanum majorana* L. herb and extracts obtained with different solvents. *J. Agric. Food Chem.* (2005) 53: 17-21.
- (33) Vardar-Ünlü G, Candan F, Sokmen A, Daferera D, Polissiou M, Sokmen M, Donmez E and Tepe B. Antibacterial and antioxidant activity of the essential oil and methanol extracts of *Thymus pectinatus* Fisch. et Mey var. *pectinatus* (Lamiaceae). *J. Agric. Food Chem.* (2003) 51: 63-67.
- (34) Mossa ATH and Nawwar GAM. Free radical scavenging and antiacetylcholinesterase activities of *Origanum majorana* L. essential oil. *Hum. Exp. Toxicol.* (2011) 30: 1501-1513.
- (35) Şahin F, Güllüce M, Daferera D, Sökmen A, Sökmen M, Polissiou M, Agar G and Özer H. Biological activities of the essential oils and methanol extract of *Origanum vulgare* ssp. *vulgare* in the Eastern Anatolia region of Turkey. *Food Control* (2004) 15: 549-557.
- (36) Hussain AI, Anwar F, Sherazi STH and Przybylski R. Chemical composition, antioxidant and antimicrobial activities of basil (*Ocimum basilicum*) essential oils depends on seasonal variations. *Food Chem.* (2008) 108: 986-995.
- (37) Ozkan A, Erdogan A, Sokmen M, Tugrulay S and Unal O. Antitumoral and antioxidant effect of essential oils and *in-vitro* antioxidant properties of essential oils and aqueous extracts from *Salvia pisidica*. *Biologia* (2010) 65: 990-996.
- (38) Ozkan A and Erdogan A. A Comparative study of the antioxidant/prooxidant effects of carvacrol and thymol at various concentrations on membrane and DNA of parental and drug resistant H1299 cells. *Nat. Prod. Commun.* (2012) 7: 1557-1560.
- (39) Chang MS, Kim WN, Yang WM, Kim HY, Oh JH and Park SK. Cytoprotective effects of *Morinda officinalis* against hydrogen peroxide-induced oxidative stress in Leydig TM3 cells. *Asian J. Androl.* (2008) 10: 667-674.
- (40) Yu JQ, Liao ZX, Cai XQ, Lei JC and Zou GL. Composition, antimicrobial activity and cytotoxicity of essential oils from *Aristolochia mollissima*. *Environ. Toxicol. Phar.* (2007) 23: 162-167.
- (41) Erdogan A and Ozkan A. Effects of *Thymus revolutus* Célak essential oil and its two major components on Hep G2 cells membrane. *Biologia* (2013) 68: 105-111.
- (42) Abdel-Massih RM, Fares R, Bazzi S, El Chami N and Baydoun E. The apoptotic and anti-proliferative activity of *Origanum majorana* extracts on human leukemic cell line. *Leukemia Res.* (2010) 34: 1052-1056.
- (43) Erdogan A and Ozkan A. A comparative study of cytotoxic, membrane and DNA damaging effects of *Origanum majorana*'s essential oil and its oxygenated monoterpene component linalool on parental and epirubicin-resistant H1299 cells. *Biologia* (2013) 68: 754-761.
- (44) Franzios G, Mirosou M, Hatzia Apostolou E, Kral J,

- Scouras ZG and Mavragani Tshipidou P. Insecticidal and genotoxic activities of mint essential oils. *J. Agric. Food Chem.* (1997) 45: 2690-2694.
- (45) Santana-Rios G, Orner GA, Amantana A, Provost C, Wu SY and Dashwood RH. Potent antimutagenic activity of white tea in comparison with green tea in the *Salmonella* assay. *Mutat. Res.* (2001) 495: 61-74.
- (46) Hoet S, Ste'vigny C, He'rent MF and Quetin-Leclercq J. Antitrypanosomal compounds from leaf essential oil of *Strychnos spinosa*. *Planta Med.* (2006) 72: 480-482.
- (47) Cal K. Skin penetration of terpenes from essential oils and topical vehicles. *Planta Med.* (2006) 72: 311-316.
- (48) Behzad S, Pirani A and Mosaddeg M. Cytotoxic activity of some medicinal plants from Hamedan District of Iran. *Iran. J. Pharm. Res.* (2014) 13: 199-205.

This article is available online at <http://www.ijpr.ir>

Archive of SID