

Preparation of a Major Metabolite of Igaratimod and Simultaneous Assay of Igaratimod and Its Metabolite by HPLC in Rat Plasma

Jun-ping Han^{a, b}, Zhen-han Zhu^a, Yue-zhang Wu^a, Wen Qian^c, Zhi-yu Li^d, Miyu Nishikawa^e, Toshiyuki Sakaki^e and Chang-qing Yang^{a, c*}

^aDepartment of Clinical Pharmacy, School of Basic Medicine and Clinical Pharmacy, China Pharmaceutical University, No. 639 Longmian Avenue, Jiangning District, Jiangsu Province, PR China. ^bDepartment of Pharmacy, The Second Affiliated Hospital of Soochow University, Suzhou City, Jiangsu Province, 215004, PR China. ^cNanjing BRT-Biomed Co., Ltd. Jiangning District, Jiangsu Province, PR China. ^dDepartment of Pharmaceutical Chemistry, School of Pharmacy, China Pharmaceutical University, No. 639 Longmian Avenue, Jiangning District, Jiangsu Province, PR China. ^eDepartment of Biotechnology, Faculty of Engineering, Toyama Prefectural University, 5180 Kurokawa, Imizu, Toyama 939-0398, Japan.

Abstract

Igaratimod is a new synthetic disease-modifying antirheumatic drug intended to treat patients with rheumatoid arthritis. A new method using recombinant human CYP450s yeast cells containing c-DNA expressed P450s was applied to identify the metabolic pathways of iguratimod and to prepare its metabolite. The metabolite was isolated, and its structure was identified by quadrupole time-of-flight-mass spectrometry and nuclear magnetic resonance. Furthermore, a selective and sensitive high performance liquid chromatography (HPLC) method was developed for the simultaneous quantification of iguratimod and its major metabolite in rat plasma for the first time. The results indicated that iguratimod was mainly metabolized to a metabolite by CYP2C9 and CYP2C19 in *in-vitro* study. The structure of the metabolite was identified as M2 (N-[3-(acetamido)-4-oxo-6-phenoxy-4H-chromen-7-yl]methanesulfonamide). HPLC assay was achieved on a C18 column using methanol-water containing 0.1% trifluoroacetic acid (55:45 v/v) at a flow rate of 1 mL/min with UV detection at 257 nm. Standard calibration curves were obtained in the concentration range of 0.5–20 µg/mL for iguratimod and its metabolite M2. The lower limits of detection of iguratimod and M2 in rat plasma were 0.1 and 0.25 µg/mL, respectively. The intra- and inter-day precision (RSD%) were within 5% for the two analytes. The average recoveries of the analytes were greater than 90%. In conclusion, recombinant human CYP450s whole-yeast transformation system could be successfully used to identify and prepare the major metabolite of iguratimod. The HPLC method we developed could be successfully applied to evaluate pharmacokinetics of iguratimod and its metabolite M2 in rats.

Keywords: Igaratimod; Recombinant human CYP450s; Metabolic pathway; Pharmacokinetics; HPLC assay.

Introduction

Igaratimod, a novel disease-modifying antirheumatic drug, which is now used in hospitals in China and Japan, has been confirmed as a highly efficacious and safe drug for

* Corresponding author:

E-mail: ycq0315@yahoo.com




Figure 1. Metabolites and possible pathways of igratimod.

rheumatoid arthritis therapy (1). Previous studies on igratimod mainly focused on its clinical efficacy, mechanisms and adverse reactions since it was firstly used to treat rheumatoid arthritis. However, there were few researches on the pharmacokinetics of igratimod.

Igratimod is extensively metabolized by the liver, and several metabolites of igratimod have been characterized in plasma from healthy male subjects following oral administration of the drug (Figure 1) (2). Previous researches reported that the major metabolites of igratimod are a deformylated form (M1) and its N-acetylated form (M2). The other metabolites are phydroxylated forms in the 6-phenoxy groups of igratimod, M1, and M2, which have been termed M4, M5, and M3, respectively (2). However, none of the studies confirmed the specific metabolic pathways of igratimod. A high-performance liquid chromatographic (HPLC) method with UV detector had been developed for the determination of igratimod (T-614) in rat plasma (3). The *in-vivo* pharmacokinetics of igratimod was consistent with the dynamics of the one-compartment model. The results of a preclinical pharmacokinetic study indicated that T-614 was absorbed rapidly and eliminated slowly in animals (4). However, none of studies

had developed a method of simultaneous determination of igratimod and its metabolite in rat plasma.

Elucidation of phase I metabolic pathways and preparation of the metabolites of new drugs are very important in the drug discovery and the clinical use of new drugs. Enzyme inhibition or induction is known to alter the activity of CYP isozymes and potentially increase the risk of side effects or toxicity of drugs. Metabolite standards of the new drug are of particular importance in the context of metabolites in safety testing (5). *In-vitro* models of drug metabolism include subcellular cell models (such as microsomes), cell line models (such as primary hepatocytes, transgenic cell lines, stem cell-induced differentiation of liver cells), recombinant metabolic enzyme models, tissue sections, *in-vitro* animal model (such as liver/intestinal perfusion model) and so on (6). Drăgan CA *et al.* successfully prepared 4'-hydroxydiclofenac at gram-scale from diclofenac using a fission yeast strain functionally co-expressing human

CPR and CYP2C9. They demonstrated that recombinant human CYP450s whole-cell yeast transformation system could be used in the confirmation of metabolic pathways and preparation of drug metabolite standard (7).

In this study, we investigated the metabolic pathways of iguratimod and prepared a main metabolite of iguratimod using a recombinant human CYP450s yeast whole-cell transformation system. The research method will provide a reference for studies on metabolic pathways and preparation of metabolites of new drugs or candidate drugs. In addition, an HPLC method for the simultaneous determination of iguratimod and its metabolite M2 in rat plasma was developed for the first time. This method will provide a basis for clinical pharmacokinetic studies of iguratimod and its metabolite M2.

Experimental

Materials

Igratimod (T-614, purity 99.2%, HPLC grade) was supplied by Jiade Medical Technology Co., Ltd. (Changzhou, China). Igratimod metabolite (M2) was isolated and purified in our laboratory (purity >98%, HPLC grade). Igratimod tablets were purchased from Simcere (Hai'kou, Hainan, China). Phenacetin (purity $\geq 98\%$), used as the internal standard (IS), was purchased from Macklin Biochemical Co., Ltd. (Shanghai, China). Recombinant human CYP450s yeast cells containing c-DNA expressed P450s (CYP1A2, CYP2C9, CYP2C19, CYP2D6, CYP2E1, and CYP3A4) were obtained from Nanjing BRT-Biomed Co., Ltd. (Nanjing, China). Other chemicals were obtained from the following sources: D-(+)-Glucose from Aladdin Industrial Corporation (Shanghai, China); di-Potassium hydrogen phosphate trihydrate and Potassium dihydrogen phosphate from Xilong Chemical Co. Ltd. (Shantou, Guangdong, China); trifluoroacetic acid (TFA) from Nanjing Chemical Reagent Co., Ltd. (Nanjing, China); Demethyl sulfoxide from Sinopharm Chemical Reagent (Nanjing, China); Ethyl acetate from Nanjing Chemical Reagent Co., Ltd. (Shanghai, China). Acetonitrile (HPLC grade) and methanol (HPLC grade) were purchased from Tedia (Fairfield, OH, USA).

Distilled water, prepared from demineralized water, was used throughout the study.

Chromatographic condition

The chromatographic system was performed by LC-2010A HT high performance liquid chromatograph (Shimadzu, Japan). The assay was performed on ACE-C18 column ($250 \times 4.6 \text{ mm} \times 5 \mu\text{m}$) (ACE, UK) and a InertSustain C18 precolumn ($4.0 \times 10 \text{ mm} \times 5 \mu\text{m}$) (Shimadzu, Japan) was fitted just before the inlet junction of the analytical column. The mobile phase consisted of methanol-water containing 0.1% TFA (55:45 v/v) at a flow rate of 1 mL/min with UV detection at 257 nm. The column temperature was maintained at 30 °C, and the injection volume was 10 μL .

Incubation of iguratimod with the recombinant human CYP450s yeast cells

The incubation mixture (4 mL final volume) of recombinant CYP450s yeast whole-cell transformation system includes 1 g recombinant CYP450s yeasts containing c-DNA expressed P450s (CYP1A2, CYP2C8, CYP2C9, CYP2C19, CYP2D6, CYP2E1 or CYP3A4), 2 mL 0.2 M K_2HPO_4 phosphate buffer (pH 7.4), 0.4 mL 20% D-(+)-Glucose solution, 1.4 mL sterile water and 0.2 mL 10 mM iguratimod dissolving in DMSO to make the final system consisting of 2% glucose solution and 0.5 mM iguratimod. Seven types of recombinant human CYP450s yeast whole-cell reaction system at seven individual 50 mL tubes were initiated at the parallel time. The mixture in each tube was incubated for 72 h at 30 °C. The samples were collected from the incubation system at 0, 24, 48, and 72 h by taking 100 μL reaction solution to 1.5 mL clean and dry tubes and adding 200 μL methanol. The samples were fully extracted by vortex-mixing for 2 min. After centrifugation at $12,000 \times g$ for 10 min, the supernatant was collected and an aliquot (10 μL) was injected into the chromatographic system for analysis.

Identification of the iguratimod metabolite in rat plasma sample

Male Sprague-Dawley rats (180-200 g) were obtained from Jiesijie experimental animal Co., Ltd. (Shanghai, China) (license key, SCXK

2013-0006) and housed with a 12 h light/12 h night cycle at ambient temperature (about 25 °C) and 60% relative humidity. Free access to food and water was allowed at all times except for fasting 12 h before the experiment but with free access to water. All animal experiments were carried out according to the Guidelines for the Care and Use of Laboratory Animals, and were approved by the Animal Ethics Committee.

To a 50 µL aliquot of rat plasma, 25 µL Phenacetin IS solution (20 µg/mL) and 25 µL acetonitrile were added. After vortex-mixing for 2 min and centrifugation (12,000×g) for 10 min, the supernatant was collected and an aliquot (10 µL) was injected into the chromatographic system for analysis.

Preparation and isolation of the iguratimod metabolite

To prepare the metabolite of iguratimod, recombinant human CYP2C9 yeast cell reaction system was used according to the method described above. Total volume of the system is 80 mL which contains 20 g recombinant CYP450s yeasts, 40 mL 0.2 M K₂HPO₄ phosphate buffer (pH 7.4), 8 mL 20% D-(+)-Glucose solution, 28 mL sterile water and 4 mL 10 mM iguratimod dissolving in DMSO. After incubation at 30 °C for 72 h, 160 mL acetic ether were added to stop the reaction. Then the acetic ether extraction solution was concentrated by a rotary evaporator and then redissolved using 10 mL methanol. Preparative HPLC was performed in an isocratic elution mode using Durashell C18 (250 mm × 10 mm, 10 µm). The isocratic mobile phase of water-methanol (52:48, v/v) containing 0.1% TFA (v/v) was run at a flow rate of 7 mL/min. Other parameters were as follows: detection wavelength, 257 nm; injection volume, 200 µL; column and ambient temperature, 25 °C. The eluents were freeze-dried by a Bilon freeze dryer to prepare the metabolite powder.

Identification of the iguratimod metabolite structure by QTOF-MS and Nuclear magnetic resonance spectroscopy

HR-MS spectral data was obtained on Agilent technologies 6520 Accurate-Mass QTOF-MS instruments (Agilent, USA). Data acquisition was performed with Agilent Mass hunter

Workstation with advanced data acquisition and data analysis capabilities. The mass spectrometer was operated in the ESI-positive mode, and the MS data were collected at the range of 320 to 490. The HNMR spectra were recorded on Bruker AV-300 (300 MHz) apparatus (Bruker, USA).

Method validation of HPLC

To assess selectivity of the method, blank plasma samples obtained from seven rats were processed by the same extraction procedure and chromatographed to determine whether endogenous components would interfere with the determination of iguratimod, iguratimod metabolite, and internal standard (Phenacetin).

The Calibration curves were prepared by assaying standard plasma samples at six concentration levels of iguratimod and metabolite over the range of 0.5-20 µg/mL, respectively. The calibration curves were constructed by plotting the peak area ratio of analytes to IS, and concentrations being used as y and x variables in a standard regression analysis. The lower limit of detection was defined as the detected concentration when the S/N ratio is 3.

Precision and accuracy were carried out in three replicates at 0.5, 10, and 20 µg/mL on the same day and on three consecutive validation days. The accuracy was expressed as relative error (RE%). The precision was calculated by relative standard deviation (RSD%). Intra- and inter-day precisions calculated as RSD (%) were required to be below 15% and accuracy as RE (%) to be within ±15%. Recoveries of the iguratimod and iguratimod metabolite were determined by comparing post-extraction standard plasma concentrations and control standard samples at corresponding concentrations.

Application to pharmacokinetic studies

The studies were approved by the Animal Ethics Committee of China Pharmaceutical University. The animals were adapted to the facilities for one week, and then fasted with free access to water overnight prior to the experiment. The iguratimod tablets were suspended in 0.1% CMC-Na aqueous solution and administered to the rats (10mg/kg bodyweight) by oral gavage. Approximately, 300 µL blood samples were




Figure 2. The representative HPLC chromatograph obtained from the analysis of reaction solutions in the recombinant human CYP2C9 and CYP2C19 reaction system after 72 h incubation. (A) CYP2C9 incubation – 0 h; (B) CYP2C9 incubation – 72 h; (C) CYP2C19 incubation – 0 h; (D) CYP2C19 incubation – 72 h (1. Igratimod; 2. igratimod metabolite).

collected from fundus venous plexus before dosing and at 0.25, 0.5, 1, 2, 3, 4, 6, 8, 10, 12 and 24 h following oral gavage. The blood samples were transferred to heparinized Eppendorf tubes and centrifuged at $4000\times g$ for 10 min to separate the plasma. The plasma samples were immediately frozen at $-80\text{ }^{\circ}\text{C}$ until analysis. The pharmacokinetic parameters of igratimod and its metabolite were calculated using Phoenix WinNonlin[®] 6.1 version. All results were represented as mean \pm SD.

Results and Discussion

Identification of the metabolic pathway of igratimod

In *in-vitro* study, igratimod was incubated with seven types of recombinant human

CYP450s (CYP1A2, CYP2C8, CYP2C9, CYP2C19, CYP2D6, CYP2E1, and CYP3A4) yeasts, respectively. After incubation at $30\text{ }^{\circ}\text{C}$ for 72 h, the reaction solution was extracted with methanol and analyzed by HPLC. We found that there was a new chromatographic peak in the recombinant human CYP2C9 and CYP2C19 transformation system with the retention time at approximately 9.064 min (Figure 2). Meanwhile, the reaction rate of igratimod in recombinant human CYP2C9 transformation system was higher than that in recombinant human CYP2C19 transformation system, with the value 50.73% and 15.83%, respectively. There were no new chromatographic peaks and the reaction rate was near 0% for the other five CYP450s transformation system. The reaction rate was calculated using the following equation:

$$\text{Reaction rate (\%)} = \frac{\text{iguratimod concentration at 0 h} - \text{iguratimod concentration at 72 h}}{\text{iguratimod concentration at 0h}} \times 100\%$$

In *in-vivo* study, we found that there was a same chromatographic peak in rat plasma after taking iguratimod, which was at the same retention time of approximately 9.064 min with that *in-vitro* study (Figure 3).

There was a new chromatographic peak after 72 h incubation when assaying the recombinant human CYP2C9 and CYP2C19 reaction solutions using HPLC. In contrast, it was not observed in the recombinant human CYP1A2, CYP2C8, CYP2D6, CYP2E1, and CYP3A4 incubation system under the present conditions, as determined by HPLC with UV detection at 257 nm. Meanwhile, we found a same chromatographic peak at the same retention time in the rat plasma with that *in-vitro* study. Therefore, we expediently assumed that iguratimod could be metabolized by CYP2C9 and CYP2C19.

In the present study, we examined the metabolic pathways of iguratimod in *in-vitro* study using the recombinant human CYP450s (CYP1A2, CYP2C8, CYP2C9, CYP2C19, CYP2D6, CYP2E1, and CYP3A4) yeast whole-cell transformation system. It is the first time to discover the metabolic pathways of iguratimod using the transformation system in this study. Among *in-vitro* recombinant human metabolic enzyme expression system, yeasts have a

more complete cell subcellular (endoplasmic reticulum) structure comparing to prokaryotic expression system. Meanwhile, yeast expression system has a relatively high expression rate and catalytic activity of recombinant enzyme, short production cycle, and it is suitable for large quantities of the preparation comparing to mammalian cell expression system (8). Yasuda *et al.* transfected human CYP1A1, 1A2, 2A6, 2B6, 2C8, 2C9, 2C18, 2C19, 2D6, 2E1 and 3A4 genes into yeast cells to prepare recombinant human CYP450s and investigated the metabolic characteristic of sesamin. They inferred that sesamin was metabolized by CYP2C9, 1A2, 2C19, and 2D6, mainly by CYP2C9 (9). Therefore, *in-vitro* recombinant human CYP450s yeast system is a novel and effective way for studying the metabolic characteristics of new drugs.

Isolation and identification of the iguratimod metabolite by QTOF-MS and HNMR

Prior to structure identification, the metabolite was prepared from recombinant human CYP2C9 yeast transformation system using preparative HPLC method. The final reaction rate of iguratimod in preparation system was 43.5%. The yield of iguratimod metabolite was 15.4% after extraction and purification. The final




Figure 3. Typical HPLC chromatograph for iguratimod and its metabolite in rat plasma. (A) Rat blank plasma; (B) Rat plasma sample at 6 h after oral administration of iguratimod + Phenacetin (1. Igaratimod; 2. iguratimod metabolite; 3. IS, Phenacetin).




Figure 4. Product ion mass spectra of igratimod metabolite.

purity of the igratimod metabolite we prepared reached at 98.8%. The chemical structure was identified by the combination of QTOF-MS and HNMR. Figure 4 shows the QTOF-MS spectra of igratimod metabolite, which formed protonated molecules [M+H] as major ion peaks. HRMS (ESI): m/z, calculated for C₁₈H₁₆N₂O₆S, found 388.0729 (M + H)⁺. Figure 5 shows the HNMR spectra of igratimod metabolite. ¹H NMR (300 MHz, DMSO-*d*₆ + D₂O): δ 9.13 (s, 1H), 7.62 (s, 1H), 7.45 (t, *J* = 7.6 Hz, 2H), 7.30 (s, 1H), 7.25 – 7.18 (m, 1H), 7.10 (d, *J* = 7.8 Hz, 2H), 3.11 (s, 3H), 2.10 (s, 3H). On this basis, the assignment of the igratimod metabolite is M2 (N-[3-(acetamido)-4-oxo-6-phenoxy-4H-chromen-7-yl]methanesulfonamide) as shown in Figure 1.

The current reaction system that we prepared a metabolite standard of igratimod, included

recombinant CYP450s yeasts containing c-DNA expressed CYP2C9, 0.2 M K₂HPO₄ phosphate buffer (pH 7.4), D-(+)-Glucose solution, sterile water and igratimod dissolving in DMSO. A previous literature had thoroughly investigated the influence of the factors such as pH, cell density and incubation time on the incubation for the whole-cell biotransformation of 4'-methyl- α -pyrrolidiobutyrophenone (MPBP) and dexamethasone (DXM) in CAD58 cultures (10). In our study, we found pH had no apparent influence on igratimod transformation rate. The temperature of yeast cell culture is usually set at 30 °C; however, the optimum temperature for human liver drug enzyme is 37 °C. Therefore, we compared the reaction rate of igratimod between 30 °C and 37 °C in the pre-study. It was found that the reaction rate of igratimod was higher at 30 °C than at 37 °C. Metabolite




Figure 5. HNMR spectra of igratimod metabolite.

formation was linear in the initial phase and leveled off in later incubation phases, reaching almost zero after approximately 72 h.

The structure of the metabolite prepared from recombinant CYP2C9 yeast incubation system was identified by QTOF-MS and HNMR. HRMS data show that m/z of igratimod metabolite is 388.0729, calculated for $C_{18}H_{16}N_2O_6S$. On the basis of HRMS and HNMR data, we confirmed the structure of igratimod metabolite as M2 (N-[3-(acetamido)-4-oxo-6-phenoxy-4H-chromen-7-yl]methanesulfonamide) (Figure 1). We inferred that igratimod may be metabolized by CYP2C9 and CYP2C19 to M2.

Method validation of HPLC

Typical chromatograms obtained from a blank, a spiked plasma sample with the analytes (at 10 $\mu\text{g/mL}$) and IS, and a plasma sample after an oral administration of igratimod are shown in Figure 6. No obvious interferences from endogenous plasma substances were observed under the chromatographic conditions.

To evaluate linearity, calibration of

igratimod and igratimod metabolite M2 (0.5–20 $\mu\text{g/mL}$) were prepared and assayed on three consecutive days. The peak area ratios of igratimod or igratimod metabolite to IS in rat plasma varied linearly over the concentration ranges. The regression equations for the drug and its metabolite were $y = 0.2311x + 0.0321$ ($r = 0.9999$) and $y = 0.1337x - 0.0022$ ($r = 0.9999$), respectively, where y refers to the peak area ratios of analytes to IS and x is the concentration of igratimod or igratimod metabolite. The present HPLC method gave the lower limit of detection for igratimod and its metabolite at 0.1 $\mu\text{g/mL}$ and 0.25 $\mu\text{g/mL}$, respectively.

Both intra- and inter-day precisions were <5.0% at three QC levels (0.5, 10.0 and 20.0 $\mu\text{g/mL}$), and RE values for igratimod and igratimod metabolite were within $\pm 10\%$. The mean extraction recoveries of igratimod at concentrations of the QC levels were 98.25, 104.12, and 97.25%, respectively. The mean extraction recoveries of igratimod metabolite M2 at concentrations of the QC levels were 95.79, 97.21, and 96.67%, respectively. These




Figure 6. High Performance Liquid Chromatography of igratimod, a igratimod metabolite M2 and Phenacetin in Rat Plasma. (A) Blank plasma; (B) Standard plasma samples; (C) Plasma samples at 6 h after oral administration of igratimod (1. Igratimod; 2. igratimod metabolite; 3. IS, Phenacetin).

results indicated that the present method has an acceptable precision and accuracy (Table 1).

Previous studies that conducted on this drug is just to determine the HPLC method of igratimod in animal plasma and in the healthy adults plasma (4, 11). However, the determination of igratimod metabolite data was not available in the previous study because igratimod metabolite had not been studied. Therefore, a new HPLC method of simultaneous determination of igratimod and its metabolite in rat plasma was developed for the first time in this study. No obvious interferences from endogenous plasma substances were observed under the chromatographic conditions (Figure 3). There was excellent correlation between the ratio of peak area and concentration for igratimod and its metabolite within the test ranges. The assay precision, accuracy and

extraction recoveries for igratimod and its metabolite were within the acceptance (Table 1). This method was successfully applied to the pharmacokinetic studies of igratimod and its metabolite after oral administration of igratimod in rats (Figure 7 and Table 2).

Application to pharmacokinetic studies

The present HPLC method was successfully applied to determining the plasma concentration of igratimod and igratimod metabolite in Sprague–Dawley rats. After a single oral gavage administration of 10 mg/kg igratimod to rats, the concentration–time profile was constructed for up to 24 h. Figure 7 shows the mean concentration–time profile of igratimod and igratimod metabolite in the rat plasma. The pharmacokinetic parameters for the drug and its metabolite M2 are shown in Table 2.

Table 1. Precision, accuracy and extraction recoveries of analytes (n = 3).

Analytes	Spiked (ug/mL)	intra-day		inter-day		Extraction Recovery	
		Precision (RSD%)	Accuracy (RE%)	Precision (RSD%)	Accuracy (RE%)	Mean (%)	Precision (RSD%)
iguratimod	0.5	2.29	-1.75	2.36	-6.79	98.25 ± 0.01	2.29
	10	0.48	4.12	0.21	3.67	104.12 ± 0.50	0.48
	20	0.15	-2.75	1.16	-1.58	97.25 ± 0.03	0.15
iguratimod metabolite	0.5	1.56	-4.21	1.90	-4.95	95.79 ± 0.01	1.56
	10	0.52	-2.79	0.29	-2.69	97.21 ± 0.50	0.52
	20	0.56	-3.33	1.19	-2.55	96.67 ± 0.11	0.56




Figure 7. Mean plasma concentration-time profiles of iguratimod and iguratimod metabolite in rat plasma after oral administration of 10 mg/kg iguratimod. (mean ± SD, n = 7). (●iguratimod; ■iguratimod metabolite M2).

Table 2. Pharmacokinetic parameters of iguratimod and its major metabolite M2 in rat plasma (mean ± SD, n = 7).

Parameters	units	iguratimod	iguratimod metabolite
$t_{1/2}$	h	5.12 ± 2.45	3.70 ± 0.85
C_{max}	mg·L ⁻¹	14.16 ± 2.24	12.56 ± 3.58
T_{max}	h	5.86 ± 2.19	9.14 ± 1.07
AUC_{0-24h}	mg·h·L ⁻¹	197.42 ± 16.58	151.79 ± 41.47
Cl_z	L·h ⁻¹ ·kg ⁻¹	0.05 ± 0.00	0.07 ± 0.02
V_z	L·kg ⁻¹	0.34 ± 0.13	0.38 ± 0.19

Conclusion

Our study demonstrated that M2 is a major metabolite of iguratimod in phase I metabolism, and iguratimod is mainly metabolized by CYP2C9 and CYP2C19, especially by CYP2C9. Meanwhile, we developed a highly selective and reproducible HPLC method for simultaneous determination of iguratimod and its metabolite in rat plasma. The method could be useful for pharmacokinetic study in rats. In this experiment, the metabolic pathways of iguratimod were verified just in 7 recombinant human CYP450s (CYP1A2, CYP2C8, CYP2C9, CYP2C19, CYP2D6, CYP2E1, and CYP3A4) whole yeast cell reaction systems. These metabolic enzymes are the metabolic subtypes of most drugs. However, there may be other phase I metabolic pathways for iguratimod, which needs further investigations.

Acknowledgement

The authors would like to acknowledge Nanjing BRT-Biomed Co., Ltd. for providing recombinant human CYP450s yeasts. This study was supported by the Program for Jiangsu Province "Shuang Chuang" Team, China (to Chang-qing YANG, Grant (2014) 155).

References

- (1) Luo Q, Sun Y, Liu W, Qian C and Jin B. A Novel Disease-Modifying Antirheumatic Drug, Igaratimod, Ameliorates Murine Arthritis by Blocking IL-17 Signaling, Distinct from Methotrexate and Leflunomide. *J. Immunol.* (2013) 191: 4969-78.
- (2) Yamaori S, Takami K, Shiozawa A and Sakuyama K. *In-vitro* Inhibition of CYP2C9-Mediated Warfarin 7-Hydroxylation by Igaratimod: Possible mechanism of iguratimod-warfarin interaction. *Biol. Pharm. Bull.* (2015) 38: 441-7.
- (3) Zhou T, Ding L, Li XM and Zhang F. Determination of iguratimod in rat plasma by high performance liquid chromatography: method and application. *Biomed. Chromatogr.* (2008) 22: 260-4.
- (4) Gao J, Zhi DG, Zeng Y, Zhang P and Liu CX. Studies on preclinical pharmacokinetics of Igaratimod (T-614). *Drug Eval. Res.* (2009) 1: 19-28.
- (5) Peters FT, Bureik M and Maurer HH. Biotechnological synthesis of drug metabolites using human cytochrome P450 isozymes heterologously expressed in fission yeast. *Bioanalysis* (2009) 1: 821-30.
- (6) Chen L, Qian W and Yang CQ. Application of recombinant human metabolic enzyme yeast system in metabolic research of new drug *in-vitro*. *Chin. J. New Drugs* (2016) 25: 277-82.
- (7) Drăgan CA, Peters FT, Bour P, Schwaninger AE, Schaan SM, Neunzig I, Widjaja M, Zapp J, Kraemer T, Maurer HH and Bureik M. Convenient gram-scale metabolite synthesis by engineered fission yeast strains expressing functional human P450 systems. *Appl. Biochem. Biotechnol.* (2011) 163: 965-80.
- (8) Battershell C, Malhotra D and Hopfinger AJ. Inhibition of dihydrofolatereductase: structure-activity correlations of quinazolines based upon molecular shape analysis. *J. Med. Chem.* (1981) 24: 812-8.
- (9) Yasuda K, Ikushiro S, Kamakura M, Ohta M and Sakaki T. Metabolism of sesamin by cytochrome P450 in human liver microsomes. *Drug Metab. Dispos.* (2010) 38: 2117-23.
- (10) Peters FT, Dragan CA and Wilde DR. Biotechnological synthesis of drug metabolites using human cytochrome P450 2D6 heterologously expressed in fission yeast exemplified for the designer drug metabolite 4-hydroxymethyl- α -pyrrolidinobutyrophenone. *Biochem. Pharmacol.* (2007) 74: 511-20.
- (11) Zhang J, Yuan Y, Li JH, Wang YY and Liu GY. Pharmacokinetics of iguratimod tablets in Chinese healthy volunteers. *Chin. J. Clin. Pharm.* (2007) 4: 234-7.

This article is available online at <http://www.ijpr.ir>