

Somatic embryogenesis: an alternative method for *in vitro* micropropagation

Sobri Hussein^{1*}, Rusli Ibrahim¹, Anna Ling Pick Kiong²

¹Agrotechnology and Bioscience Division, Malaysian Nuclear Agency, Bangi, 43000 Kajang, Selangor, Malaysia

²Department of Bioscience, Faculty of Engineering and Science, Universiti Tunku Abdul Rahman, 53300 Setapak, Kuala Lumpur, Malaysia

Abstract

Plant tissue culture is an alternative method of commercial propagation and is being used widely for the commercial propagation of a large number of plant species, including many medicinal plants. Somatic embryogenesis is a process by which asexual or somatic cells are induced to form embryos in culture. Somatic embryogenesis is a multi-step regeneration process starting with formation of pro-embryogenic masses, followed by somatic embryo formation, maturation and regeneration. This paper outlined the important processes involved in the somatic embryogenesis.

Keywords: Somatic embryogenesis; Micropropagation; Somatic embryo; Regeneration; Plant tissue culture.

INTRODUCTION

Plants are unique in their ability to produce somatic embryos (Jayasankar *et al.*, 2001). All somatic cells within a plant contain the entire set of information necessary to create a complete and functional plant (Wang, 1997). Somatic embryogenesis is another alternative to traditional vegetative propagation methods as it offers a rapid large-scale propagation system (Fuentes *et al.*, 2000; Find *et al.*, 1998; Capuana and Debergh, 1997). Furthermore, the plantlets produced through somatic embryogenesis are reported as true-to-type (Martin, 2003b).

Somatic embryogenesis is a process whereby embryos develop from somatic cells or tissues in tissue culture achieve maturity and subsequently germinate to form normal plants (Witjaksono, 1997). Von Arnold and co-workers (2002) further defined somatic

embryogenesis as a process in which a bipolar structure resembling a zygotic embryo develops from a non-zygotic cell without vascular connection with the original tissue. Most somatic embryos including those from woody species pass through the same stages of development as a zygotic embryo; the globular, heart-shaped, torpedo- and cotyledonary stage (Sobri *et al.*, 2005; Ramos *et al.*, 2003; Choi *et al.*, 2002a; Tejavathi *et al.*, 2000; Choi *et al.*, 1999). In woody species, the proliferating embryogenic tissues are often secondary embryos (Ibaraki and Kurata, 2001). Secondary somatic embryogenesis is a phenomenon whereby new somatic embryos are initiated from somatic embryos (Vasic *et al.*, 2001). Although similar structurally and physiologically, somatic embryos are lack of protective seed coats, nutritive accessory tissues and the quiescent resting phase as in zygotic embryos (Jayasankar *et al.*, 2001). Wang (1997) stated that somatic embryos contain embryo specific macro-molecules like storage proteins and lipids.

Somatic embryos can be obtained via either direct or indirect somatic embryogenesis. In direct embryogenesis, the embryos originate directly from tissues in the absence of conspicuous callus proliferation (Aly *et al.*, 2002) whereas indirect embryogenesis involves the callus proliferation and embryogenic tissues precede embryo development (Fuentes *et al.*, 2000). Explants from which direct embryogenesis is most likely to occur include ovules, zygotic and somatic embryos, and seedlings (von Arnold *et al.*, 2002). Induction of somatic embryos directly from plant tissue is the most desirable approach in order to avoid somaclonal variation (Aoshima, 2005). Furthermore, it reduces the time required for plant regeneration, which may be beneficial to minimize culture-induced genetic changes (Fuentes *et al.*, 2000). Efficient systems based

*Correspondence to: Sobri Bin Hussein, Ph.D.

Tel: +603 89250510; Fax: +603 8928256

E-mail: sobri@nuclearmalaysia.gov.my

on direct somatic embryogenesis have been developed for numerous species including alfalfa, forest trees, celery, red clover, *Camellia japonica* and chickpea (Yantcheva *et al.*, 1998).

In vitro propagation through somatic embryogenesis has been reported in many medicinal plants such as *Eurycoma longifolia* (Sobri *et al.*, 2005), *Ocimum basilicum* L. (Gopi and Ponnuragan, 2006) as well as in a few spruce and pine species (Haggman *et al.*, 2005; Arya *et al.*, 2000). For instance, more than 500 somatic embryo plantlets of loblolly pine were successfully established in the field in the year 1991 to 1993 (Tang *et al.*, 1998). Somatic embryogenesis of woody plants was achieved in the 1960's with *Citrus* spp., *Biota orientalis*, *Santalum album* and *Zamia integrifolia*; however, it was not until the 1980's that woody plant somatic embryogenesis was more widely reported (Karkonen, 2001). For many applications, somatic embryos have powerful advantages for mass propagation; that is, the high multiplication rate, the ease of use of liquid medium, the handling of enormous numbers of embryos at one time and the possible use of bioreactors (Hamidah *et al.*, 1997). Thus, it is a more preferred pathway than organogenesis because it offers better potential for multiplication (Yantcheva *et al.*, 1998). Besides, plant regeneration by somatic embryogenesis provides an ideal *in vitro* system for basic studies of plant cell biology and embryo development (Find *et al.*, 1998). Somatic embryogenesis also provides an ideal experimental process for investigating differentiation and for understanding the mechanisms of expression of totipotency in plant cells (Arya *et al.*, 2000). In addition, the plantlets arising from embryoids have a tap root system which is superior to adventitious roots (Sita, 1986). Thus, somatic embryogenesis can be one of the important tools especially for the plants that do not or have low percentage of tap root formation in the micropropagation system (Sobri *et al.*, 2006a).

Stages of somatic embryogenesis: Somatic embryogenesis is a multi-step regeneration process starting with formation of pro-embryogenic masses, followed by somatic embryo formation, maturation, desiccation and plant regeneration (von Arnold *et al.*, 2002). Although the somatic embryogenesis protocols largely depend on plant species and cultivars, the production of somatic embryos generally involves four main processes: (1) induction, (2) maintenance of embryogenic cultures, and (3) development of embryos (4) Regeneration of the embryos (Korban, 2005; Ibaraki and Kurata, 2001).

Initiation of embryogenic callus: Embryogenic callus is defined as a type of callus that is either able to produce somatic embryos or is capable of embryogenesis when it is subjected to proper culture conditions (Gairi and Rashid, 2004). It is usually easy to distinguish between embryogenic and non-embryogenic callus based on its morphology and colour (von Arnold *et al.*, 2002). According to Sun *et al.* (2003) and Yang *et al.* (2003) embryogenic callus tends to be either (1) compact, yellowish-white with nodular structures and slow growing or (2) highly friable with embryos interspersed among masses of dry loosely organized cells. Meanwhile, non-embryogenic callus also has two distinct types: (1) translucent-to-yellow-to-brown callus with a wet, rough, crystalline, unorganized appearance and composed of rather large elongated cells or (2) highly organized tissue composed of a loose callus matrix with proliferating root primordia (Kebebew *et al.*, 1998; Fei, 1997). In terms of the biochemical content, von Aderkas *et al.* (2002) stated that embryogenic callus contains twice the amount of starch compared to non-embryogenic callus and also overall amino acid content comparable to that of the zygotic embryos. Early identification and selection of somatic embryogenesis in plants can also be carried out using the biochemical markers such as protein and isoenzyme banding profiles (Sobri *et al.*, 2006b). Sobri *et al.* (2006b) reported that two proteins at 21 and 25 kDa were highly accumulated in the embryogenic callus of *Eurycoma longifolia*.

Somatic embryos have been induced from a variety of plant tissues, most frequently from zygotic embryos (Corredoira *et al.*, 2002), immature inflorescence (Lauzer *et al.*, 2000), roots (Chand and Sahrawat, 2002), leaves (Zhang *et al.*, 2000), hypocotyls segments (Choi *et al.*, 1999) or directly from zygotic buds or cotyledons (Kevers *et al.*, 2002). However, almost all somatic embryos in conifers and deciduous trees are derived from zygotic embryos (Wang, 1997). The conditions under which somatic embryogenesis has been achieved are quite variable for different woody species (Karkonen, 2001). In a majority of studies, auxins, particularly 2,4-D have been employed for the induction of somatic embryos (Umehara and Kamada, 2005; Gumerova *et al.*, 2003; Pareek and Kothari., 2003; Mary and Jayabalan, 1997). For example, initiation of embryogenic cultures of some woody species such as *Citrus clementina*, *C. sinensis* and *C. limon* require continuous exposure to 2,4-D (Kayim and Koc, 2006) while in the case of *Gymnema sylvestri*, a combination of 2,4-D and BA managed to produce the secondary embryos (Kumar *et al.*, 2002b). In some other cases such as rice (Gairi and

Rashid, 2004) and *Cucumis melo* L. (Kintzios *et al.*, 2002), an initial pretreatment of 2,4-D for two to three days is sufficient to induce secondary embryogenesis that will continue for years on a basal medium. Meanwhile, supplementation of other auxin, NAA to the medium enhanced the embryogenic response in another woody species, *Sesbania sesban* var. *bicolor* (Shahana and Gupta, 2002).

Proliferation of embryogenic callus: Embryogenic callus is often accompanied by non-embryogenic callus which grows faster than the embryogenic callus. Thus, repeated selection of embryogenic callus for subculture is critical (Fei, 1997). Subculture interval also has a great impact on maintenance of embryogenic callus. Embryogenic cultures of some species and some genotypes can be subcultured for a prolonged period on medium containing plant growth regulators, and still retain their full embryogenic potential (von Arnold *et al.*, 2002). Somaclonal variation has been shown to increase with the prolonged *in vitro* culture (Lakshmanan and Taji, 2000). Normally, embryogenic callus is maintained and proliferated in a medium similar to that used for initiation (Kevers *et al.*, 2002; Anand *et al.*, 2000; Selles *et al.*, 1999). However, the degree of embryo differentiation which takes place in the presence of auxin varies in different species (Hamidah *et al.*, 1997).

Development and maturation of somatic embryos: Somatic embryo formation through callus culture often starts when embryogenic callus is transferred to media whereby the concentration of auxin or auxin-like compounds is reduced or omitted (Umehara and Kamada, 2005; Yang *et al.*, 2003) although in some cases, somatic embryogenesis and maturation occur at the same growth regulator concentration (Martin, 2003c). With the depletion of auxin, the block on the expression of those genes required for the transition to the heart shape is removed (von Arnold *et al.*, 2002).

One of the most important events occurring during embryo development is the differentiation of the shoot and apical meristems (Stasolla and Yeung, 2003). During the maturation stage, the somatic embryos undergo various morphological (Schuller *et al.*, 2000) and biochemical changes (Blanco *et al.*, 1997). Incomplete maturation of somatic embryos is one of the most significant factors that account for low rates of plant conversion (Witjaksono, 1997). Somatic embryo maturation can be controlled by treatments with abscisic acid (Blochl *et al.*, 2005), sucrose and desiccation (Thomas, 2006; Jheng *et al.*, 2006). Other

supplements that influence development include osmoticants and gelling medium (von Aderkas *et al.*, 2002). Reports of maturation of somatic embryos for various conifers species emphasized the importance of reducing the osmotic water potential of the medium by increasing the concentration of sucrose or supplementing the medium with mannitol, sorbitol or polyethylene glycol (PEG MW 4000) in the presence of abscisic acid (Garin *et al.*, 2000).

Germination of embryos and plant conversion: According to Stasolla and Yeung (2003), the terms germination refers only to roots elongation, which may not be necessarily followed by shoot growth while conversion refers to growth of a functional shoot and root system and it is through this process that viable plantlets are produced. Germination of somatic embryos and regeneration into plants has been reported to be low for most species (Jayasankar *et al.*, 2001) probably due to the structural abnormality of somatic embryos (Lee *et al.*, 2002). In addition, only mature embryos with a normal morphology and which have accumulated enough storage materials and acquired desiccation tolerance at the end of maturation develop into normal plants (von Arnold *et al.*, 2002; Walker and Parrott, 2001).

Somatic embryos usually develop into small plants, comparable to the seedlings, on culture medium lacking plant growth regulators (Stefanello *et al.*, 2005; Park *et al.*, 2005). However, there are cases where auxin and cytokinin stimulate germination (von Arnold *et al.*, 2002). In *Zingiber officinale* Rosc. for instance, supplementation of 3.0 mg l⁻¹ BA kinetin in the regeneration medium converted the somatic embryos into plantlets (Guo and Zhang, 2005). Meanwhile, for some species such as *Picea glauca*, inclusion of extra compounds like casein hydrolysate at 1.0 g l⁻¹ (w/v) is also required (Garin *et al.*, 2000).

Regeneration of cell suspension: Liquid medium processes have been investigated widely with the aim of obtaining synthetic seeds on an industrial scale (Aberlenc-Bertossi *et al.*, 1999). However, application of various technologies for plant breeding purposes, viable protoplast and cytoplasm fusions, organelle transfer, *in vitro* selection and some molecular techniques rely on the availability of cell suspension cultures which maintain their regeneration potential (Remotti, 1995). Cell suspension cultures are generally initiated from compact or friable callus of which the latter type is generally considered to be the most suitable (Luczkiewicz *et al.*, 2002; Figueiredo *et al.*,

2000). The cell suspension culture composed of a heterogeneous population of embryogenic and non-embryogenic cells (Holme, 1998). The embryogenic cells contains a dense cytoplasm showing an evident nucleus and containing starch grains while the non-embryogenic cells were large, vacuolated without an evident nucleus with no cell division (Tribulato *et al.*, 1997).

The somatic embryos in the liquid medium underwent similar development stages as in the embryos cultured in the solid medium (Anand *et al.*, 2000). The presence of starch, which is often related to the embryogenic cells and is considered to be an indicator of development towards somatic embryos, has also been observed in the embryogenic cell suspension cultures of *Lilium longiflorum* Thunb (Tribulato *et al.*, 1997). However, Nhut *et al.* (2006) reported that the number of somatic embryos derived from embryogenic calli of *Lilium longiflorum* cultured in liquid medium, especially at a volume of 20 ml, were shown to be more than in solidified medium, 170 rather than 28, and the forms of somatic embryos in different developmental steps (globular, heart and cotyledon-shaped) were clearly distinguished under microscope. Besides, Choi *et al.* (1999) also observed that the somatic embryos formed on agar medium had larger cotyledons and a more intact epidermis compared to embryos formed in liquid medium. The lower somatic embryo multiplication on solid media may be due to binding of water and absorption of minerals and plant growth regulators by the gelling agents resulting in the depleted supply to the growing embryos (Sood *et al.*, 2002).

Unlike the limited reports on somatic embryogenesis in suspension culture of monocots, systems for somatic embryogenesis have been established using suspension cultures in some species of dicotyledons such as carrot, celery and caraway (Kim and Soh, 1996). Several studies involving embryogenic suspension cultures of woody species have also been reported in *Eleutherococcus senticosus* (Choi *et al.* 2002b; Choi *et al.*, 1999), *Valeriana edulis* spp. *procera* (Castillo *et al.*, 2000), *Picea abies* and *Picea sitchensis* (Find *et al.*, 1998). The embryogenic cell line obtained from cell suspension culture will be useful for production of useful secondary metabolite by metabolic engineering using genetic transformation and for cryo-preservation of genetic resources (Kim *et al.*, 2003), for large scale multiplication, efficient in vitro selection of putative genetic transformants as well as for the production of non-chimeric transgenics (Thiruvengadam *et al.*, 2006).

References

- Aberlenc-Bertossi F, Noirot M, Duval Y (1999). BA enhances the germination of oil palm somatic embryos derived from embryogenic suspension cultures. *Plant Cell Tissue Organ Cult.* 56: 53-57.
- Aly MAM, Rathinasabapathi B, Kelley K (2002). Somatic embryogenesis in perennial statice *Limonium bellidifolium*, plumbaginaceae. *Plant Cell Tissue Organ Cult.* 68: 127-135.
- Anand RP, Ganapathi A, Anbazhagan R, Vengadesan G, Selvaraj N (2000). High frequency plant regeneration via somatic embryogenesis in cell suspension cultures of cowpea, *Vigna unguiculata* (L.) walp. *In Vitro Cell Dev Biol.* 36: 475-480.
- Aoshima Y (2005). Efficient embryogenesis in the callus of tea (*Camellia sinensis*) enhanced by the osmotic stress or antibiotics treatment. *Plant Biotechnol.* 22: 277-280.
- Arya S, Kalis RK, Arya ID (2000). Induction of somatic embryogenesis in *Pinus roxburghii* Sarg. *Plant Cell Rep.* 19: 775-780.
- Blanco MDLA, Nieves N, Sanchez M, Borroto CG, Castillo R, Gonzalez JL, Escalona M, Baez E, Hernandez Z (1997). Protein changes associated with plant regeneration in embryogenic calli of sugarcane (*Saccharum* sp.). *Plant Cell Tissue Organ Cult.* 51:153-158.
- Bloch A, March GG, Sourdioux M, Peterbauer T, Richter A (2005). Induction of raffinose oligosaccharide biosynthesis by abscisic acid in somatic embryos of alfalfa (*Medicago sativa* L.). *Plant Sci.* 168: 1075-1082.
- Castillo P, Marquez J, Rubluo A, Hernandez G, Lara M (2000). Plant regeneration from callus and suspension cultures of *Valeriana edulis* ssp. *Procera* via simultaneous organogenesis and somatic embryogenesis. *Plant Sci.* 151: 115-119.
- Chand S, Sahrawat AK (2002). Somatic embryogenesis and plant regeneration from root segments of *Psoralea corylifolia* L. an endangered medicinally important plant. *In vitro Cell Dev Biol.* 28: 33-38.
- Capuana M, Debergh PC (1997). Improvement of the maturation and germination of horse chestnut somatic embryos. *Plant Cell Tissue Organ Cult.* 48: 23-29.
- Choi YE, Kim JW, Yoon ES (1999). High frequency of plant production via somatic embryogenesis from callus or cell suspension cultures in *Eleutherococcus senticosus*. *Ann Bot.* 83: 309-314.
- Choi YE, Ko SK, Lee KS, Yoon ES (2002a). Production of plantlets of *Eleutherococcus sessiliflorus* via somatic embryogenesis and successful transfer to soil. *Plant Cell Tissue Organ Cult.* 69: 201-204.
- Choi YE, Lee KS, Kim EY, Kim YS, Han JY, Kim HS, Jeong JH, Ko KS (2002b). Mass production of Siberian ginseng plantlets through large-scale tank culture of somatic embryos. *Plant Cell Rep.* 21: 24-28.
- Corredoira E, Vieitez AM, Ballester A (2002). Somatic embryogenesis in Elm. *Ann Bot.* 89: 637-644.
- Fei SZ (1997). *In vitro* regeneration of buffalograss [*Buchloe dactyloides* (Nutt.) Engelm.]. *Ph.D Thesis*, University of Nebraska, USA.
- Figueiredo SFL, Simoes C, Albarello N, Viana VRC (2000). *Rollinia mucosa* cell suspension cultures: Establishment and growth conditions. *Plant Cell Tissue Organ Cult.* 63: 85-92.
- Find JI, Kristensen MMH, Norgaard JV, Krogstrup P (1998). Effect of culture period and cell density on regrowth following cryopreservation of embryogenic suspension cultures of Norway Spruce and Sitka spruce. *Plant Cell Tissue Organ Cult.* 53:

- 27-33.
- Fuentes SRL, Calheiros MBP, Manetti-Filho J, Vieira LGE (2000). The effects of silver nitrate and different carbohydrate sources on somatic embryogenesis in *Coffea canephora*. *Plant Cell Tissue Organ Cult.* 60: 5-13.
- Gairi A, Rashid A (2004). TDZ-induced somatic embryogenesis in non-responsive caryopses of rice using a short treatment with 2,4-D. *Plant Cell Tissue Organ Cult.* 76: 29-33.
- Garin E, Bernier-Cardou M, Isabel N, Klimaszewska K, Plourde A (2000). Effect of sugars, amino acids and culture technique on maturation of somatic embryos of *Pinus strobes* on medium with two gellan gum concentrations. *Plant Cell Tissue Organ Cult.* 62: 27-37.
- Gopi C, Ponmurugan P (2006). Somatic embryogenesis and plant regeneration from leaf callus of *Ocimum basilicum* L. *J Biotechnol.* 126: 260-264.
- Gumerova EA, Galeeva EI, Chuyenkova SA, Rummyantseva NI (2003). Somatic embryogenesis and bud formation on cultured *Fagopyrum esculentum* hypocotyls. *Russian J Plant Physiol.* 50: 640-645.
- Guo Y, Zhang Z (2005). Establishment and plant regeneration of somatic embryogenic cell suspension cultures of the *Zingiber officinale* Rosc. *Sci Hortic.* 107: 90-96.
- Haggman H, Vuosku J, Sarjala T, Jokela A, Niemi K (2005). Somatic embryogenesis of pine species from functional genomics to plantation forestry. In *Somatic Embryogenesis: Plant Cell Monographs*. Second edition. A Mujib, Smaaj J. New York: Springer-Verlag Berlin. P: 119-140.
- Hamidah M, Abdul Karim AG, Debergh P (1997). Somatic embryogenesis and plant regeneration in *Anthurium scherzerianum*. *Plant Cell Tissue Organ Cult.* 48: 189-193.
- Holme IB (1998). Growth characteristics and nutrient depletion of *Misanthus × ogiformis* Honda 'Giganteus' suspension cultures. *Plant Cell Tissue Organ Cult.* 53: 143-151.
- Ibaraki Y, Kurata K (2001). Automation of somatic embryo production. *Plant Cell Tissue Organ Cult.* 65: 179-199.
- Jayasankar S, van Aman M, Li Z, Gray DJ (2001). Direct seeding of grapevine somatic embryos and regeneration of plants. *In Vitro Cell Dev Biol.* 37: 476-479.
- Jheng FY, Do YY, Liauh YW, Chung JP, Huang PL (2006). Enhancement of growth and regeneration efficiency from embryogenic callus cultures of *Oncidium* 'Gower Ramsey' by adjusting carbohydrate sources. *Plant Sci.* 170: 1133-1140.
- Karkonen A (2001). Plant tissue cultures as models for tree physiology: somatic embryogenesis of *Tilia cordata* and lignin biosynthesis in *Picea abies* suspension cultures as case studies. *Ph.D Thesis*, University of Helsinki, Finland.
- Kayim M, NK Koc (2006). The effects of some carbohydrates on growth and somatic embryogenesis in citrus callus culture. *Sci Hortic.* 109: 29-34.
- Kebebew A, Gaj MD, Maluszynski M (1998). Somatic embryogenesis and plant regeneration in callus culture of tef, *Eragrostis tef* (Zucc.) Trotter. *Plant Cell Rep.* 18: 154-158.
- Kevers C, Gasper T, Dommès J (2002). The beneficial role of different auxins and polyamines at successive stages of somatic embryo formation and development of *Panax ginseng* in vitro. *Plant Cell Tissue Organ Cult.* 70: 181-188.
- Kim JW, Soh WY (1996). Plant regeneration through somatic embryogenesis from suspension cultures of *Allium fistulosum* L. *Plant Sci.* 114: 215-220.
- Kim SW, In DS, Kim TJ, Liu JR (2003). High frequency somatic embryogenesis and plant regeneration in petiole and leaf explant cultures and petiole-derived embryogenic cell suspension cultures of *Hylomecon vernalis*. *Plant Cell Tissue Organ Cult.* 74: 163-167.
- Kintzios S, Sereti E, Bluchos P, Drossopoulos JB, Kitsaki CK, Liopa-Tsakalidis A (2002). Growth regulator pretreatment improves somatic embryogenesis from leaves of squash (*Cucurbita pepo* L.) and melon (*Cucumis melo* L.). *Plant Cell Rep.* 21: 1-8.
- Korban SS (2005). Somatic embryogenesis in Rose: Gene expression and genetic transformation. In *Somatic Embryogenesis: Plant Cell Monographs* (2) eds A Mujib, Smaaj J. pp 247-257. New York : Springer-Verlag Berlin.
- Kumar S, Agrawal V, Gupta SC (2002b). Somatic embryogenesis in the woody legume *Calliandra tweedii*. *Plant Cell Tissue Organ Cult.* 71: 77-80.
- Lakshmanan P, Taji A (2000). Somatic embryogenesis in leguminous plants. *Plant Biology* 2: 136 -148.
- Lee KS, Lee JC, Soh WY (2002). High frequency plant regeneration from *Aralia cordata* somatic embryos. *Plant Cell Tissue Organ Cult.* 68: 241-246.
- Luczkiewicz M, Zarate R, Dembinska-Migas W, Migas P, Verpoorte R (2002). Production of pulchelin E in hairy roots, callus and suspension cultures of *Rudbeckia hirta* L. *Plant Sci.* 163: 91-100.
- Martin KP (2003b). Plant regeneration through somatic embryogenesis on *Holostemma adakodien*, a rare medicinal plant. *Plant Cell Tissue Organ Cult.* 72:79-82.
- Martin KP (2003c). Plant regeneration through direct somatic embryogenesis on seed coat explants of cashew (*Anacardium occidentale* L.). *Sci Hortic.* 98: 299-304.
- Mary RJ, Jayabalan N (1997). Influence of growth regulators on somatic embryogenesis in sesame. *Plant Cell Tissue Organ Cult.* 49:67-70.
- Nhut DT, Hanh NTM, Tuan PQ, Nguet LTM, Tram NTH, Chinh NC, Nguyen NH, Vinh DN (2006). Liquid culture as a positive condition to induce and enhance quality and quantity of somatic embryogenesis of *Lilium longiflorum*. *Sci Hortic.* 110: 93-97.
- Pareek A, Kothari SL (2003). Direct somatic embryogenesis and plant regeneration from leaf culture of ornamental species of *Dianthus*. *Sci Hortic.* 98: 449-459.
- Park SY, Ahn JK, Lee WY, Murthy HN, Paek KY (2005). Mass production of *Eleutherococcus koreanum* plantlets via somatic embryogenesis from root cultures and accumulation of eleutherosides in regenerants. *Plant Sci.* 168: 1221-1225.
- Ramos EMJ, Palma GRD, Hernandez-Sotomayor SMT (2003). Changes in phosphatidylinositol and phosphatidylinositol monophosphate kinase activities during the induction of somatic embryogenesis in *Coffea arabica*. *Physiologia Plantarum* 119: 270-277.
- Remotti PC (1995). Primary and secondary embryogenesis from cell suspension cultures of *Gladiolus*. *Plant Sci.* 107:205-214.
- Schuller A, Kirchner-Neß R, Reuther G (2000). Interaction of plant growth regulators and organic C and N components in the formation and maturation of *Abies alba* somatic embryos. *Plant Cell Tissue Organ Cult.* 60: 23-31.
- Selles M, Viladomar F, Bastida J, Codina C (1999). Callus induction, somatic embryogenesis and organogenesis in *Narcissus confusus*: correlation between the state of differentiation and the content of galanthamine and related alkaloids. *Plant Cell Rep.* 18: 646-651.
- Shahana S, Gupta SC (2002). Somatic embryogenesis in *Sesbania sesban* var. *bicolor*: a multipurpose fabaceous woody species.

- Plant Cell Tissue Organ Cult.* 69: 289-292.
- Sita LG (1986). Sandalwood (*Santalum album* L.). In *Biotechnology in Agriculture and Forestry 1: Trees* I. ed. Y.P.S. Bajaj. pp 363-374. New York: Springer-Verlag Berlin.
- Sobri H, Rusli I, Anna LPK (2006a). Adventitious shoots regeneration from root and stem explants of *Eurycoma longifolia* Jack-An important tropical medicinal plant. *Int J Agr Res.* 1: 183-193.
- Sobri H, Rusli I, Anna LPK (2006b). Potential biochemical markers for somatic embryos of *Eurycoma longifolia* Jack. *J Plant Biol.* 49: 97-101.
- Sobri H, Rusli I, Anna LPK, Mohd Fadzillah N, Daud SK (2005). Micropropagation of *Eurycoma longifolia* via formation of somatic embryogenesis. *Asian J Plant Sci.* 4: 472-485.
- Sood A, Ahuja PS, Sharma M, Sharma OP, Godbole S (2002). *In vitro* protocols and field performance of elites of an important bamboo *Dendrocalamus hamiltonii* Nees et Arn. Ex Munro. *Plant Cell Tissue Organ Cult.* 71: 55-63.
- Stasolla C, Yeung EC (2003). Recent advances in conifer somatic embryogenesis: improving somatic embryo quality. *Plant Cell Tissue Organ Cult.* 74: 15-35.
- Stefanello S, LLD Vesco, Ducroquet JPHJ, Nodari RO, Guerra MP (2005). Somatic embryogenesis from floral tissues of feijoa (*Feijoa sellowiana* Berg). *Sci Hort.* 105: 117-126.
- Sun Y, Zhang X, Jin S, Liang S, Nie Y (2003). Somatic embryogenesis and plant regeneration in wild cotton (*Gossypium klotzschianum*). *Plant Cell Tissue Organ Cult.* 75: 247-253.
- Tang W, Quyang F, Guo ZC (1998). Plant regeneration through organogenesis from callus induced from mature zygotic embryos of loblolly pine. *Plant Cell Rep.* 17: 557-560.
- Tejavathi DH, Lakshmi Sita G, Sunita AT (2000). Somatic embryogenesis in Flax. *Plant Cell Tissue Organ Cult.* 63: 155-159.
- Thiruvengadam M, Mohamed SV, Yang CH, Jayabalan N (2006). Development of an embryogenic suspension culture of bitter melon (*Momordica charantia* L.). *Sci Hort.* 109: 123-129.
- Thomas TD (2006). Effect of sugars, gibberellic acid and abscisic acid on somatic embryogenesis in *Tylophora indica* (Burm.f.) Merrill. *Chinese J Biotechnol.* 22: 465-471.
- Tribulato A, Remotti PC, Loffler HJM, Tuyl JMV (1997). Somatic embryogenesis and plant regeneration in *Lilium longiflorum* Thunb. *Plant Cell Rep.* 17: 113-118.
- Umehara M, Kamada H (2005). Development of the embryo proper and the suspensor during plant embryogenesis. *Plant Biotechnol.* 22: 253-260.
- Vasic D, Alibert G, Skoric D (2001). Protocols for efficient repetitive and secondary somatic embryogenesis in *Helianthus maximiliani*. *Plant Cell Rep.* 20: 121-125.
- von Aderkas P, Rohr R, Sundberg B, Gutmann M, Dumont-BeBoux N, Lelu MA (2002). Abscisic acid and its influence on development of the embryonal root cap, storage product and secondary metabolite accumulation in hybrid larch somatic embryos. *Plant Cell Tissue Organ Cult.* 69: 111-120.
- von Arnold S, Sabala I, Bozhkov P, Dyachok J, Filonova L (2002). Developmental pathways of somatic embryogenesis. *Plant Cell Tissue Organ Cult.* 69: 233-249.
- Walker DR, Parrott WA (2001). Effect of polyethylene glycol and sugar alcohols on soybean somatic embryo germination and conversion. *Plant Cell Tissue Organ Cult.* 64: 55-62.
- Wang X (1997). Somatic embryo induction and plant regeneration in American Ginseng (*Panax quinquefolium* L.). *M.Sc. Thesis*, University of Guelph, Canada.
- Witjaksono T (1997). Development of protocols for Avocado tissue culture: somatic embryogenesis, protoplast culture, shoot proliferation and protoplast fusion. *Ph.D. Thesis*, University of Florida, USA.
- Yang YG, Guo YM, Guo Y, Guo ZC, Lin JX (2003). Regeneration and large-scale propagation of *Phragmites communis* through somatic embryogenesis. *Plant Cell Tissue Organ Cult.* 75: 287-290.
- Yantcheva A, Vlahova M, Antanassov A (1998). Direct somatic embryogenesis and plant regeneration of carnation (*Dianthus caryophyllus* L.). *Plant Cell Rep.* 18: 148-153.