

Assessment of the genetic diversity of almond (*Prunus dulcis*) using microsatellite markers and morphological traits

Ali Fathi¹, Behzad Ghareyazi¹, Ali Hagnazari², Mohammad Reza Ghaffari¹, Seyed Mostafa Pirseyedi¹, Saeid Kadkhodaei¹, Mohammad Reza Naghavi³, Mohsen Mardi^{1*}

¹Department of Genomics, Agricultural Biotechnology Research Institute of Iran (ABRII), Mahdasht Road, P.O. Box 31535-1897, Karaj, I.R. Iran ²Department of Agronomy and Plant Breeding, Faculty of Agriculture, University of Zanjan, P.O. Box 45195-313, Zanjan, I.R. Iran ³Department of Agronomy and Plant Breeding, Faculty of Agriculture, University of Tehran, P.O. Box 31587-11167, Tehran, I.R. Iran

Abstract

The genetic diversity among 56 almond (*Prunus dulcis*) genotypes was analysed using 35 microsatellite markers and 14 morphological traits. Analysis of morphological traits revealed a wide range of variation among the studied genotypes. Out of 35 simple sequence repeats (SSRs) markers, 25 were polymorphic, producing 215 alleles that varied from 2 to 16 with an average of 8.76 alleles per locus. Regression analyses revealed a positive correlation between the *CPPCT03* locus and kernel yield, kernel percentage, grain weight, leaf length and tree altitude. The results of analysis of molecular variance (AMOVA) indicated that approximately 4.5% of genetic variance was observed between the collection sites. Based on SSR data, cluster analyses showed that the studied almond genotypes were classified into five main groups. The results of the present study showed that microsatellite markers could be successfully used to assay genetic diversity among Iranian almond landraces/cultivars and to identify informative markers for improving traits in breeding programs.

Keywords: *Prunus dulcis*; Genetic relationship; Microsatellite; Informative markers.

INTRODUCTION

Almond [*Prunus dulcis* (Miller) D.A. Webb, syn. *Prunus amygdalus* Batsch] occupies a very peculiar

place among fruit trees (Miller *et al.*, 1989). Because of almond's tolerance to cold, drought and salinity, it is considered an important tree crop and is cultivated in different climatic regions of Iran. Breeding practices in *Prunus* face unique challenges resulting from the narrow genetic background of commercial cultivars (Scorza *et al.*, 1985). Morphological traits such as seed and kernel size, kernel yield, and blooming time are usually used for cultivar identification in almond (De iorgio and Polignano, 1999). However, morphological traits are limited because of their environmental fluctuations.

In recent years, molecular markers have been used to study genetic diversity and cultivar identification of peach and almond (Shiran *et al.*, 2007; Sorkheh *et al.*, 2007; Amirbakhtiar *et al.*, 2006; Kadkhodaei *et al.*, 2006; Sanchez-Pérez *et al.*, 2006; Xie *et al.*, 2006; Testolin *et al.*, 2000, 2004; Aranzana *et al.*, 2003). Methods based on knowledge provided by advances in molecular genetics, notably molecular markers, promise faster and more efficient approaches to cultivar improvement. In fact important tools such as molecular markers, maps, DNA sequences, and quantitative trait loci (QTLs) have been developed and made available to researchers, and applications at the breeding program level have already started (Dirlewanger *et al.*, 2004). Recently, DNA microarray-based genome composition analysis has also been used in comparative genomic studies of trees (Martinez-Gomez *et al.*, 2007). The objectives of the present study are to investigate the genetic diversity of major Iranian almond

*Correspondence to: Mohsen Mardi, Ph.D.
Tel: +98 261 2700845; Fax: +98 261 2700845
E-mail: mardi@abrii.ac.ir

landraces/cultivars, to identify their relationship to important foreign cultivars, and to introduce informative markers for important nut traits using microsatellite markers.

MATERIALS AND METHODS

Plant materials: Fifty-one *Prunus dulcis* landraces/cultivars from different provinces of Iran along with three and two registered cultivars from Spain and USA, respectively, were used in this study (Table 1). The trees with similar ages were sown in a randomized complete block design, with four replications, at the experimental field of the Agricultural Biotechnology Research Institute of Iran (ABRII), Isfahan.

Phenotypic analysis: Fourteen independent morphological traits including leaf shape, leaf length (cm), leaf width (cm), petiole length (cm), flowering duration (day), tree altitude (cm), frostbite kernel yield (g), kernel length (cm), kernel width (cm), kernel thickness (cm), nut weight (g), kernel nut weight (g) and kernel percentage were recorded, based on food and agriculture organization (FAO).

Microsatellite analysis: Total genomic DNA was extracted according to the method described by Doyle and Doyle (1987), with minor modifications. Thirty-five simple sequence repeat (SSR) markers, isolated from peach and almond were used in this study (Testolin *et al.*, 2004; Dirlewanger *et al.*, 2002). Amplification reaction products were separated on a 6% (w/v) dena-

Table 1. Almond landraces/cultivars included in this study.

No.	Genotype name	Landrace/ Cultivar	Collection site	No.	Genotype name	Landrace/ Cultivar	Collection site
1	Post nazok1 (pk1)	Cultivar	Shiraz	30	H 8	Landrace	Hamadan
2	Monagha Shiraz	Cultivar	Shiraz	31	H 9	Landrace	Hamadan
3	S 8	Landrace	Shiraz	32	H 22	Landrace	Hamadan
4	S 21	Landrace	Shiraz	33	H20	Landrace	Hamadan
5	S 27	Landrace	Shiraz	34	H 6	Landrace	Hamadan
6	Post nazok 2 (pk2)	Cultivar	Shiraz	35	H 27	Landrace	Hamadan
7	S18	Landrace	Shiraz	36	H 5	Landrace	Hamadan
8	S17	Landrace	Shiraz	37	H 30	Landrace	Hamadan
9	S7	Landrace	Shiraz	38	H 7	Landrace	Hamadan
10	Sangi	Cultivar	Shiraz	39	H 4	Landrace	Hamadan
11	Mamaei1	Cultivar	Isfahan	40	H 18	Landrace	Hamadan
12	Mamaei2	Cultivar	Isfahan	55	H0	Landrace	Hamadan
13	Tadjeri	Cultivar	Isfahan	41	Ferragnes	Cultivar	France
14	Dobahre1	Cultivar	Isfahan	42	Sahand	Cultivar	Azerbaijan
15	Monagha Najafabad	Cultivar	Isfahan	43	Spain 200	Cultivar	Spain
16	Dobahre2	Cultivar	Isfahan	44	Shokofe	Cultivar	Azerbaijan
17	103	Landrace	Isfahan	45	Yalda	Cultivar	Azerbaijan
18	101-1	Landrace	Isfahan	46	Nonpareil	Cultivar	USA
19	101-2	Landrace	Isfahan	47	AR (1)	Landrace	Arak
20	Rabii	Cultivar	Isfahan	48	AR (2)	Landrace	Arak
21	H 12	Landrace	Hamadan	49	AR (3)	Landrace	Arak
22	H 15	Landrace	Hamadan	50	AR (4)	Landrace	Arak
23	H 16	Landrace	Hamadan	51	AR (5)	Landrace	Arak
24	H 17	Landrace	Hamadan	52	AR (8)	Landrace	Razan
25	H 11	Landrace	Hamadan	53	AR (6)	Landrace	Arak
26	H 21	Landrace	Hamadan	54	AR (7)	Landrace	Arak
27	H 10	Landrace	Hamadan	56	Spain 230	Cultivar	Spain
28	H 13	Landrace	Hamadan	57	Harir	Cultivar	Azerbaijan
29	H 19	Landrace	Hamadan	-	-	-	-

The names of the landraces/cultivars have been identified by their collection sites.

Table 2. Measured morphological traits.

Trait	Mean	Minimum	Maximum	Percentage of coefficient of Variation (CV%)
Leaf shape	8.95	1	14	55
Leaf length (cm)	5.26	3.5	8	19
Leaf width (cm)	1.68	1	2.2	17
Petiole length (cm)	17	10	28	24
Flowering duration (day)	8.94	4	16	31
Tree altitude (cm)	130.33	60	290	37
Frostbite	1.5	1	2	33
Kernel yield (gr)	2.3	0.03	11.4	107
Kernel length (cm)	3.05	2.5	3.91	10
Kernel width (cm)	1.99	1.67	2.51	11
Kernel thickness (cm)	1.14	1.23	1.85	12
Nut weight (gr)	34.66	14	56.7	33
Kernel nut weight (gr)	9.33	6.29	16.8	24
Kernel percentage (%)	28.72	15.91	81.11	47

turing polyacrylamide gel using a Sequi-Gen GT Sequencing Cell 30 cm gel apparatus (BioRad Laboratories Inc., Hercules, CA, USA). The amplified fragments were detected by the silver staining method as described by Bassam *et al.* (1991). The gels were visually scored by two independent observations.

Data analysis: Each polymorphic fragment was scored as either present (1) or absent (0) across all genotypes. The data were used to calculate the similarity matrix among cultivars employing simple matching coefficients. The similarity matrix was then used to construct dendrograms using the unweighted pair group method with arithmetic averages (UPGMA). This was achieved by employing the sequential, agglomerative, hierarchical, and nested clustering (SAHN) using the numerical taxonomy and multivariate analysis system (NTSYS-PC), version 2.00 (Rohlf, 1998). Observed heterozygosity (H_o) and expected heterozygosity (H_e) were calculated using the POPGENE version 1.32 (Yeh *et al.*, 1997). The degree of polymorphism was quantified using the polymorphic information content (PIC). Probability of identity (PI) was estimated according to Paetkau *et al.* (1995). Analysis of molecular variance (AMOVA) was performed using the Arlequin version 2.00 (Schneider *et al.*, 2000) to determine genetic variation (Nei, 1972). Average value of the Shannon index was also measured (Shannon and Weaver, 1949). Informative markers were determined by stepwise regression using the SPSS software version 10.0 for windows (SPSS Inc., Chicago, IL).

RESULTS

Morphological trait analysis: Mean, maximum, minimum and the percentage of coefficient of variation (CV%) of 14 morphological characters are shown in Table 2. A large diversity in the characters was observed, indicating a high level of variation in the studied plant materials.

SSR marker analysis: The results of this study showed cross amplification ability of microsatellite markers among the studied almond genotypes. Out of 35 SSR markers, 25 were polymorphic and produced 215 alleles. The number of alleles per locus ranged from 2 to 16, with an average of 8.76 (Table 3). Average value of the Shannon index was 1.79, which varied from 0.35 in *UDP96-008* to 2.6 in *CPPCT3*. Mean H_e across microsatellite loci ranged from 0.92 in *CPPCT3* to 0.17 in *UDP96-008*. The highest level of observed heterozygosity was found in *XAM18* and *CPPCT22* and the lowest in *UDP96-008*. According to PI, the most informative loci were *UDP98-412* and *CPPCT3* with values of 0.041 and 0.042, respectively. PIC for these two loci was greater (0.7) than others. The least informative locus was *XAM04* with PI of 0.98 and PIC of 0.159, followed by *XAM18* with PI and PIC values of 0.494 and 0.0018, respectively. The average of PI and PIC values for all loci were 0.258 and 0.475, respectively (Table 3). Rare polymorphic alleles (*i.e.* those with a frequency of ≤ 0.005) and their weights were determined for the pur-

Table 3. Characterized SSR markers amplified from almond (*Prunus dulcis*).

Locus name	Sequence primers (5'→3')	Average Size Base Pair (bp)	Number of alleles	Shannon Index	Ho	He	PIC	PI
<i>XAM01</i>	ATAAATATATACACACACACAC CATATAGGGTCAAGGGAGTG	175	---	---	---	---	---	---
<i>XAM02</i>	CGTGAGGTCTCACTCTCTC ATTTAAGGGTCTGGTCA	146	9	85.1	660.0	834.0	449.0	215.0
<i>XAM03</i>	GCAAGAACGGTTTCTTTTC GATGGACCAACTCAAGC	191	---	---	---	---	---	---
<i>XAM04</i>	TACATTTATCCCCGGTA GAAGCTCCATTCTTGTA	156	3	03.1	537.0	625.0	159.0	98.0
<i>XAM05</i>	CACACACAAACACAAATGC TTGTGCTTTCATGGAC	172	10	96.1	518.0	873.0	676.0	083.0
<i>XAM06</i>	TCTCCAAGGGGATAAGCA AGGCACCTGTCCCCTACA	156	---	---	---	---	---	---
<i>XAM07</i>	CGCTTTGCATACATACAAGT AGGAACTGGGATTAGAGA	149	---	---	---	---	---	---
<i>XAM08</i>	ACATCTCTCTCCTCCATGC TCTCTGGCAGCACAAAGC	220	11	22.2	673.0	888.0	658.0	054.0
<i>XAM09</i>	TCACATACACGTTGGGTTTC TGTGATTTGTGTGTGTGC	157	9	97.1	571.0	850.0	69.0	071.0
<i>XAM10</i>	ATTGTTTTCCCTGGTA GAATCTCAACTCGGAAACG	94	---	---	---	---	---	---
<i>XAM11</i>	CCGGGGCTTTATAAAT TGTGATGGCCAGAGCTT	199	9	89.1	673.0	836.0	632.0	134.0
<i>XAM12</i>	CCTGTCACAAAGATGCAA CATTTTCCAGTAGTCCA	162	2	56.0	350.0	374.0	311.0	419.0
<i>XAM13</i>	AATACACACGGCCACAC AAGCATCGTCACTAGCC	165	11	10.2	777.0	860.0	545.0	102.0
<i>XAM14</i>	CCATCGCTTGCATT CCGTGTGTGTTGTGTG	140	---	---	---	---	---	---
<i>XAM15</i>	AACTATAAAAATACACACACACA CATCATCGGCTTTATTAG	193	---	---	---	---	---	---
<i>XAM16</i>	GCACCAAAACACAACCTGA GTGTTGCCAATGTTGAT	172	8	74.1	673.0	810.0	38.0	243.0
<i>XAM17</i>	CACGTACATTTGTGACTGC GTGTAATGCCACAGATGC	163	---	---	---	---	---	---
<i>XAM18</i>	CGTCTCATTTTCCCATT CGATGGAGGAGCACT	174	10	83.1	910.0	815.0	0018.0	494.0
<i>XAM19</i>	CCGTGATACACTAACAACT TGCCAAAGTAAAGTGCCTA	175	12	30.2	732.0	892.0	566.0	087.0

Table 3. Continued

Locus name	Sequence primers (5'→3')	Average Size Base Pair (bp)	Number of alleles	Shannon Index	Ho	He	PIC	PI
<i>XAM20</i>	AGAAAGCTGCACCTGGTA GCTTATTCGTGTGTG	138	9	06.2	821.0	867.0	57.0	13.0
<i>UDP98-409</i>	TGATGGGTTTTATGGTTTC GGACTCTTATCCTCTATCAACA	129	---	---	---	---	---	---
<i>UDP96-015</i>	TTGACCTATTTGTTCTGTC TAGTCAACAACATCCCCCG	174	---	---	---	---	---	---
<i>UDP96-008</i>	TGTACACACCCCTCAGCCTG GCTGAGGTTTCAGGTGAGTG	165	---	---	---	---	---	---
<i>UDP98-412</i>	GGGAAAGTTTCTGCTGCAC TGAAAGACGACGATGATGA	129	---	---	---	---	---	---
<i>Pchgms1</i>	GTAATAATGCCCATTTGTCAATC ATCATTGAACTACGTCAATCCTC	194	---	---	---	---	---	---
<i>PS7a2</i>	GGGAAATAGATAAGATG TAATGGTGGTGTTCATT	-	---	---	---	---	---	---
<i>CPPCT3</i>	GTAACGAAGAAGTTACGGG AACTGTCGCTGCTGGGTT	160	16	60.2	750.0	921.0	707.0	042.0
<i>CPPCT4</i>	TCATTCCGAAAGACGACCGT GTCTAGGCACGTTGCTAG	250	---	---	---	---	---	---
<i>CPPCT16</i>	AATTCCCTATGGAAATTAGA CGCATATTATAGGTAGGAAA	191	13	29.2	771.0	886.0	609.0	112.0
<i>CPPCT17</i>	GTGACATGCATGCACATAACA TGCAAAATGCAATTCATAAAGG	177	12	16.2	690.0	863.0	626.0	113.0
<i>CPPCT22</i>	CAATTAGCTAGAGAAATTATTG GACAAGAAAGCAAGTAGTTG	240	5	31.1	771.0	690.0	053.0	535.0
<i>CPPCT24</i>	TTCTCCCAAAAACCAAAACC TCATTGGCTGCTAAGTGTCTT	180	6	50.1	571.0	763.0	41.0	42.0
<i>CPPCT27</i>	GAGCAGTTCATAAGTTGGAAC CGATAAAGATTTGACTGCATG	114	11	12.2	842.0	866.0	513.0	155.0
<i>CPPCT30</i>	TGAATAATTGTTCTCAAATTC CTCTAGGCAAGAGATGAGA	198	5	92.0	368.0	466.0	185.0	556.0
<i>CPPCT33</i>	TCAGCAAACTAGAAACAACC TTGCAATCTGGTTGATGTT	151	9	07.2	740.0	873.0	55.0	17.0

Ho, He, PIC, and PI are observed heterozygosity, expected heterozygosity, polymorphic information content and probability of identity, respectively. The loci (*XAM01* to *XAM20*) have been determined by Testolin et al. (2004) and the others by Dirlewanger et al. (2002).

Table 4. Rare polymorphic alleles and their weight for use in almond identification.

Locus name	Rare allele weight	Genotype name
XAM06	145	H 12
XAM02	190	Mamaei 1
XAM02	170	H 21
XAM05	212	AR (3)
XAM05	147	Sangi
XAM08	195	AR (5)
XAM08	307	H 13
XAM08	240	Nonpareil and S 8
XAM09	131	Shokofe
XAM16	183	Fragness
XAM16	160	S 21
XAM18	119	H 19
CPPCT16	204	H 8
CPPCT03	225	Rabii
CPPCT03	186	Spain 230 and Sahand
CPPCT17	147	AR (6)
CPPCT27	79	H 4
CPPCT30	250	Shokofe
UDP96-008	147	Spain 200

For genotype and locus names see Table 1 and 3, respectively.

pose of rapid cultivar identification (Table 4). Regression analyses revealed that there was a positive correlation between the *CPPCT03* locus and kernel yield ($\beta = 0.424$), kernel percentage ($\beta = 0.49$), grain weight ($\beta = 0.35$), leaf length ($\beta = 0.32$) and tree altitude ($\beta = 0.327$) (Table 5).

Based on sampling sites, average *He* was 0.697 and the largest heterozygosity was observed for cultivars from Hamadan (0.731). The results of AMOVA indicated that approximately 4.5% of genetic variance belonged to between collected sites (Table 6). Based on SSR data, the studied almond genotypes were classified into five main groups (Fig. 1). The first cluster included some landraces and cultivars from the Shiraz, Isfahan, Hamadan and Arak provinces. The second cluster included two sub-clusters: the first sub-cluster contained 4 landraces from the Shiraz province and the second sub-cluster contained registered cultivars from Spain, USA and Azerbaijan. Two landraces from Shiraz and Arak provinces were gathered into cluster III. One registered cultivar from USA (HO) and one registered cultivar from Azerbaijan (Harir) were located in two distinct clusters (IV and V).

DISCUSSION

The results of this study support those of Sosinski *et al.* (2000), regarding the cross amplification ability of microsatellite markers across the *Prunus* species. High level of heterozygosity for all loci (0.697) can be attributed to cross pollination and the self-incompatibility nature of almond. The high values of polymorphic loci (71%), average number of alleles per locus (8.76), *He* (0.775), average polymorphism information content (0.475) and PI (0.258) observed in this study indicate that SSR markers are able to identify genetic variation among the studied almond genotypes. According to PI and PIC values, *CPPCT3*, *UDP98-412*, *UDP96-409*, *XAM05*, *XAM08*, *XAM09*, *XAM15* and *XAM19* are the best loci for further studies of almond genetic diversity. The percentage of polymorphic SSR loci (71%) in this study was much higher than that estimated for RFLPs (21.9%), suggesting that SSRs can act as better systems for almond cultivar identification (Eldredge *et al.*, 1992).

During this research, alleles were identified that correlated with yield-related traits. The allele belonging to the *XAM09* locus had a positive correlation with blooming duration (0.418) (Table 5). In addition, *CPPCT17* was found to be an informative marker for nut weight, average kernel thickness and leaf width (Table 5).

In this investigation, cluster analyses showed that most Iranian landraces are well separated from the Spanish and American (USA) cultivars, indicating that they may be native to Iran. However, Shiraz almond landraces are assigned to the same group as the Spanish and American cultivars. A possible explanation is that they might carry a common genetic background. According to the results of this study, SSR data failed to separate genotypes based on their sampling sites. Germplasm migration or insufficient SSR markers can explain this incomplete separation. The results show that Iranian registered cultivars including Yalda, Shokofe and Sahand are similar to the foreign cultivars.

Informative markers are most applicable for breeding purposes. These markers have previously been used in the identification of peach and nectarine varieties (Manubens *et al.*, 1999). A combination of molecular and morphological data is the best choice to find informative markers. In summary, results of the present study reveal that microsatellite markers can be

Table 5. Regression between morphological and molecular data to define informative markers.

Trait	Locus name	Adjusted R ²	P-value	Standard β
Leaf shape	XAM13	0.105	0.001	-0.34
Leaf length	CPPCT27	0.153	0.00	0.403
Leaf length	UDP96-412	0.141	0.00	0.389
Leaf length	CPPCT16	0.093	0.003	0.322
Leaf length	CPPCT03	0.091	0.003	0.32
Leaf width	CPPCT17	0.188	0.00	0.425
Leaf width	XAM08	0.107	0.001	-0.343
Petiole length	XAM02	0.118	0.001	0.359
Petiole length	XAM15	0.088	0.003	-0.314
Flowering duration	XAM09	0.167	0.00	0.418
Flowering duration	CPPCT27	0.128	0.00	-0.372
Flowering duration	Pchgm1	0.121	0.01	0.362
Flowering duration	XAM11	0.074	0.007	0.292
Tree altitude	XAM08	0.13	0.00	-0.375
Tree altitude	CPPCT24	0.089	0.005	-0.305
Tree altitude	XAM11	0.077	0.006	-0.296
Tree altitude	CPPCT03	0.096	0.002	0.327
Tree altitude	XAM13	0.097	0.002	0.328
Frostbite	XAM20	0.138	0.00	0.385
Kernel yield	CPPCT03	0.17	0.00	0.424
Kernel yield	XAM19	0.122	0.001	0.364
Kernel length	UDP96-08	0.137	0.00	-0.384
Kernel length	XAM15	0.133	0.00	0.378
Kernel length	XAM16	0.072	0.007	0.288
Kernel width	CPPCT03	0.176	0.00	-0.431
Kernel width	XAM02	0.087	0.004	-0.313
Kernel width	XAM13	0.128	0.00	-0.372
Kernel thickness	UDP98-409	0.147	0.00	0.396
Kernel thickness	XAM15	0.074	0.007	-0.292
Kernel thickness	CPPCT17	0.129	0.00	0.373
Kernel thickness	Pchgm1	0.101	0.002	0.334
Kernel thickness	XAM08	0.083	0.004	-0.306
Nut weight	XAM08	0.125	0.001	-0.367
Nut weight	CPPCT17	0.102	0.002	0.336
Nut weight	CPPCT17	0.076	0.006	0.295
Nut weight	CPPCT03	0.112	0.001	0.35
Nut weight	CPPCT02	0.083	0.005	-0.305
Kernel weight	CPPCT33	0.149	0.00	0.398
Kernel weight	Pchgm1	0.093	0.003	0.323
Kernel percentage	CPPCT03	0.232	0.00	0.491
Kernel percentage	XAM18	0.039	0.039	-0.244

For locus name see Table 3.

Table 6. Analysis of molecular variance (AMOVA) and variance components for total genetic differentiation in almond based on collection sites.

Source of variation	Degree of freedom	Mean squared	Variance component	Percentage of variation	P value
Among collection sites	4	4.98	0.116	4.41	0.001
Within collection sites	109	2.26	2.51	95.43	-
Total	113	7.25	2.63	100	-




Figure 1. Dendrogram showing the relationships between 57 almond accessions using simple matching index and unweighted pair group method whit arithmetic mean (UPGMA).

successfully used to assay genetic diversity among Iranian almond landraces/cultivars and to identify informative markers for breeding of important traits.

Acknowledgments

We would like to thank the Agricultural Biotechnology Research Institute of Iran (ABRII) for funding this study.

References

- Amirbakhtiar N, Shiran BH, Moradi H, Sayed-Tabatabaei BE (2006). Molecular characterization of almond cultivars using microsatellite markers. *ISHS Acta Horticulture 726: IV International Symposium on Pistachios and Almonds*. Tehran, Iran.
- Aranzana MJ, Pineda A, Cosson P, Dirlwanger E, Ascasibar J, Cipriani G, Ryder CD, Testolin R, Abbott A, King GJ, Iezzoni AF, Arus P (2003). A set of simple-sequence repeat (SSR) markers covering the *Prunus* genome. *Theor Appl Genet*. 106:

- 819-825.
- Bassam BJ, Caetano-Anollés G, Gresshoff PM (1991). Fast and sensitive silver staining of DNA in polyacrylamide gels. *Anal Biochem.* 196: 81-84.
- De iorgio D, Polignano G (1999). Evaluating almond biodiversity of cultivars grown a germoplasm collection field in south of Italy. Proc. *ISCO Conference*, May 23-28, Purdue University, West Lafayette, Indiana, USA.
- Dirlewanger E, Cosson P, Tavaud M, Aranzana MJ, Poizat C, Zanetto A, Arus P, Laigret F (2002). Development of microsatellite markers in peach [*Prunus persica* (L.) Batsch] and their use in genetic diversity analysis in peach and sweet cherry (*Prunus avium* L.). *Theor Appl Genet.* 105: 127-138.
- Dirlewanger E, Graziano E, Joobeur T, Garriga-Caldere F, Cosson P, Howad W, Arus P (2004). Comparative mapping and marker-assisted selection in Rosaceae fruit crops. *PNAS.* 101: 9891-9896.
- Doyle JJ, Doyle JL (1987). A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem Bull.* 19: 11-15.
- Eldredge L, Ballard R, Baird W, Abbott A, Morgens P, Callahan A, Scorza R, Monet R (1992). Application of RFLP analysis to genetic mapping in peaches. *HortScience* 27: 160-163.
- Kadkhodaei S, Aghdai SRT, Grigorian V, Moghadam M, Hashemi SMM (2006). A study on genetic variation among some wild almond species using RAPD markers. *ISHS Acta Horticulture 726: IV International Symposium on Pistachios and Almonds*. Tehran, Iran.
- Manubens A, Lobos S, Jadue Y, Toro M, Messina R, Liadser M, Seelenfreund D (1999). DNA isolation and AFLP fingerprinting of nectarine and peach varieties (*Prunus persica*). *Plant Mol Biol Rpt.* 17: 255-267.
- Martinez-Gomez P, Sanchez-Perez R, Dicenta F, Howad W, Arus P, Gradziel TM (2007). Genome mapping and molecular breeding in plants. *Springer Berlin Heidelberg.* 4: 229-242.
- Miller PJ, Parfitt DE, Weinbaum SA (1989). Outcrossing in peach. *HortScience* 24: 359-360.
- Nei M (1972). Genetic distance between collections. *Amer Naturalist.* 106: 283-292.
- Paetkau D, Calvert W, Stirling I, Strobeck C (1995). Microsatellite analysis of population structure in Canadian polar bears. *Mol Ecol.* 4: 347-354.
- Rohlf FJ (1998). *NTSYSpc: Numerical Taxonomy and Multivariate Analysis System*, Version 2.02, Exeter Software, Setauket, New York.
- Sanchez-Pérez R, Dicenta F, Martinez-Gomez P, Howad W, Arus P (2006). Construction of linkage map and QTL analysis of agronomic traits in almon using SSR markers. *ISHS Acta Horticulture 726: IV International Symposium on Pistachios and Almonds*. Tehran, Iran.
- Schneider S, Roessli D, Excoffier L (2000). *Arlequin: A Software for Population Genetics Data Analysis*, Version 2.000 Genetics and Biometry Laboratory, Dept. of Anthropology, University of Geneva, Switzerland.
- Shannon CE, Weaver W (1949). *The mathematical theory of communication*. Urbana, Ill: Univer. of Illinois Press.
- Shiran B, Amirbakhtiar N, Kiani S, Mohammadi SH, Seyed-Tabatabaei BT, Moradi H (2007). Molecular characterization and genetic relationship among almond cultivars assessed by RAPD and SSR markers. *Sci Hort.* 111: 280-292.
- Scorza R, Mehlenbacher SA, Lighner GW (1985). Inbreeding and coancestry of freestone peach cultivars of the eastern United States and implications for peach germplasm improvement. *J Am Soc Hort Sci.* 110 : 547-552.
- Sorkheh K, Shiran B, Gradziel, TM, Epperson BK, Mrtinez-Gomez P, Asadi E (2007). Amplified fragment length polymorphism as a tool for molecular characterization of almond germplasm: genetic diversity among cultivated genotypes and related wild species of almond, and its relationships with agronomic traits. *Euphytica* 156: 327-344.
- Sosinski B, Gannavarapu M, Hager LD, Beck E, King GJ, Ryder CD, Rajapakse S, Baird WV, Ballard RE, Abbott AG (2000). Characterization of microsatellite markers in peach [*Prunus persica* (L.) Batsch]. *Theor Appl Genet.* 101: 421-428.
- Testolin R, Marrazzo T, Cipriani G, Quarta R, Verde I, Dettori M, Pancaldi M, Sansavini S (2000). Microsatellite DNA in peach (*Prunus persica* L. Batsch) and its use in fingerprinting and testing the genetic origin of cultivars. *Genome* 43: 512-520.
- Testolin R, Messina R, Lain O, Marrazzo MT, Huang WG, Cipriani G (2004). microsatellites isolation in almond from an AC-repeat enriched library. *Mol Ecol Notes.* 4: 459-461.
- Xie H, Sui Y, Chang FQ, Xu Y, Ma RC (2006). SSR allelic variation in almond (*Prunus dulcis* Mill.). *Theor Appl Genet.* 112: 366-72.
- Wright S (1951). The genetical structure of collections. *Ann Eugen.* 15: 323.
- Yeh FC, Yang RC, Boyle T (1997). POPGENE version 1.21. (URL <http://www.ualberta.ca/~fyeh>).