

Molecular characterization of a *Salmonella* Typhimurium isolate from Caspian pony

Taghi Zahraei Salehi^{1*}, Mohammad Javad Gharagozlou², Nemat Shams³, Omid Madadgar¹, Bahar Nayeri Fasaee¹, Ramak Yahyaraeyat¹

¹Department of Microbiology, Faculty of Veterinary Medicine, University of Tehran, P.O. Box 14155-6453, Tehran, I.R. Iran ²Department of Pathology, Faculty of Veterinary Medicine, University of Tehran, P.O. Box 14155-6453, Tehran, I.R. Iran ³Department of Microbiology, Faculty of Veterinary Medicine, University of Lorstan, P.O. Box 465, Khorram-Abad, I.R. Iran

Abstract

Typhoid disease or salmonellosis is a common sickness in horses. In several epidemiological studies in hospitalized horses, several serotypes of *Salmonella* often are predominant in nosocomial infections. Transportation, overcrowding, dehydration, oral antimicrobial therapy and infections are the risk factors which may activate latent or subclinical salmonellosis. In this study, the occurrence of typhoid due to *Salmonella* serogroup B was considered in a Caspian ponies flock kept in a husbandry center of ponies around Tehran. During transportation of 19 ponies, two pregnant ponies aborted and four cases died because of acute septicemia. Pathological and bacteriological follow up showed salmonellosis. A multiplex polymerase chain reaction (m-PCR) assay was used for detection and identification of *Salmonella* to confirm pathological and bacteriological studies. *Salmonella* Typhimurium was isolated from bone marrow, mesenteric lymph nodes, liver and intestinal contents of died pony. *Salmonella* was not isolated from stools of other ponies. Pulsed Field Gel Electrophoresis (PFGE) and antibiotic susceptibility test were also performed. PFGE pattern was similar to the other collected isolates which have existed since more than 30 years ago in Iran. Because of importance of salmonellosis in ponies, using rapid methods are recommended to confirm the presence of *Salmonella*. Results showed that m-PCR permit to evaluate samples more rapidly than other methods and also can detect multiple genes simultaneously like virulence factors which declare virulence of the isolates and have surveillance signifi-

cance.

Keywords: Caspian Pony; *Salmonella* Typhimurium; Transportation; Multiplex PCR; PFGE

INTRODUCTION

In several epidemiological studies in hospitalized horses, several *Salmonella* serotypes such as *Salmonella* Typhimurium, *S. Newport*, *S. Anatum*, *S. Agona*, Heidelberg and *S. Ohio* have been commonly isolated and one or two serotypes often are predominant in nosocomial infections. *Salmonella enterica* serovar Typhimurium is the most frequently isolated serovar worldwide and is considered as one of the most virulent serotypes, affecting horses of all ages. (Gay, 1995; Walker *et al.*, 1995; Mainar-Jaime *et al.*, 1998; Ernst *et al.*, 2004). Typhoid disease or salmonellosis is a common disease of hospitalized horses. The spectrum of disease associated with *Salmonella* infections ranges from fecal shedding without clinical signs to septic shock and death. Many of the risk factors for salmonellosis in horses have been elucidated through descriptive studies of outbreaks on breeding farms and in hospitalized horses. In addition, a limited number of case-control studies have compared affected horses with unaffected horses to investigate potential risk factors. Stress factors that have been associated with salmonellosis in horses include transportation, antimicrobial administration, intestinal surgery, changes in diet, parturition, anesthesia and anthelmintic treatment (House *et al.*, 1999).

*Correspondence to: Taghi Zahraei Salehi, Ph.D.
Tel: +98 21 61117052; Fax: +98 21 66427517
E-mail: tsalehi@ut.ac.ir

In addition, dosage of bacteria and virulence are important factors in incidence of disease. Diagnosis and isolation of bacteria is very difficult in subclinical cases because of a few and or alternative fecal shedding. For laboratorial diagnosis, the amount of bacteria must be at least 100 particles/gr. However, not all patients shedding *Salmonella* may be detected by bacterial culture of fecal samples. Nowadays, advanced PCR based on oligonucleotide primers called m-PCR has been developed. This technique is more quickly and sensitive than bacterial culture (Aabo *et al.*, 1993). The aim of this study was molecular characterization of *Salmonella enterica* serovar Typhimurium isolated from Caspian ponies.

MATERIALS AND METHODS

Samples: In the present study, during transportation of 19 Caspian ponies, two pregnant ponies aborted and four cases died because of acute septicemia. Samples were obtained in the pathology laboratory of veterinary faculty and included intestinal content, mesenteric lymph node, liver and bone marrow which were collected from aborted and died Caspian ponies and feces from other live ponies. Samples were placed into plastic bags with ice pack and quickly transported to the Department of Microbiology, Faculty of Veterinary Medicine, University of Tehran. The Samples were inoculated into selenite-cystein broth (Merck, Darmstadt, Germany) for overnight enrichment at 37°C, and then plated on MacConkey agar (Merck) for primary selection. Presumptive *Salmonella* isolates were confirmed using conventional biochemical tests [triple sugar iron (TSI), urease test, MR-VP, Indole product and citrate utilization test] and serological

agglutination (Bacto-*Salmonella* O and H antisera; Difco™; Becton Dickinson and Company, Franklin Lakes, MI, USA).

Oligonucleotide primers: For m-PCR assay, four primer sets were selected. ST139 and ST141, specific for *Salmonella* genus (Rahn *et al.*, 1992) and the RfbJ, FliC and FljB, specific for the *rfbJ*, *fliC* and *fljB* genes of *Salmonella* Typhimurium or other *Salmonella* serovars with similar antigenic properties (Lim *et al.*, 2003). ST139 and ST141 primers are specific to detect *InvA* gene (one of *Salmonella* virulence genes which indicate to “invasion”). Also RfbJ, FliC and FljB primers are specific to detect O:4 (one of *Salmonella* LPS genes), H:i and H:1,2 (two of *Salmonella* flagella genes which indicate to motility), respectively. Primer sets, *inv-A*, RfbJ, FljB and FliC chosen, have been successful to detect *Salmonella* Typhimurium (Zahraei Salehi *et al.*, 2007). The primers sequences and their corresponding genes are shown in Table 1.

DNA extraction and amplification: A single colony of each isolate on agar plate was picked and suspended in 200 µl of distilled water. After vortexing, the suspension was boiled for 5 min, and 50 µl of the supernatant was collected after centrifuging for 10 min at 14,000 rpm. Polymerase chain reaction was performed with 10 µl of DNA sample, 5 mM KCl, 1.5 mM MgCl₂, 10 mM Tris-HCl pH 8.5, 1 µM of each primer, 200 µM dNTPs (Fermentas, Latvia) and 1 U of Taq DNA polymerase (Fermentas, Latvia) in a final volume of 25 µl. Amplifications were performed in a DNA thermocycler (Techne, TC-512, Cambridge, UK). The m-PCR protocol consisted of the following steps: The initial denaturation step of 5 min at 95°C; 30 cycles, with considering of 1 min at 95°C, 1 min at

Table 1. Primers sets used for detection of *Salmonella* serovars and identification of *Salmonella* Typhimurium.

Primers	Target gene	Amplicon (bp)	Reference
ST139-s: 5'-GTGAAATTATCGCCACGTTCCGGCAA-3' ST141-as: 5'-TCATCGCACCGTCAAAGGGAACC-3'	<i>invA</i>	284	Rahn <i>et al.</i> , 1992
Rfbj-s: 5'-CCAGCACCAAGTTCCAAGTTGATAC-3' Rfbj-as: 5'-GGCTTCCGGCTTTATTGGTAAGCA-3'	<i>rfbJ</i>	663	Lim <i>et al.</i> , 2003
Flic-s: 5'-ATAGCCATCTTTACCAGTTCCCCC-3' Flic-as: 5'-GCTGCAACTGTTACAGGATATGCC-3'	<i>fliC</i>	183	Lim <i>et al.</i> , 2003
Fljb-s: 5'-ACGAATGGTACGGCTTCTGTAACC-3' Fljb-as: 5'-TACCGTCGATAGTAACGACTTCGG-3'	<i>fljB</i>	526	Lim <i>et al.</i> , 2003

65°C and 30 s 72°C; and a final extension step of 7 min at 72°C. The PCR products were subjected to electrophoresis in 1.2% (w/v) agarose gel, stained with ethidium bromide and photographed under UV transilluminator. In each PCR run, a negative (distilled water) and a positive (*Salmonella* Typhimurium ATCC14028) control tubes were included (Zahraei Salehi *et al.*, 2007).

Pulsed Field Gel Electrophoresis (PFGE): Pulsed-field gel electrophoresis was performed according to the procedures developed by the Centers for disease Control and Prevention (CDC) for molecular subtyping of *Escherichia coli* O157:H7, non-typhoidal *Salmonella* serovars and *Shigella sonnei* and as previously described (Centers for Disease Control and Prevention, 2004). Briefly, agarose-embedded DNA was digested overnight in a water bath at 37°C with 50 U of *Xba*I (Fermentas, Latvia). The restriction fragments were separated by electrophoresis in 0.5X Tris-borate-EDTA (TBE) buffer at 14°C for 20 h in 6 V/Cm using a CHEFF DR II electrophoresis system (Gene Navigator, Pharmacia, Sweden) with pulse times of 2.2 to 63.8 s. The gels were stained with ethidium bromide (1 µg/ml) and destained with the buffer remained in the electrophoresis apparatus for 60-90 min and then images were captured with trans illuminator. Also isolates presenting DNA smear patterns were retested. The size standard used for all gels was *Xba*I-digested DNA from *Salmonella* Braenderup strain H9812

(ATCC no BAA-664), the universal size standard used by all PulseNet laboratories.

Antibiotic susceptibility: Antibiotic susceptibility test was performed by the standard disk diffusion method in Mueller-Hinton agar, and the results were interpreted in accordance to the criteria of the National Committee for Clinical Laboratory Standards (National Committee for Clinical Laboratory Standards, 2001). The strain was screened for resistance to the following antibiotics: cephalexin (CN, 30 µg), oxytetracycline (T, 30 µg), trimethoprim (TMP, 5 µg), lincospectin (LP, lincomycin/spectinomycin 15/200), enrofloxacin (NFX, 5 µg), trimethoprim sulfamethoxazole (SXT), nalidixic acid (NA, 30 µg), nitrofurantoin (FM, 300 µg), ampicillin (AM,10 µg), chloramphenicol (C: 30 µg), kanamycin (K, 30 µg), streptomycin (S, 10 µg) and ceftiofur (CFTIO, 30 µg) (Bahar afshan Co, Tehran, Iran).

RESULTS

Isolation and identification of *Salmonella* Typhimurium: Cultured samples were positive for *Salmonella enterica* serovar Typhimurium. Also the




Figure 1. Multiplex PCR result of some serotype of *Salmonella* for confirming of *S. Typhimurium*. 183 bp: *fliC*; 284 bp: *invA*; 526bp: *fljB*; 663bp: *rfbJ*; M: 100 bp DNA ladder (Fermentas, Latvia); C: Positive control (*S. Typhimurium* ATCC14028); 1: *S. Typhimurium* isolated from pony; 2: *S. Enteritidis* (wild type); 3: *S. Paratyphi B* (wild type); 4: *S. Typhimurium* (wild type); 5: *E. coli* (wild type); 6: negative control. Numbers have been explained on the basis of bp.




Figure 2. PFGE by *Xba*I enzyme digestion of some *S. Typhimurium* isolates from: 1: pony 2: cat 3,4,7,8: chicken 5,6: cow 9: *S. Typhimurium* with ATCC 14028 and M: *S. Braenderup* H9812 Marker (*Salmonella* PFGE marker according to PulseNet protocol).

samples were positive with serotyping (1,4,5,12:i:1,2). Four amplified product (663, 526, 284 and 183 bp) were found in all specimens that had serovar Typhimurium, they corresponded to the *rfbJ*, *fljB*, *invA* and *fliC* genes of this serovar respectively. In *Salmonella enterica* serovar Enteritidis (1, 9, 12; g, m:-) only one PCR product (284 bp) was amplified from the *invA* gene. In *Salmonella enterica* serovar Paratyphi B (1,4,5,12:b:1,2) three positive bands (284, 526 and 663 bp) were amplified corresponding to the *invA*, *fljB* and *rfbJ* genes respectively (Fig. 1). Eleven bands were presented in PFGE pattern of the isolate. Bands sizes were almost 40, 70, 90, 230, 260, 300, 380, 550, 670, 730 and 780 kb. This pattern was like to the other PFGE patterns isolated from cat (2006; 1979), sparrow (2005) and parrot (2005) in Iran which their information were documented in surveillance system of our laboratory. Comparison between PFGE pattern of the isolate with the standard size and *S. Typhimurium* with ATCC 14028 was presented in Figure 2.

Antibiotic susceptibility: The isolate presented resistance against cephalosporin, oxytetracycline and streptomycin but were sensitive to oxytetracycline (T), trimethoprim (TMP), lincospectin (LP, lincomycin/spectinomycin), enrofloxacin (NFX), trimethoprim sulfamethoxazole (SXT), nalidixic acid (NA), ampicillin (AM), chloramphenicol (C), kanamycin (K) and ceftiofur (CFTIO).

DISCUSSION

Salmonellosis is a commonly encountered infectious disease of horses (Walker *et al.*, 1995). *Salmonella* are among the most frequent causes of acute diarrhea in horses and the incidence seems to be increasing (van Duijkeran *et al.*, 1995). *Salmonella* Typhimurium is the most common serotype identified and is considered as one of the most virulent serotypes affecting horses of all ages. A number of other serotypes with apparent varying degrees of virulence have also been reported to cause salmonellosis in the horses such as *S. Enteritidis*, *S. Newport*, *S. Anatum*, *S. Java*, *S. Saintpaul*, *S. Kerfeld*, *S. Thompson*, *S. Heidelberg*, *S. Hadar*, *S. Infantis*, *S. Derby*, *S. Oranienburg*, *S. Hindmarsh* (Ernst *et al.*, 2004; van Duijkeran *et al.*, 2002; Mainar-Jaime *et al.*, 1998; Daniel *et al.*, 1997; van-Duijkeran *et al.*, 1995; Walker *et al.*, 1995; Traub-Dargatz *et al.*, 1990). Recently, the multiresistant *S.*

Typhimurium phage type DT104 was the most common phage type isolated from horses correspond with those found in human, pigs and cattle and have highly zoonotic significance (van Duijkeren *et al.*, 2002).

The predominance of one or two *Salmonella* serotypes in herd and hospital of horses in the most outbreaks of salmonellosis, suggests that many of these outbreaks reflect nosocomial infections. In addition, some case-control studies have compared affected with control horses to investigate potential risk factors. Stress factors that have been associated with salmonellosis in horses include transportation, antimicrobial administration, intestinal surgery, changes in diet, food deprivation, dehydration, colic, gastrointestinal tract disease, anesthesia and anthelmintic treatment (Ernst *et al.*, 2004; House *et al.*, 1999; Mainar-Jaime *et al.*, 1998).

Owen *et al.* 1983 reported that transportation has a major role in reactivating the *Salmonella* infection in ponies. Diarrhea due to a reactivation of the *Salmonella* infection occurred greater than 3 days after stress, although maximal shedding of organisms occurred within 24 h (Owen *et al.*, 1983).

In this study, it seems that transportation of ponies, changing in diet and probably deprivation during transportation, have been the most important risk factors and in accordance to predominance of *S. Typhimurium*, the outbreak has had nosocomial identity. Following of source of the infection in water supplies and environments of the herd, *Salmonella* was not detected. There were not any food sources for examination but it seems that only food materials could be the source of initial infection. PFGE pattern of the isolate revealed similarity with the other collected isolates which have existed since more than 30 years ago in Iran (unpublished data). This similarity indicates that the strain probably present for several years in this region. So, according to permanent and reliable profiles generated in PFGE method, we concluded and confirmed that main reason of such epidemic with systemic property in pony, is stress factors such as transportation, changing in diet and probably deprivation during transportation that introduce optimum conditions for *Salmonella* invasion or reactivation of latent *Salmonella* in carriers.

Antibiotic susceptibility test indicated that the isolate does not have any noticeable antibiotic resistance pattern. So, it seems that pathogenesis of this strain is related to risk factors or susceptibility of pony (as host) to *S. Typhimurium*.

The greatest difficulty in identifying of the out-

breaks is detection of the first source of infection for determination and separation of nosocomial infection with one source from those acquired prior to admission (House *et al.*, 1999). Therefore, sensitivity and rapidity of the methods which are applied for *Salmonella* detection are important. The limit of detection for most culture techniques is around 100 *Salmonella* organism/g of the feces. Subclinical infections in horses are more difficult to detect because less organism are shed and shedding may be intermittent (Mainar-Jaime *et al.*, 1998).

Advanced PCR based techniques, such as multiplex PCR, are logically the most sensitive methods which could be applied for detection of even one organism/g of feces. Amavisit *et al.* (2001) indicated, while the sensitivity of the PCR assay was less than culture of feces for *Salmonella*, its sensitivity on fecal samples obtained from horses, was much greater than culture method. They detected *Salmonella* DNA in 40% of fecal samples using the PCR assay while *Salmonella* was isolated by culturing from only 2% of the samples. Also, this method performs in less than 5-7 h and actually decreases the time of identification (Amavisit *et al.*, 2001). Due to correlation between *inv-A* virulent gene and clinical signs such as fever, bloody diarrhea and prolapsed rectum in ponies, it seems m-PCR by presenting multiple virulent genes (Such as *inv-A*), confirms the virulent properties of an isolate and is useful for molecular characterization of the isolate which causes an epidemic.

Therefore, in accordance to low prevalence of *Salmonella* shedding in horses about 1.7%, low drainage of *Salmonella* bacteria from feces (Mainar-Jaime *et al.*, 1998), virulence and zoonotic importance of *Salmonella* and for molecular characterization of an isolate which causes *S. Typhimurium* epidemics, we need powerful method to detect it in *equidae*. M-PCR is suggested to be employed as a rapid and sensitive method for identifying the source of *Salmonella* especially in outbreaks of *S. Typhimurium* and even in carrier status in *equidae*.

References

- Aabo S, Rasmussen OR, Sorensen PD, Olson JE (1993). *Salmonella* identification by polymerase chain reaction. *Mol Cell Probes*. 7: 171-178.
- Amavisit P, Browning GF, Lightfoot D, Church S, Anderson GA, Whithear KG, Markham PF (2001). Rapid PCR detection of *Salmonella* in horse faecal samples. *Vet Microbiol*. 79: 63-74.
- Centers for Disease Control and Prevention (2004). One day (24-28 h) standardized laboratory protocol for molecular subtyping of *Escherichia coli* O157:H7, non-typhoidal *Salmonella* serotypes, and *Shigella sonnei* by pulsed field gel electrophoresis (PFGE). Pulse Net, USA.
- Clinical Laboratory Standard Institute (2001). Performance standards for antimicrobial susceptibility testing; 11th informational supplement, vol. 21, no 1. M100-S11 NCCLS, Wayne, PA.
- Daniel RG, Barrow P, Szmolleny G, Wood M (1997). *Salmonella* Hindmarsh infection in sheep and ponies. *Vet Res*. 141: 203-203.
- Ernst NS, Hernandez JA, MacKay RJ, Brown MP, Gaskin JM, Nguyen AD, Giguere S, Colahan PT, Troedsson MR, Haines GR, Addison IR, Miller BJ (2004). Risk factors associated with fecal *Salmonella* shedding among hospitalized horses with signs of gastrointestinal tract disease. *JAVMA*. 225: 275-281.
- Gay J (1995). Bovine herds salmonellosis. *Am. J Infect Control*. 23: 56-62.
- House JK, Mainar-Jaime RC, Smith BP, House A, Kamiya DY (1999). Risk factors for nosocomial *Salmonella* infection among hospitalized horses. *J Am Vet Med Assoc*. 214: 1511-1516.
- Lim YH, Hirose K, Izumiya H, Arakawa E, Takahashi H, Terajima J, Itoh K, Tamura K, Kim S, Watanabe H(2003). Multiplex polymerase chain reaction assay for selective detection of *Salmonella enterica* serovar Typhimurium. *Jpn J Infect Dis*. 56: 151-155.
- Mainar-Jaime RC, House JK, Smith BP, Hird DW, House AM, Kamiya DY (1998). Influence of fecal shedding of *Salmonella* organisms on mortality in hospitalized horses. *JAVMA*. 213: 1162-1166.
- Owen RA, Barnum DA (1983). Effects of transportation, surgery, and antibiotic therapy in ponies infected with *Salmonella*. *Am J Vet Res*. 44: 46-50.
- Rahn K, De Grandees SA, Clarke RC, Galen JE, Pinocchio C, Curtis III R, Giles CL (1992). Amplification of an *inv-A* sequence of *Salmonella* Typhimurium by polymerase chain reaction as a specific method of detection of *Salmonella*. *Mol Cell Probes*. 6: 271-279.
- Traub-Dargatz JL, Salman MD, Jones RL (1990). Epidemiologic study of *salmonellae* shedding in the feces of horses and potential risk factors for development of the infection in hospitalized horses. *JAVMA*. 196: 1617-1622.
- van-Duijkeran E, van-Klingeren B, Vulto AG, Marianne M, Sloet OO, Breukink HJ, van Miert ASJPAM (1995). In vitro susceptibility to antimicrobial drugs of 62 *Salmonella* strains from horses in The Netherlands. *Epidemiol Infect*. 45: 19-26.
- van Duijkeren E, Wannet WJB, Heck MEOC, van Pelt W, van Oldruitenborgh-Oosterbaan MMS, Smit JAH, Houwers DJ (2002). Sero types, phage types and antibiotic susceptibilities of *Salmonella* strains isolated from horses in The Netherlands from 1993 to 2000. *Vet Microbiol*. 86: 203-212.
- Walker RL, de Peralta TL, Villanueva MR, Snipes KP, Madigan JE, Hird DW, Kasten RW (1995). Genotypic and phenotypic analysis of *Salmonella* strains associated with an outbreak of equine neonatal salmonellosis. *Vet Microbiol*. 43: 143-50.
- Zahraei Salehi T, Tadjbakhsh H, Atashparvar N, Nadalian MG, Mahzounieh MR (2007). Detection and identification of *Salmonella Typhimurium* in bovine diarrhoeic fecal samples by immunomagnetic separation and multiplex PCR assay. *Zoonoses Public Health*. 54: 231-236.