JOURNAL OF ENTOMOLOGICAL SOCIETY OF IRAN 2017, 37(2), 155–168



نامه انجمن حشره شناسی ایران ۱۹۸ – ۱۳۹ (۲), ۱۳۹۹

Biochemical characterization of digestive proteases of *Pieris brassicae* (Lepidoptera: Pieridae) feeding on cruciferous hosts

Fariba Mehrkhou

Department of Plant Protection, Faculty of Agriculture, Urmia University, Urmia, Iran. * Corresponding author, E-mail: f.mehrkhou@urmia.ac.ir

Abstract

Pieris brassicae is one of the most serious pests of cultivated crucifers in the world. The present experiment is intended to determine the proteolytic digestion of fifth instar larvae and the effects of four cruciferous hosts to *P. brassicae* nutritional physiology. *Brassica oleracea var. botrytis, Brassica oleracea var. capitata, Brassica oleracea var. italica* and *Brassica oleracea var. viridis* were assessed under laboratory conditions at $25 \pm 2^{\circ}$ C, $60 \pm 5\%$ RH and a photoperiod of 16:8 (L: D) h. The maximum total proteolytic activity in the midgut extract was observed at pH 11, suggesting the presence of serine protease. The lowest rate of tryptic activity and both the highest general proteolytic (0.11 ± 0.01 Umg⁻¹) and chymotryptic activities (0.45±0.002 Umg⁻¹) were observed in *B. oleracea var. viridis*. The proteolytic activity of *P. brassicae* on this host plant indicates that *B. oleracea var. viridis* is not a suitable host for the pest likely due to the presence of some secondary chemicals or proteinaceous protease inhibitors.

Keywords: Pieris brassicae, Digestive enzymes, Proteolytic activity, Cole crops

ویژگیهای بیوشیمیایی پروتئازهای گوارشی پروانه سفیده بزرگ کلم Pieris brassicae (Lepidoptera: Pieridae) با تغذیه از میزبانهای مختلف خردلیان

فریبا مهرخو گروه گیاهپزشکی، دانشکده کشاورزی، دانشگاه ارومیه * مسئول مکاتبات، یست الکترونیکی: f.mehrkhou@urmia.ac.ir

چکیدہ

پروانه سفیده بزرگ کلم (Pieris brassicae (Lepidoptera: Pieridae یکی از مهمترین آفات مخرب خردلیان در اغلب کشورهای جهان می باشد. هدف از این تحقیق، تعیین آنزیمهای پروتئولیتیک لارو سن پنجم سفیده بزرگ کلم و تأثیر چهار و تم مختلف کلم بر فیزیولوژی تغذیه آن می باشد. در این تحقیق ارقام مختلف کلم شامل کلم گل، کلم برگی سفید، بروکلی و کلم برگی قرمز استفاده شد. آزمایشات در آزمایشگاه، تحت شرایط دمایی ۲°۲ ± ۲۵، رطوبت نسبی ۵ ± ۲۰ درصد و دوره نوری ۱۲ ساعت روشنایی و ۸ ساعت تاریکی انجام گرفت. بیشترین میزان فعالیت پروتئولیتیک عصاره روده میانی کلم قرمز بود، در حالی که بیشترین میزان فعالیت تریپروتئازی می باشد. کمترین میزان فعالیت تریپتیکی در کیم قرمز بود، در حالی که بیشترین میزان فعالیت ویژه پروتئولیتیکی (۱۰/۰ ± ۱۱/۱ واحد آنزیمی/ میلی گرم) و پروتئولیتیکی (۲۰۱۰ ± ٤۵/ واحد آنزیمی/ میلی گرم) در لاروهای تغذیه شده از کلم قرمز مشاهده شد، فعالیت پروتئولیتیکی پروانه سفیده کلم روی میزبانهای مذکور بیانگر ناماسب بودن کلم قرمز در میان میزبانهای مورد مطالعه بود که دلیل آن می تواند احتمالا به وجود ترکیبات شیمای ثانویه و یا مهارکنده های پروتئازی موجود با این رقم مورد مطالعه بود که دلیل آن می تواند احتمالا به وجود ترکیبات شیمای ثانویه و یا مهارکنده های پروتئازی موجود با این رقم مورد مطالعه بود که دلیل آن می تواند احتمالا به وجود ترکیبات شیمایی ثانویه و یا مهارکنده های پروتئازی موجود با این رقم مرتبط باشد.

دریافت: ۱۳۹۵/۹/۱، یذیرش: ۱۳۹٦/۱/۲۳.

Introduction

It has been estimated that damage from insect pests alone causes more than 40% of yield loss on vegetable crops annually (Hasan & Ansari, 2011). Pajmon (1999) listed 38

insect pests of Cole crops, of which *Pieris brassicae* L. remains one of the most destructive onethrough developing stages (seedling, vegetative, and flowering) (Bhalla & Pawar, 1977; Sachan & Gangwar, 1980; Siraj, 1999; Lal & Ram, 2004; Blatt *et al.*, 2008; Bhandari *et al.*, 2009; Zibaee, 2012; Kumar *et al.*, 2015). This pest infests 83 species of food plants belonging to Cruciferae (Jainulabdeen & Prasad, 2004; Raqib, 2004). Garden nasturtium was used to feed the larvae of *P. brassicae* prolifically (Kumar *et al.*, 2015) on which a larva of *P. brassicae* consumed 74 - 80 cm² leaf area (Younas *et al.*, 2004).

Resistance to insecticides has been developed in *P. brassicae* (Sharma and Gupta, 2009). Modification of economically important crops by adding insecticidal proteins is one of the alternative methods, which can be used to control pests. The first step in designing inhibitor-transgenic crops is the understanding of biochemical characterization of the enzyme activities in the midgut of the target pest (Oppert, 2000; Oppert *et al.*, 2000; Wilhite *et al.*, 2000).

The herbivorous lepidopteran larvae feed actively on plants to gather nutritional components for their development and progression into the reproductive adult phase. Proteins serve as primary constituents in the development, growth and fecundity of insects and are digested by protease. Any disorder to digestion causes impair in insect development and fecundity of the adults. Srinivasan *et al.* (2005, 2006) reported that among proteins, serine proteases play an important role in digestion processes in lepidopteran larvae. These enzymes have been shown to have a high pH optimum of 8–11, which is consistent with the alkaline conditions in the midgut (Zibaee, 2012).

The biological (Hasan & Ansari, 2010a, b; Mehrkhou & Taheri-Sarhozaki, 2014) and nutritional indices (Mehrkhou *et al.*, 2013) of large white butterfly, on Cole crops have been previously studied. Zibaee (2012) investigated the enzyme diversity of *P. brassicae* on cabbage. Kumar *et al.* (2015) evaluated the midgut serine proteases and alternative host plant utilization in *P. brassicae*, to design sustainable pest management strategies, including transgenic approaches using genes encoding plant protease inhibitors.

The present study was conducted to identify types of proteases in the midgut of *P. brassicae* and their responses to different host plants to use the data in IPM program in order to improve the control strategies against this pest.

Material and methods

Host plants

Seeds of four Cole crops namely, *Brassica oleracea var. botrytis*, *Brassica oleracea var. capitata*, *Brassica oleracea var. viridis* and *Brassica oleracea var. italica* were grown in a research field at the Department of Crop Protection of Urmia University (Urmia, Iran).

Insect rearing and dissections

The eggs of *P. brassicae* were collected from the leaves of each host plant and kept in plastic rearing containers (14 cm width×20 cm length×8 cm height) covered with fine nylon mesh containing a piece of moist cotton for providing the ventilation and moisture, respectively. The neonate larvae were taken out and reared to their fifth instar. The experiments were conducted in growth chambers at temperature of $25\pm2^{\circ}$ C, $60\pm5\%$ R.H. and a photoperiod of 16: 8 h (L:D). For dissection, fifth instar larvae of *P. brassicae* were anaesthetized on ice and their midguts homogenized immediately in distilled water. The crude gut homogenate was transferred to 1.5 ml tubes and centrifuged at 13,000 rpm for 15 min at 4 °C. The clear supernatant was transferred to a pre-chilled Eppendorf tube. The samples were stored at -20° C until further use.

Protein quantification

The protein concentrations of gut digestive enzymes were determined according to the Bradford's method (1976). Bovine serum albumin (BSA; Bio-Rad, biorad.com) was used as a standard (0.1, 0.4, 0.7, 1.0, 1.2, 1.6 and 1.8 mg/ml).

Midgut's pH and caseinolytic activity

General proteolytic activity in the midgut of *P. brassicae* was measured by using 2% azocasein (Sigma-Aldrich, A2765) as a substrate at broad pH range (pH 5-12) of universal buffer (50 mM sodium acetate-phosphate-borate), according to Elpidina *et al.* (2001) with slight modification. The universal buffer (90 μ l), 2% azocasein solution (30 μ l) and enzyme solution (15 μ l) were used in reaction, which incubated in Benmary (at 37° C for 60 min), and trichloroacetic acid (TCA) 30% (30 μ l) was used to stop the reaction. Precipitation was achieved by cooling at 4° C for 60 min and the reaction mixture centrifuged at 12000 (×g) for 15 min. We added 50 μ l of 2 M NaOH to the supernatant (100 μ l) and the absorbance was recorded 440 nm. All assays were carried out in triplicate. The blank consisted of all mentioned portions except for the enzyme solution.

Specific proteolytic activity

Determining the optimum endoproteolytic activity was conducted at extent pH range of universal buffer on susceptible cultivar (*Brassica oleracea var. capitata*). The optimum pH was selected for determining of endoproteolytic activity on other host plants. Trypsinlike and chymotrypsin-like activities (as two sub-classes of serine proteases) were assayed using the concentration 1 mM of Na-benzoyl-L-arginine-p-nitroanilide hydrochloride (Sigma-Aldrich, 3279) and 1 mM of N-succinyl–alanine–alanine–proline–phenylalanine–pnitroanilide (Sigma-Adrich, S7388) as substrates, respectively. The reaction mixture included 85 μ l of universal buffer (50 mM, pH 5-12), 5 μ l of each mentioned substrate, and 10 μ l of the enzyme solution. All assays were carried out in triplicate along with appropriate blanks (Sarboland *et al.*, 2017).

Protease inhibitors assay

The effect of different inhibitors on proteolytic activities of the midgut was determined. The used inhibitors and their concentrations are as follows: Serine protease inhibitors, PMSF (phenylmethyl sulfonyl fluoride1,1 mM; Sigma-Aldrich, P7626); trypsin inhibitor, TLCK (N α -p-tosyl-L-lysine chloromethyl ketone, 0.1 mM; Sigma-Aldrich, T5012); chymotrypsin inhibitor, TPCK (N-tosyl-L-phenyl-alanine chloromethyl ketone, 0.1 mM; Sigma-Aldrich, T7254). To determine the effect of these compounds on enzyme activities, the enzymes were pre-incubated with the inhibitors at room temperature for 15 min, then the substrate was added before performing assays (Hosseininaveh *et al.*, 2007).

Statistical analysis

All data were obtained from a complete randomized design and compared with oneway analysis of variance (ANOVA) followed by Tukey's test once significant differences were determined at $P \le 0.05$ (SPSS, 2010).

Results

Midguts'pH

Optimum pH of general proteolytic activity was obtained for each host plant separately. General proteolytic activity of fifth instar larval gut extracts reared on four different Cole crops are shown in Fig. 1. Results of the study demonstrated the presence of serine proteases in the midgut of *P. brassicae* after feeding on different host plants. The optimal pH for general proteolytic activity in the midgut of *P. brassicae* was obtained over a broad alkaline pH range (pH 8 to pH 11), with maximum activity at pH 11 using azocasein (2%) as a substrate (Fig.1). In highly alkaline conditions (pH 11), over 80% of maximum activity was recorded.

Caseinolytic and endoproteolytic activity

Specific protease activity was assayed with synthetic substrates for trypsin-like (BApNA) and chymotrypsin-like (SAAPFpNA) enzymes at different pH (5-12) on *Brrasica oleracea var. capitata.* Results showed a continuous increase from pH 5 to 12. The substrates were hydrolyzed in alkaline conditions, with an activity peak recorded at pH 11, indicating presence of trypsin- and chymotrypsin-like proteinases (Fig. 2). Among the

different Cole crops (F=17.22; df=3,8; P<0.01), protease activity was the highest in the larvae fed on *B. oleracea var. viridis*, although it was not significantly different among other three host plants. A midgut serine proteinase value was estimated 0.057, 0.068, 0.11 and 0.045 Umg⁻¹ on *B. oleracea var. botrytis, capitata, viridis and italica*, respectively (Fig. 3a).

There was a significant difference among different host plants, in terms of trypsin (F=20.35; df= 3,8; P<0.01) and chymotrypsin (F= 40.82; df= 3,8; P<0.01) activity. Trypsin activity of larvae, which fed on *B. oleracea var. capitata* and *viridis*, was significantly higher than other host plants. A gut trypsin proteinase specific activity value was estimated 0.27, 0.42, 0.45 and 0.28 (U mg⁻¹) substrate hydrolyzed on *B. oleracea var. botrytis, capitata, viridis and italica*, respectively (Fig. 3b). Chymotrypsin activity (SAAPF*p*NA as substrate) was high on *B. oleracea var. botrytis*-fed larvae but not significantly different from larvae reared on the rest of Cole crops (Fig.3b). The chymotrypsin- like activity was estimated 0.77, 0.41, 0.62 and 0.30 (U mg⁻¹) on *B. oleracea var. botrytis, capitata, viridis and italica*, respectively.



Fig. 1. Effect of pH on *P. brassicae* larval gut protease activity. Midgut extracts were assayed for total proteolytic activity by using Azocasein substrate on different of host plants. Activities are expressed as a change in Elisa reader at 440 nM/min due to hydrolysis of substrate. Each point represents mean of three independent estimations associated with standard errors.

Effect of protease inhibitors on proteolytic activity

General proteolytic activity of larval midgut extract was further characterized using general (PMSF) and specific (TLCK, TPCK) proteinase inhibitors (Table 1). Significant inhibition of protease activity by PMSF, TLCK, and TPCK suggested that serine proteinases were mainly responsible for protein digestion in the midgut. The highest inhibition by PMSF and TLCK were observed on *B. oleracea var. viridis* (Table 1).

 Table 1. The effects of some protease inhibitors, on the general proteolytic activity from midgut extract of *Pieris brassicae* larvae using azocasein as substrate, on different Cole crops.

		% Inhibition	
Cole crops	PMSF (1mM)	TLCK (0.1mM)	TPCK (0.1mM)
B. O. Var. Viridis	29.13 ± 1.30^{a}	57.58 ± 1.39^{a}	22.38 ± 4.32^{b}
B. O. Var. Capitata	24.83 ± 1.29^{ab}	46.18 ± 1.23^{ab}	38.51 ± 2.20^{b}
B. O. Var. Botrytis	20.23 ± 1.46^{b}	23.33 ± 0.06^{b}	$49.26\pm4.55^{\rm a}$
B. O. Var. Italica	21.88 ± 1.76^{b}	34.41 ± 1.75^{b}	43.65 ± 3.22^{ab}

Means with different letters in each column are significantly different at P < 0.01 based on Tukey's HSD multiple range test.

Discussion

Some biochemical process including regulating enzymatic reactions related to digestion, control the solubility and toxicity of stomach poisons, and determine gut flora caused by gut pH (House, 1974; Berenbaum, 1980; Terra & Ferreira, 1994).

Screening of the pH-activity profile of digestive peptidases in the midgut homogenate of the larval *P. brassicae* was accomplished under different pH conditions by azocaseine, at a wide range of pH (5-12). The results show one peak of activity, at alkaline (pH 11). It is likely due to the presence of serine protease activity. Further biochemical characterization of *P. brassicae* larval digestive proteases using general, specific substrates and synthetic inhibitors reveal that serine proteinases are dominant proteases, as is typical in other lepidopteran larvae (Johnston *et al.*, 1991; Gatehouse *et al.*, 1997; Chougule *et al.*, 2008). Similar findings have been reported for other lepidopteran species: 8-11 in *Ectomyelois ceratoniae* Zeller (Pyralidae) (Razavi- Tabatabaei *et al.*, 2011), 12 in *Helicoverpa armigera* Hübner (Naseri *et al.*, 2010) although, Rahimi-Namin *et al.* (2011) found that, the optimal proteolytic activity in the midgut of *P. brassicae* occurs at pH 8.

Larvae of *P. brassicae* exhibit complex and diverse forms of proteolytic digestion that is influenced by their host plants. Within different Cole crops, the highest general proteolytic activity was in the larvae reared on *B. oleracea var. viridis*, likely due to the hyperproduction of proteases by midgut cells of *P. brassicae* in response to protease inhibition by PIs. Zibaee (2012) stated higher proteolytic activity in the midgut of fifth instar larvae of *P. brassicae* (36.62 U mg⁻¹) by Hemoglobine. This differences can be related to the type of general substrate (Sarboland *et al.*, 2017).



Fig. 2. Effect of pH on *P. brassicae* larval gut, which were assayed for endoproteolityic activity by using BApNA and SAAPFpNA substrates on *Brassica Oleracea. var. botrytis.* Activities are expressed as a change in Elisa reader at 405 nM/min due to hydrolysis of substrates on cauliflower. Each point represents mean of three independent estimations associated with standard errors.

The presence of trypsin-like serine proteases in the larval midgut of *P. brassicae* was clearly demonstrated by the hydrolysis of the trypsin-specific substrate BApNA and inhibition of general proteolytic activity by TLCK. Similarly, chymotrypsin-like were also demonstrated by the hydrolysis of SAAPFpNA and by the general proteolytic inhibition by the inhibitor TPCK. Within Cole crops, the trypsin activity in the larvae reared on B. oleracea var. botrytis and italica was the lowest. It can be resulted from the inhibition of tryptic activity by PIs of these two host plants that decrease the activity of trypsin-like enzymes in midgut extracts of the larvae fed on these crops. However, the larvae fed on B. oleracea var. botrytis had the highest chymotryptic activity compared with the other Cole crops. The biochemical analysis of P. brassicae larval digestive enzymes using specific substrates showed that serine proteinases (trypsin and chymotrypsin) had the highest activity (Gatehouse et al., 1999; Hegedus et al., 2003; Chougule et al., 2008). Broadway (1989) found the highest activity of tryspin and chymotrypsin-like proteases in the midgut of P. rapae L.. Zibaee (2012) observed higher tryptic activity compared with chymotryptic in *P. brassicae*, which fed on cabbage leaves. It has been proven that insects that are reared on different host plants, haved different digestive enzymatic responses (Saikia et al., 2011).

Some protease inhibitor compounds were used for further proof of the proteolytic profile in the midgut of *P. brassicae*, which affecting serine proteases, significantly changed proteolytic activity, confirming the presence of serine proteases in the midgut extract of *P. brassicae*. PMSF, as a serine protease inhibitor, significantly decreased proteolytic activity. Although the highest general proteolytic activity was observed on *B. oleracea var. viridis*, but the lowest chymotryptic activity was possibly related to the presence of some serine protease inhibitors in this plant, resulting in hyperproduction of trypsin-like enzymes in response to the inhibition of these enzymes.



Fig. 3. Specific serin protease (a), Trypsin- (b) and chymotrypsin- -like (b) activities of midgut extracts from *Pieris brassicae* larvae reared on different brassicaceous Cole crops using Azocasin, BApNA and SAAPFpNA (pH 11) as substrates respectively. Bars represent means of three independent estimations associated with standard error.

The effects of examined different Cole crops on biological, life table and nutritional indices of large white butterfly were studied by Mehrkhou *et al.* (2013) and Mehrkhou and Taheri- Sarhouzaki (2014). The larvae which were fed on cabbage (*B. oleracea var. capitata*) had the lowest rate of mortality and the shortest developmental time, which was consistent with Hasan & Ansari (2010 a,b) who had found the cabbage the most suitable host plant comparing to mustard, radish and cauliflower. *B. oleracea var. viridis* was found to be unsuitable because of the highest rate of mortality and the longest developmental time.

The eco-physiological parameters show that the highest enzymetic activity was observed in the larvae which were fed on *B. oleracea var. viridis*, although their biological and life table parameters were at the lowest rate. Some insects exhibit an interesting flexibility in adapting to various host plants by altering the specificities of their gut proteases in response to qualitative changes in dietary protein content, esp. when the existing proteases are ineffective and/or inefficient for digestion (Gatehouse *et al.*, 1997). Mechanisms of insect resistance to PIs include the upregulation of enzymes that degrade the PIs (Yang*et al.*, 2009), the induction of enzymes that resist inactivation by PIs

(Broadway, 1996), and overproduction of enzymes to maintain normal levels of gut proteolysis (Brioschi *et al.*, 2007).

B. oleracea var. viridis was the least suitable species for inflicting longer development time, high rates of mortality, and the lowest chymotryptic activity. The larvae which fed on *B. oleracea var. capitata* were heavier and showed higher relative growth rate, shorter developmental time, as well as higher survival rate and tryptic activity. Naseri *et al.* (2010) studied the ecophysiologica aspects of *H. armigera*, on different soybean varieties and mentioned that the larvae which were reared on L17 variety of soybean had both the longest and the lowest biological and demographic parameters while their enzymetic activities were at the highest levels. They argued that the larvae had to increase enzymetic activity by hyperproduction of chymotrypsin- and elastase-like enzymes in response to the inhibition of these enzymes.

Generally, digestive enzymatic studies of insects in response to feeding on different host plants have different applications such as assessment of resistance and susceptibility of host plants (Naseri *et al.*, 2010; Saadat *et al.*, 2014; Sarboland *et al.*, 2017), identifying of resistance factor and antifeedant compounds (Lewis *et al.*, 1997), production of crops with elevated levels of endogenous resistance by plant proteinasese inhibitors and glucosinolates (Ryan, 1990; Hosseininaveh *et al.*, 2007; Velasco *et al.*, 2008).

Host plants have critical roles in the fitness of herbivorous insects; it is due to the fact that, the host components could directly affect their performance via influence on the digestive process (Chown and Nicolson, 2004). Changes of proteolytic activity of different host plants and genetically modified varieties have been reported in different insects (Chen, 2008). Ecological parameters especially population growth rate, food quality, presence of secondary components (e.g. glucosinolates) and possible inhibitors from a wide range of Cole crops are substantial for designing stable planting system.

Acknowledgments

We are grateful to Urmia University, office of Vice Chanceller, for funding this research.

References

- **Berenbaum, M.** (1980) Adaptive significance of midgut pH in larval lepidoptera. *The American Naturalist* 115(1), 138–146.
- Bhalla. O. P. & Pawar, A. D. (1977) A survey study of insect and non-insect pests of economic Importance in Himachal Pradesh. Tiku and Tiku, Kitab Mahal, Bombay, p. 80.

- Bhandari, K., Sood, P., Mehta, P. K., Choudhary, A. & Prabhakar, C. S. (2009) Effect of botanical extracts on the biological activity of *Granulosis virus* against *Pieris brassicae*. *Phytoparasitica* 37, 317–322.
- Blatt, S. E., Smallegange, R. C., Hess, L., Harvey, J. A., Dicke, M. & van Loon, J. J. A. (2008) Tolerance of *Brassica nigra* to *Pieris brassicae* herbivory. *Botany*, 86, 641– 648.
- Brioschi, D., Nadalini, L. D., Bengtson, M. H., Sogayar, M., Moura, D. S. & Silva-Filho, M. C. (2007) General up regulation of *Spodoptera frugiperda* trypsins and chymotrypsins allows its adaptation to soybean proteinase inhibitor. *Insect Biochemistry and Molecular Biology* 37, 1283–1290.
- Broadway, R. M. (1989) Characterization and ecological implications of midgut proteolytic activity in larval *Pieris rapae* and *Tricoplusia ni. Journal of Chemical Ecology* 15, 2101–2113.
- Broadway, R. M. (1996) Dietary proteinase inhibitors alter complement of midgut proteases. Archives of Insect Biochemistry and Physiology 32, 39–53.
- Chen, M. S. (2008) Inducible direct plant defense against insect herbivores: a review. *Journal of Insect Science* 15, 101–114.
- Chougule, N. P., Doyle, E., Fitches, E. & Gatehouse, J. A. (2008) Biochemical characterization of midgut digestive proteases from *Mamestra brassicae* (cabbage moth; Lepidoptera: Noctuidae) and effect of soybean Kunitz inhibitor (SKTI) in feeding assays. *Journal of Insect Physiology* 54, 563–572.
- Chown, S. L. & Nicolson, S. W. (2004) Insect Physiological Ecology Mechanisms and Patterns. Oxford University Press, p 254.
- Elpidina, E. N., Vinokurov, K. S., Gromenko, V. A., Rudenshaya, Y. A., Dunaevsky, Y. E. & Zhuzhikov, D. P. (2001) Compartmentalization of proteinases and amylases in *Nauphoeta cinerea* midgut. *Archives of Insect Biochemistry and Physiology* 48, 206–216.
- Gatehouse, A. M. R., Davidson, G. M., Newwll, C. A., Merryweather, A., Hamilton,
 W. D. O., Burgess, E. P. J., Gilbert, R. J. C. & Gatehouse, J. A. (1997) Trangenic potato plants with enhanced resistance to the tomato moth, *Lacanobia oleracea*: growth room trials. *Molecular Breeding* 3, 49–63.
- Gatehouse, L. N., Shannon, A. L., Burgess, E. P. J. & Christeller, J. T. (1997). Characterization of major midgut proteinase cDNAs from *Helicoverpa armigera* larvae and changes in gene expression in response to four proteinase inhibitors in the diet. *Insect Biochemistry and Molecular Biology* 27, 929-944.
- Gatehouse, A. M. R., Norton, E., Davison, G. M., Babbe, S. M., Newell, C. A. & Gatehouse, J. A. (1999) Digestive proteolytic activity in larvae of tomato moth,

Lacanobia oleracea; Effects of plant proteinase inhibitors in vitro and in vivo. *Journal* of *Insect Physiology* 45, 545–558.

- Hasan, F. & Ansari, M. S. (2010a) Effect of different cole crops on the biological parameters of *Pieris brassicae* (L.) (Lepidoptera: Pieridae) under laboratory conditions. *Journal of Crop Science and Biotechnology* 13, 195–202.
- Hasan, F. & Ansari, M. S. (2010b) Population growth of *Pieris brassicae* (L.) (Lepidoptera: Pieridae) on different cole crops under laboratory conditions. *Journal of Pest Science* 84, 179–186.
- Hasan, F. & Ansari, M. S. (2011) Toxic effects of neem-based insecticides on *Pieris brassicae* (Linn.). *Crop Protection* 30, 502–507.
- Hegedus, D., Baldwin, D., O'Grady, M., Braun, L., Gleddie, S., Sharpe, A., Lydiate, D. & Erlandson, M. (2003). Midgut proteases from *Mamestra configurata* (Lepidoptera: Noctuidae) larvae: Characterization, cDNA cloning and expressed sequence tag analysis. *Archives of Insect Biochemistry and Physiology*, 5330–47.
- Hemming, J. D. C. & Lindroth, R. L. (2000) Effects of phenolic glycosides and protein on gypsy moth (Lepidoptera: Lymantriidae) and forest tent caterpillar (Lepidoptera: Lasiocampidae) performance and detoxication activities. *Environmental Entomology* 29, 1108–1115.
- Hosseininaveh, V., Bandani, A. R., Azmayeshfard, P., Hosseinkhani, S. & Kazzazi, M. 2007. Digestive Proteolytic and Amylolytic Activities in *Trogoderma granarium* Everts (Dermestidae: Coleoptera). *Journal of Stored Products Research* 43,515–522.
- House, H. L. (1974) Digestion pp. 63–117. in: Rockstein, M. (Eds), Physiology of Insecta. Academic Press, New York,
- Jainulabdeen, S. & Prasad, S. K. (2004) Severe infection of cabbage butterfly, *Pieris brassicae* (L.) on six species of brassica and effect of abiotic factor on its population dynamics. *Journal of Entomological Research* 28(3), 193–197.
- Johnston, K. A., Lee, M. J., Gatehouse, J. A. & Anstee, J. H. (1991) The partial purification and characterization of serin protease activity in midgut of larval *Helicoverpa armigera. Journal of Insect Biochemistry* 21 (4), 389–397.
- Kumar, R., Bhardwaj, U., Kumar, P. & Mazumdar-Leighton S. (2015). Midgut serine proteases and alternative host plant utilization in *Pieris brassicae* L. *Invertabrate Physiology*, doi:10.3389/fphys.2015.00095
- Lal, M. N. & Ram, B. (2004) Cabbage butterfly, *Pieris brassicae* L. An upcoming menance for Brassicae oilseed crop in Northern India. *Cruciferae Newsletter* 25, 83– 86.
- Lewis, W. J., Van Lenteren, J.C., Phatak, S. C., & Tumlinson III, J. H. (1997) A total system approach to sustainable pest management. Proceedings of National Academy of Sciences of the United States of America 94, 12243–12248.

- Mehrkhou, F., Mahmoodi, L. & Mousavi, M. (2013) Nutritional indices parameters of large white butterfly *Pieris brassicae* (Lepidoptera: Pieridae) on different cabbage crops. *African Journal of Agricultural Research* 8, 3294–3298.
- Mehrkhou, F. & Taheri-Sarhozaki, M. (2014) Life table parameters of large white butterfly *Pieris brassicae* (Lepidoptera: Pieridae) on different cabbage varieties. Archives of Phytopathology and Plant Protection 47(12), 1444–1453.
- Naseri, B., Fathipour, Y., Moharramipour, S., Hosseininaveh, V. & Gatehouse, A. M.
 R. (2010) Digestive proteolytic and amylolytic activities of *Helicoverpa armigera* in response to feeding on different soybean cultivars. *Pest Management Science* 66, 1316–1323.
- **Oppert, B.** (2000) Transgenic plants expressing enzyme inhibitors and the prospects for biopesticide development. pp. 83–95. in Koul, O. & Dhaliwal, G. S. (Eds) Advances in Biopesticide Research, Hardwood Academic, Amsterdam,
- **Oppert, B., Hartzer, K. & Smith, C. M.** (2000) Digestive proteinases of alfalfa weevil, *Hypera postica* (Gyllenhal) (Coleoptera; Curculionidae). Transactions of the Kansas Academy of Science, (1903) 103 (3/4), 99–110.
- Pajmon, A. (1999) Pest of cabbage. Sodob. Kemetij 32: 537-540.
- Rahimi-Namin, F., Naseri, B., Arghand, A. & Azizi, M. (2011) Effect of pH and temperature on protease and amylase activities of *Pieris brassicae* (Lepidoptera: Pieridae). 2nd Iranian Pest Management Congress, Kerman University, p. 93.
- Razavi Tabatabaei, P., Hosseininaveh, V., Goldansaz, S. H. & Talebi, K. (2011) Biochemical characterization of digestive proteases and carbohydrases of the carob moth, *Ectomyelois ceratoniae* (Zeller) (Lepidoptera: Pyralidae). *Journal of Asia-Pacific Entomology* 14, 187–194.
- **Raqib, A.** (2004) Population dynamic of cabbage butterfly and cabbage aphid on five cultivars of cauliflower at Peshawar. *Asian Journal of Plant Science* 3, 391–393.
- Ryan, C. A. (1990) Proteinase inhibitors in plants: genes for improving defenses against insects and pathogens. *Annual Review of Phytopathology* 28, 425–449.
- Saadat, D., Bandani, A. R. & Dastranj, M. (2014) Comparison of the developmental time of *Bracon hebetor* (Hymenoptera: Braconidae) reared on five different lepidopteran host species and its relationship with digestive enzymes. *European Journal of Entomology* 111 (4), 495–500.
- Sachan, J. N. & Gangwar, S. K. (1980) Vertical distribution of important pest of cole crop in Meghalya as influenced by environment factors. *Indian Journal of Entomology* 42, 414–421.
- Saikia, M., Singh, Y. T., Bhattacharya, A. & Mazumdar-Leighton, S. (2011) Expression of

- diverse midgut serine proteinases in the sericigenous Lepidoptera Antheraea assamensis (Helfer) is influenced by choice of host plant species. Insect Molecular Biology 1–13.
- Sarboland, S., Mehrkhou, F. & Imani, M. (2017) Gut Proteolytic Profile of Larval Callosobruchus maculatus (Coleoptera: Chrysomelidae) in Response to Feeding on Different Fabaceous Host Plants. Journal of Agricultural Science and Technology 19, 121–132.
- Sharma, A. & Gupta, R. (2009) Biological activity of some plant extracts against *Pieris* brassicae (Linn.). Journal of Biopesticide 2(1), 26–31.
- Siraj, Q. (1999) Chemical Control and Estimation of Loses Caused by *Pieris brassicae* on Cauliflower (Seed Crop) in Swat. M.Sc. Thesis, NWFP Agriculture University, Peshawar, Pakistan, p. 40.
- Srinivasan, A., Giri, A. P., Harsulkar, A. M., Gatehouse, J. A. & Gupta, V. S. (2005) A Kunitz trypsin inhibitor from chickpea (*Cicer arietinum* L.) that exerts antimetabolic effect on podborer (*Helicoverpa armigera*) larvae. *Plant Molecular Biology* 57, 359–374.
- Srinivasan, A., Giri, A. P. & Gupta, V. S. (2006) Structural and functional diversities in lepidopteran serine proteases. *Cell Molecular Biology Letters* 11, 132–154.
- Terra, W. R. & Ferreira, C. (1994) A review, Insect digestive enzymes: properties, compartmentalization and function. *Journal of Comparative Biochemistry and Physiology* 109 B (1), 1–62.
- Velasco, P., Soengas, P., Vilar, M. & Cartea, E. M. (2008) Comparison of Glucosinolate Profiles in Leaf and Seed Tissues of Different *Brassica napus* Crops. *Journal of the American Society for Horticultural Science* 133(4), 551–558.
- Wilhite, S. E., Elden, T. C., Brzin, J. & Smigocki, A. C. (2000) Inhibition of cysteine and aspartyl proteinases in the alfalfa weevil midgut with biochemical and plant-derived proteinase inhibitors. *Insect Biochemistry and Molecular Biology* 30, 1181–1188.
- Yang, L., Fang, Z., Dicke, M., van Loon, J. J. & Jongsma, M. A. (2009) The diamondback moth, *Plutella xylostella*, specifically inactivates Mustard Trypsin Inhibitor 2 (MTI2) to overcome host plant defence. *Insect Biochemistry and Molecular Biology* 39, 55–61.
- Younas, M., Naeem, M., Raquib, A. & Masud, S. (2004) Population dynamics of *Pieris brassica* on five cultivar of cauliflower at Peshawar. *Asian Journal of Plant Science* 3, 391–393.
- Zibaee, A. (2012) Digestive enzymes of large cabbage white butterfly, *Pieris brassicae* L. (Lepidoptera: Pieridae) from developmental and site of activity perspectives. Italian *Journal of Entomology* 79 (1), 16–23.

Zibaee, A. & Hajizadeh, J. (2012) Proteolytic activity in *Plagiodera versicolora* Laicharting (Coleoptera: Chrysomelidae): Characterization of digestive proteases and effect of host plants. *Journal of Asia-Pacific Entomology* 16, 329–334.