

Effect of Curcumin on Doxorubicin-induced Cytotoxicity in H9c2 Cardiomyoblast Cells

¹Leila Hosseinzadeh, ²Javad Behravan, ³Fatemeh Mosaffa, ⁴Gholamreza Bahrami, ⁵Ahmad Reza Bahrami
⁶*Gholamreza Karimi

Abstract

Objective(s)

Doxorubicin (DOX), a widely used chemotherapeutic agent can give rise to serve cardiotoxicity by inducing apoptosis. Curcumin, the active compound of the rhizome of *Curcuma longa* L. has anti-inflammatory, antioxidant and anti-proliferative activities. Curcumin has been identified to increase cytotoxicity in several cancer cell lines in combination with DOX, but there is no study about its effect and DOX on normal cardiac cells. Therefore, in the present study, we evaluated the effect of curcumin on apoptosis induced by DOX in H9c2 rat heart-derived cells.

Materials and Methods

Cell viability was determined by MTT assay. Also, activation of caspase-3 was evaluated by spectrophotometry. Quantitative real time RT-PCR was used to evaluate the expression of c-IAP1. Detection of intracellular DOX accumulation was performed by flow cytometry.

Results

No toxicity observed when the cells exposed for 1 hr to different concentrations of curcumin, but pretreatment of cells with curcumin increased cytotoxicity of DOX in a dose dependent manner. Analysis of caspase-3 activation showed that curcumin pretreatment increased caspase-3 activation. RT-PCR analysis clearly showed that curcumin significantly decreased mRNA gene expression of c-IAP1 compared to cells treated with DOX alone. Pretreatment of H9c2 cells with DOX and curcumin had no effect on the intracellular accumulation of DOX.

Conclusion

Our observations indicated that subtoxic concentrations of curcumin sensitize H9c2 cells to DOX-induce apoptosis. These results suggest that the use of curcumin in combination with DOX in malignancy must be reevaluated.

Keywords: Apoptosis, Curcumin, Doxorubicin, H9c2 cells

1- Medical Toxicology Research Center, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

2- Biotechnology Research Center, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

3- Biotechnology Research Center, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

4- Medical Biology Research Center, Kermanshah University of Medical Sciences, Kermanshah, Iran

5- Faculty of Sciences, Ferdowsi University of Mashhad, Mashhad, Iran

6- Medical Toxicology Research Centre, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

*Corresponding author: Tel: +98-511-8823256; Fax: +98-511-8823251; email: karimig@mums.ac.ir

Introduction

Since the late 1960s, the anthracycline antibiotic doxorubicin (DOX, adriamycin) has been one of the most largely prescribed chemotherapeutic drugs for the treatment of a variety of hematologic and solid malignancies, such as leukemia, bladder, lung and breast cancer, Hodgkin's and non-Hodgkin's lymphomas, and ovarian cancer (1, 2). Unfortunately, in addition to its potent anti-tumor activity, the use of DOX is associated with a number of unwanted side effects, especially with serious cardiac toxicity which can progress to end-stage heart failure (2). The exact mechanism of DOX-induced cardiotoxicity and its progression to heart failure are not fully understood yet. Different **hypothesis** have been proposed to explain DOX induced cardiotoxicity. The supposed mechanism is the redox activation to a semi-quinone intermediate and the formation of reactive oxygen species (ROS) which ultimately results in myocytes apoptosis (3, 4). Antioxidants can inhibit DOX-induced cardiomyocyte apoptosis suggesting that ROS is involved in apoptotic cell death (5, 6).

Curcumin from *Curcuma longa* is a nutraceutical compound reported to possess therapeutic properties against a variety of diseases ranging from cancer to cystic fibrosis (7). The strong antioxidant activity of curcumin makes it an interesting candidate for use in counteracting oxidative stress-induced damage (8, 9). Curcumin has been proposed as a putative agent in the treatment of some cardiovascular disorders (10).

On the other hand, a few published studies have indicated that curcumin is able to increase DOX effectiveness in the cancer cells (7, 11). No data are available if these effects are related to **tumor** cells or it could be observed in normal cardiac cells.

For above reasons, in this study we aimed to assess the effect of curcumin pretreatment on DOX-induced apoptosis in H9c2 cells. Due to its well known antioxidant properties, we hypothesized that curcumin would be able to prevent the DOX-induced cytotoxicity in H9c2 cells. The results that we present here differ from the previous hypothesis and show

curcumin was not able to protect against DOX-mediated cardiotoxicity.

Materials and Methods

Materials

MTT [3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyltetrazoliumbromide], Curcumin and DOX were bought from Sigma Aldrich (St Louis, MO, USA). Cell culture medium, penicillin-streptomycin and fetal bovine serum (FBS) were purchased from Gibco (Gibco, Grand Island, NY, USA). Caspase-3 colorimetric assay kit was from abcam (abcam superscript® plc, USA). High pure RNA isolation kit was from Roche (Mannheim, Germany). Real time RT-PCR kit was bought from Invitrogen (Carlsbad CA, USA).

Methods

Cell culture

Rat heart cell line H9c2 was obtained from American Type Culture Collection (ATCC, CRL-1446). The H9c2 cells were grown in Dubblico modified Eagle medium (DMEM) with 4 mM L-glutamine, 4.5 g/l glucose, 1 mM sodium pyruvate, 10% (v/v) heat inactivated FBS, penicillin G (100 U/ml) and streptomycin (100 mg/ml) at 37 °C in 95% CO₂ humidified incubator. The myoblastic population would become depleted rapidly if the cultures were allowed to become confluent. Thus, to prevent loss of myoblastic population, cells were subcultured when 70% of the flask was covered with cells and plated (25000-30000 cells/cm²) in 25-cm flask for flow cytometry studies and in 6-well plate for other studies.

Cell viability assay

Cellular toxicities of DOX and curcumin were analyzed in H9c2 cells using MTT method. Cells were plated onto 96-well plates at a density of 2.0×10^4 cells/well and in a volume of 200 µl. Stock solutions of curcumin and DOX were prepared in dimethyl sulfoxide (DMSO). The final concentration of the vehicle in the medium was always 0.5%. One day after seeding, 2 µl of the DMSO containing DOX or curcumin at different concentrations was added to each well. At appropriate time intervals, the medium was

removed and replaced by 100 μ l of 0.5 mg/ml of MTT in growth medium and then the plates transferred to a 37 °C incubator for 3-4 hr. Supernatants were removed and the reduced MTT dye was solubilized with DMSO (100 μ l /well). Absorbance was determined on an ELISA plate reader with a test wavelength of 570 nm and a reference wavelength of 630 nm to obtain sample signal ($OD_{570}-OD_{630}$). Percentage of proliferation was calculated using the following formula:

Percent of control proliferation = $(OD_{test}/OD_{control}) \times 100$.

IC₅₀ values were calculated by plotting the log₁₀ of the percentage of proliferation versus drug concentration.

Caspase-3 activity

The activity of caspase-3 was determined by the abcam colorimetric caspase-3 kit according to the manufacturer's instructions. This assay is based on spectrophotometric detection of the chromophore *p*-nitroanilide (*p*NA) after cleavage from the labeled substrate 7-amino-4-trifluoromethyl coumarin conjugated *p*NA (DEVD-*p*NA) in equal amount of cells protein lysates. Briefly, 1×10^6 cells were collected and lysed with 50 μ l of chilled lysis buffer and incubated on ice for 10 min. Cell lysates were centrifuged at maximum speed for 5 min at 4 °C, after which 50 μ l of 2 \times reaction buffer/dithiothreitol (DTT) mix and 5 μ l of 1 mM caspase-3 substrate (DEVD-*p*NA) were added to each reaction and incubated at 37 °C for 1 hr. The *p*NA light emission was quantified using a microplate reader at 400- or 405- nm. Comparison of the absorbance of *p*NA from an apoptotic sample with an un-induced control allowed determination of the fold increase in caspase-3 activity. The protein content was determined by the Bradford method using the bovine serum albumin as a standard.

Real time RT-PCR analysis of the inhibitor of apoptosis protein (c-IAP1)

Total RNA of H9c2 cells was extracted using high pure isolation kit (Roche, Mannheim, Germany) according to the manufacturer's instructions. Quality and quantity of total RNA was assessed by spectrophotometry (NanoDropTM 2000, USA) and

samples stored at -80 °C until use. Quantitative specific RNA expression was performed in one step with cybr-green invitrogen kit. Specific primer sequences were as follows: For c-IAP1, sense: 5'-CTGGGAACCGAACGATGATC-3'; anti-sense: 5'-AGCCACCATCACAACAAAAGC-3'. For β -actin, sense: 5'-TTGCTGATCCACATCTGCTG-3'; anti-sense: 5'-GACAGGATGCAGAAGGAGAT-3'. Thermal cycler conditions were 15 min at 50 °C for cDNA synthesis, 10 min at 95 °C followed by 40 cycles of 15 s at 95 °C to denature the DNA and 45s at 60 °C to anneal and extend the template. We did melting curve analysis to ascertain specification by continuous acquisition from 65 °C -95 °C with a temperature transient rate of 0.1 °C/S. All reactions were performed in triplicate in a Stratagene MX 3000P system (USA). The values obtained for the target gene expression were normalized to β -actin and analyzed by the relative gene expression $-\Delta\Delta CT$ method where $-\Delta\Delta CT = (CT_{target} - CT_{\beta-actin})_{Unknown} - (CT_{target} - CT_{\beta-actin})_{Calibrator}$.

DOX uptake assay

For determination of DOX accumulation, cells were seeded in six well plates at a density of 2×10^5 cells / well and exposed to DOX, with or without 1 hr pretreatment with different concentrations of curcumin. After treatment, cells were trypsinized, washed twice with ice-cold PBS. After resuspending in ice-cold PBS, the intracellular DOX fluorescence immediately was measured using a PartecTM cytometer (Germany) equipped with a standard argon laser for 488-nm excitation and a 575-nm band pass (FL2) filter. Ten thousand (10^4) gated events were collected for all the samples. Debris was eliminated by gating on forward versus side scatter. In the above procedure, treatment only with vehicle was considered to measure of cellular auto fluorescence. Flow cytometry data were processed and analyzed using FloMax version 2.52.

Statistical analysis

Each experiment was performed at least three times, and the results were presented as mean \pm SEM. One-way analysis of variance

(ANOVA) followed by Tukey's test was used to compare the differences between means. Statistical significances were considered at $P < 0.05$.

Results

Cell viability after exposure to DOX and curcumin alone

The viability of H9c2 cardiomyoblast cells was evaluated after 18, 22 and 28 hr of exposure to different concentrations of DOX. Cell viability was evaluated by the MTT method. As shown in Figure 1, DOX-induced cytotoxicity was time and dose dependent. The mean \pm SEM IC_{50} values were $5.25 \pm 0.051 \mu M$, $3.73 \pm 0.082 \mu M$ and $1.64 \pm 0.039 \mu M$ for 18, 22 and 28 hr exposure to DOX, respectively. The appropriate incubation time for complete dose-response cytotoxicity assay was 22 hr.

In order to set curcumin at concentrations which are nontoxic to cells but could prevent DOX-induced apoptosis, we also examined the effects of different concentrations of curcumin on cell viability in H9c2 cells. The cells were incubated with curcumin ($1-50 \mu M$) for 1, 2 and 24 hr. The percentage of cell viability under different treatment conditions has been shown in Figure 2. The Figure clearly revealed that one hr treatment with curcumin had no cytotoxic effects at the concentration up to $50 \mu M$, while 2 hr and 24 hr exposure to curcumin induced cytotoxicity at the concentrations more than $30 \mu M$ and $10 \mu M$, respectively. Therefore, one hr pretreatment with curcumin was then selected.




Figure 1. Time- and dose-dependent DOX-induced H9c2 cytotoxicity. The cell viability was determined by MTT assay as described in material and methods. Data are expressed as the mean \pm SEM of three separate experiments ($n = 6$).




Figure 2. Cell viability of H9c2 cells after exposure to curcumin (CUR). Cells were treated with different concentrations of curcumin ($5-50 \mu M$) for 1 hr, 2 hr and 24 hr. The cell viability was determined by MTT assay. Data are expressed as the mean \pm SEM of three separate experiments ($n = 6$).




Figure 3. The effect of curcumin on DOX cytotoxicity in H9c2 cells. Cell pretreated with different concentrations of curcumin (CUR) 1 hr before exposure to A) $3 \mu M$ of DOX and B) indicated concentrations ($0.5, 1.5, 3$ and $6 \mu M$) of DOX. Data are expressed as the mean \pm SEM of three separate experiments ($n = 6$). ### $P < 0.01$ vs. Control, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ vs. DOX treated cells.

Effect of curcumin pretreatment on DOX-induced cell death

For evaluation of curcumin pretreatment on DOX-induced cytotoxicity, H9c2 cells were pretreated for 1 hr with curcumin ($0.5-15 \mu M$), then the medium was changed and cells were

Effect of Curcumin on DOX-induced Cytotoxicity

treated with DOX (3 μ M) for 22 hr. Surprisingly, as shown in Figure 3A, compared to control the presence of curcumin exacerbated the extent of cell death, which was significant at higher concentrations of curcumin. For further investigation, concentration–response curve was performed for DOX to access the IC_{50} value with or without 1 hr pretreatment with curcumin (5–15 μ M). The results showed that one hour pretreatment with curcumin shifted the concentration–response curves to a lower IC_{50} values in a dose dependent manner (Figure 3B) (Table 1).

Effect of curcumin pretreatment on caspase-3 activation

Activation of caspase cascade is critical in the initiation of apoptosis in various biological systems. A member of this family, caspase-3 has been identified as being a key mediator of apoptosis in mammalian cells. Therefore, to investigate the type of cell death involved in the experiments outlined above, the activity of caspase-3 was examined. The results indicated that pretreatment with curcumin significantly increased caspase-3 activation compared to DOX-treated cells (Figure 4).

Effect of curcumin on c-IAP1 expression

In order to complete our results and also to investigate how curcumin pretreatment increased DOX-induced apoptosis, we examined the mRNA expression of a member of the Inhibitors of apoptosis family cIAP1. Real time RT-PCR analysis showed a reduction of the expression level of c-IAP1 after 22 hr treatment with DOX. However, pretreatment with curcumin led to a significant down-regulation of cIAP1 compared to cells treated with DOX alone (Figure 5).

Effect of curcumin pretreatment on intracellular DOX accumulation

To examine whether curcumin-enhanced DOX potency was a result of its inhibition of any other or undefined drug-transporters, which may recognize DOX as a substrate, we evaluated the effect of curcumin on DOX accumulation in H9c2 cells. There was no significant difference in DOX accumulation between cells exposed to DOX (3 μ M) and cells pretreated with curcumin (5–15 μ M) before exposure to DOX (Figure 6).

Table 1. Effect of curcumin pretreatment on DOX-induced cytotoxicity in H9c2 cells.

group	$IC_{50} \pm SEM$	95% CI*
DOX	3.6 ± 0.049	3.477–3.726
DOX+ CUR (5 μ M)	2.5 ± 0.105^a	2.233–2.270
DOX+CUR (10 μ M)	1.82 ± 0.098^a	1.568–2.072
DOX+CUR (15 μ M)	0.95 ± 0.064^a	0.7855–1.115

Data are expressed as the mean \pm SEM of three separate experiments.

^a $P < 0.001$ vs. DOX treated cells.

* 95% confidence interval.




Figure 4. The effect of curcumin on caspase-3 activity in H9c2 cells. Cell pretreated with different concentrations of curcumin (CUR) 1 hr before exposure to 3 μ M of DOX. Caspase-3 activity measured by colorimetric detection of p-nitroanilide and expressed as percentage of control. Data are expressed as the mean \pm SEM of three separate experiments (n= 6). ## $P < 0.01$ vs. control, * $P < 0.05$, *** $P < 0.001$ vs. DOX treated cells.




Figure 5. Real-time RT-PCR analysis of effect of curcumin on cIAP1 mRNA expressions. Cell pretreated with different concentrations of curcumin (CUR) 1 hr before exposure to 3 μ M of DOX. Normalization relative to β -actin was performed. Levels of mRNA are expressed relative to control cardiomyoblast in the mean (SEM) values derived from three independent experiments, ** $P < 0.01$, *** $P < 0.001$ vs. DOX treated cells.




Figure 6. The effect of curcumin on DOX accumulation in H9c2 cells. Cell pretreated with different concentrations of curcumin (CUR) 1 hr before exposure to 3 μ M of DOX. DOX accumulation measured by flow cytometry and expressed as percent of DOX. A) Representative of flow cytometry with FL2 plots of H9c2 cells in different groups B) Column bar graph of mean cell fluorescent for FL2. Data are expressed as the mean \pm SEM of three separate experiments (n= 6).

Discussion

DOX is well known cancer chemotherapeutic agent used to treat a wide variety of human malignancies (4, 12). A major adverse effect of DOX treatment in cancer patients is the onset of cardiomyopathy and heart failure (13). Reactive Oxygen Species (ROS) derived from redox activation of DOX were proposed to be responsible for DOX-induced apoptosis in cardiac cells (6, 14, 15). Therefore, antioxidant treatment has been proposed as a potential strategy in the prevention of apoptosis in cardiac cells.

Curcumin is known for its antioxidant properties and act as free radical scavenger by inhibiting lipid peroxidation and oxidative DNA damage (16). Moreover, it has been proposed as a candidate to treat cardiovascular disease involving oxidative stress (11). There are reports that show curcumin is able to inhibit apoptosis,

through its well known antioxidant properties in normal cells (8, 17, 18).

On the other hand, it has been reported in studies that curcumin increases cytotoxicity of DOX in *cancer cells* (10,19, 20). Therefore, it is important to evaluate whether a similar phenomenon occurs in cardiomyoblast which are the main target of DOX toxicity. Consequently, our study was designed for evaluation of curcumin pretreatment against DOX cytotoxicity in H9c2 cells. At first, cytotoxicity of DOX and curcumin in different exposure times were evaluated. The results showed that one hour exposure to different concentrations of curcumin had no cytotoxic effect in H9c2 cells. However, curcumin was able to induce cytotoxicity at high concentrations, when duration of exposure exceeded one hour. It may not be a simple stoichiometric reaction. The function of curcumin was biphasic: it was non toxic at low concentrations, while higher doses promoted cell death. This is in line with a study that demonstrated 24 hr treatment with 5 μ M curcumin resulted marginal decrease in cell viability in murine macrophage RAW 264.7 cells (21). Additionally, another study showed that the concentrations of curcumin higher than 20 μ M were able to induce toxicity in the MES23.5 cells (8). Although, curcumin has been described as a protective agent in several cell models, we observed here that pretreatment to subtoxic concentrations of curcumin greatly increased cytotoxicity compared to level reached with the DOX alone. The present findings corroborate similar findings by Ortiz-Ortiz, *et al* who recently reported that curcumin enhanced paraquat induced cytotoxicity in N27 mesencephalic cells (16).

It has been reported that DOX treatment was able to induce cytotoxicity in cardiac cells through induction of apoptosis (2, 15). Many proteins are involved in this complex process. Caspases, a family of cysteine-dependent aspartate-directed proteases, play a critical role in the initiation and execution of apoptosis. Among this family of caspases, caspase 3, in particular, is believed to be one of the most commonly involved caspases in the execution of apoptosis in various cell types (22). Therefore, we assayed caspase-3 activation for

characterizing the nature of cell death that occurred in response to the pretreatment with curcumin. Curcumin increased DOX-induced caspase-3 activation in a dose dependent manner.

To confirm that synergistic effect of curcumin was due to apoptosis, we investigated the mRNA expression of a member of the inhibitors of apoptosis family cIAP1. IAPs have shown a remarkable ability of blocking cell death. It was demonstrated that caspase activation not only requires proteolytic cleavage of the enzyme themselves, but removal of inhibitory influences (IAPs) is also necessary (10). Curcumin has been reported to down-regulate the anti-apoptotic IAP proteins in human renal cells (23). Also, it was able to decrease cIAP1 and cIAP2 expression in human glioblastoma T988G cells (24).

In the present study, DOX slightly decreased inhibition of apoptotic protein c-IAP1 mRNA level after 22 hr, but curcumin potentiated this declining.

Thus, evaluation of caspase-3 activation and c-IAP1 expression, demonstrated that curcumin is able to potentiate the apoptotic effect of DOX in H9c2 cells. These results are in contrast with previously described findings that curcumin is able to protect normal cells against some cytotoxic agents (8, 17, 18).

Curcumin could also chemosensitize resistant tumor cells to DOX through an increase in the accumulation of DOX in the human carcinoma cell line (25). Therefore, we investigated whether curcumin pretreatment had an effect on the DOX accumulation in cardiomyoblast cells. The results indicated that the drug toxicity enhancement can not be explained by alteration of cellular DOX levels.

Several mechanisms may explain how

curcumin mediates its effect on cardiac myoblast cells. Oxidative stress plays an important role in DOX-induced apoptosis in cardiac cells. Recent studies have showed that curcumin, although a potent antioxidant with ROS scavenging properties was able to enhance ROS formation in different models. For instance rapid ROS generation by curcumin leads to caspase-dependent and independent apoptosis in L929 cells (26). Also, curcumin increased oxidative stress induced by paraquat via inhibition of nitric oxide synthase (NOS) (16). Therefore, it is possible that curcumin enhanced DOX-induced apoptosis through generation of ROS. However, there is still the lack of information on specific molecular mechanisms induced by synergistic combination of curcumin and DOX because curcumin influences many cellular signaling pathways (27), that play major role in the control of cell proliferation and survival.

Further studies are in progress in our laboratory to determine molecular mechanisms involved in synergistic effect of curcumin and DOX in rat cardiac muscle H9c2 cells.

Conclusion

For the first time our findings demonstrated that curcumin enhanced the apoptosis induced by DOX in H9c2 cells. These results suggest that the use of curcumin in combination with DOX in malignancy must be reevaluated.

Acknowledgment

The results presented here are extracted from the PhD thesis of L Hosseinzadeh. This study was financially supported by the Research Council of Mashhad University of Medical Sciences, Mashhad, Iran.

References

1. Tokarska-Schlattner M, Zaugg M, Zuppinger C, Wallimann T, Schlattner U. New insights into doxorubicin-induced cardiotoxicity: The critical role of cellular energetics. *J Mol Cell J Mol Cell* 2006; 41:389-405.
2. Wang S, Konorev EA, Kotamraju S, Joseph J, Kalivendi S, Kalyanaraman B. Doxorubicin induces apoptosis in normal and tumor cells via distinctly different mechanisms: Intermediacy of H₂O₂- and p53-dependent pathways. *J Biol Chem* 2004; 279:25535-25543.
3. Spallarossa P, Garibaldi S, Altieri P, Fabbi P, Manca V, Nasti S, *et al.* Carvedilol prevents doxorubicin-induced free radical release and apoptosis in cardiomyocytes *in vitro*. *J Mol Cell Cardiol* 2004; 37:837-846.
4. Mizutani H, Tada-Oikawa S, Hiraku Y, Kojima M, Kawanishi S. Mechanism of apoptosis induced by doxorubicin through the generation of hydrogen peroxide. *Life Sci* 2005; 76:1439-1453.

5. Kim D-S, Woo E-R, Chae S-W, Ha K-C, Lee G-H, Hong S-T, *et al*. Plantainoside D protects adriamycin-induced apoptosis in H9c2 cardiac muscle cells via the inhibition of ROS generation and NF-[kappa]B activation. *Life Sci* 2007; 80:314-323.
6. Han X, Ren D, Fan P, Shen T, Lou H. Protective effects of naringenin-7-O-glucoside on doxorubicin-induced apoptosis in H9C2 cells. *Eur J Pharmacol* 2008; 581:47-53.
7. Duvoix A, Blasius R, Delhalle S, Schneckeburger M, Morceau F, Henry E, *et al*. Chemopreventive and therapeutic effects of curcumin. *Cancer Lett* 2005; 223:181-190.
8. Wang J, Du XX, Jiang H, Xie JX. Curcumin attenuates 6-hydroxydopamine-induced cytotoxicity by anti-oxidation and nuclear factor-kappaB modulation in MES23.5 cells. *Biochem Pharmacol* 2009; 78:178-183.
9. Chen W, Zhao Z, Li L, Wu B, Chen Sf, Zhou H, *et al*. Hispolon induces apoptosis in human gastric cancer cells through a ROS-mediated mitochondrial pathway. *Free Radic Biol Med* 2008; 45:60-72.
10. Fiorillo C, Becatti M, Pensalfini A, Cecchi C, Lanzilao L, Donzelli G, *et al*. Curcumin protects cardiac cells against ischemia-reperfusion injury: effects on oxidative stress, NF-kB, and JNK pathways. *Free Radic Biol Med* 2008; 45:839-846.
11. Notarbartolo M, Poma P, Perri D, Dusonchet L, Cervello M, D'Alessandro N. Antitumor effects of curcumin, alone or in combination with cisplatin or doxorubicin, on human hepatic cancer cells. Analysis of their possible relationship to changes in NF-kB activation levels and in IAP gene expression. *Cancer Lett* 2005; 224:53-65.
12. Wold LE, Aberle Ii NS, Ren J. Doxorubicin induces cardiomyocyte dysfunction via a p38 MAP kinase-dependent oxidative stress mechanism. *Cancer Detect Prev* 2005; 29:294-299.
13. Gewirtz DA. A critical evaluation of the mechanisms of action proposed for the antitumor effects of the anthracycline antibiotics adriamycin and daunorubicin. *Biochem Pharmacol* 1999; 7:727-741.
14. Jang YM, Kendaiah S, Drew B, Phillips T, Selman C, Julian D, *et al*. Doxorubicin treatment in vivo activates caspase-12 mediated cardiac apoptosis in both male and female rats. *FEBS Lett* 2004; 577:483-490.
15. Spallarossa P, Fabbi P, Manca V, Garibaldi S, Ghigliotti G, Barisione C, *et al*. Doxorubicin-induced expression of LOX-1 in H9c2 cardiac muscle cells and its role in apoptosis. *Biochem Biophys Res Commun* 2005; 335:188-196.
16. Ortiz-Ortiz MA, Morán JM, Bravosánpedro JM, González-Polo RA, Niso-Santano M, Anantharam V, *et al*. Curcumin enhances paraquat-induced apoptosis of N27 mesencephalic cells via the generation of reactive oxygen species. *Neurotoxicology* 2009; 30:1008-1018.
17. Srinivasan M, Rajendra Prasad N, Menon VP. Protective effect of curcumin on [gamma]-radiation induced DNA damage and lipid peroxidation in cultured human lymphocytes. *Mutat Res* 2006; 611:96-103.
18. Bishnoi M, Chopra K, Kulkarni SK. Protective effect of curcumin, the active principle of turmeric (*Curcuma longa*) in haloperidol-induced orofacial dyskinesia and associated behavioural, biochemical and neurochemical changes in rat brain. *Pharmacol Biochem Behav* 2008; 88:511-522.
19. Choi BH, Kim CG, Lim Y, Shin SY, Lee YH. Curcumin down-regulates the multidrug-resistance *mdr1b* gene by inhibiting the PI3K/Akt/NF [kappa] B pathway. *Cancer Lett* 2008; 259:111-118.
20. Chuang SE, Yeh PY, Lu YS, Lai GM, Liao CM, Gao M, *et al*. Basal levels and patterns of anticancer drug-induced activation of nuclear factor-kB (NF-kB), and its attenuation by tamoxifen, dexamethasone, and curcumin in carcinoma cells. *Biochem Pharmacol* 2002; 63:1709-16.
21. Kunwar A, Sandur SK, Krishna M, Priyadarsini KI. Curcumin mediates time and concentration dependent regulation of redox homeostasis leading to cytotoxicity in macrophage cells. *Euro J Pharmacol* 2009; 611:8-16.
22. Garg S, Narula J, Chandrashekhar Y. Apoptosis and heart failure: clinical relevance and therapeutic target. *J Mol Cell Cardiol* 2005; 38:73-79.
23. Woo JH, Kim YH, Choi YJ, Kim DG, Lee KS, Bae JH, *et al*. Molecular mechanisms of curcumin-induced cytotoxicity: Induction of apoptosis through generation of reactive oxygen species, down-regulation of Bcl-X and IAP, the release of cytochrome c and inhibition of Akt. *Carcinogenesis* 2003; 24:1199-1212.
24. Karmakar S, Banik NL, Patel SJ, Ray SK. Curcumin activated both receptor-mediated and mitochondria-mediated proteolytic pathways for apoptosis in human glioblastoma T98G cells. *Neurosci Lett* 2006; 407:53-62.
25. Zhang HZ, Yang L, Ren LM, Liu SM. Effects of curcumin combining with adriamycin to kill KB and KBv200 cells. *Chin Pharmacol Bull* 2001; 17:702-704.
26. Thayyullathil F, Chathoth S, Hago A, Patel M, Galadari S. Rapid reactive oxygen species (ROS) generation induced by curcumin leads to caspase-dependent and -independent apoptosis in L929 cells. *Free Radic Biol Med* 2008; 45:1403-1412.
27. Reuter S, Eifes S, Dicato M, Aggarwal BB, Diederich M. Modulation of anti-apoptotic and survival pathways by curcumin as a strategy to induce apoptosis in cancer cells. *Biochem Pharmacol* 2008; 76:340-351.