



## The relaxant effect of *Nigella sativa* on smooth muscles, its possible mechanisms and clinical applications

Rana Keyhanmanesh <sup>1</sup>, Zahra Gholamnezhad <sup>2</sup>, Mohammad Hossien Boskabady <sup>2\*</sup>

<sup>1</sup> Tuberculosis and Lung Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

<sup>2</sup> Neurogenic Inflammation Research Centre and Department of Physiology, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

### ARTICLE INFO

#### Article type:

Review article

#### Article history:

Received: Jul 23, 2014

Accepted: Nov 20, 2014

#### Keywords:

Clinical application

*Nigella sativa*

Possible mechanisms

Relaxant effect

Thymoquinone

### ABSTRACT

*Nigella sativa* (*N. sativa*) is a spice plant which has been traditionally used for culinary and medicinal purposes. Different therapeutic properties including the beneficial effects on asthma and dyspnea, digestive and gynecology disorders have been described for the seeds of *N. sativa*. There is evidence of the relaxant effects of this plant and some of its constituents on different types of smooth muscle including rabbit aorta, rabbit jejunum and trachea. The relaxant effect of *N. sativa* could be of therapeutic importance such as bronchodilation in asthma, vasodilation in hypertension and therapeutic effect on digestive or urogenital disorders. Therefore in the present article, the relaxant effects of *N. sativa* and its constituents on smooth muscles and its possible mechanisms as well as clinical application of this effect were reviewed.

#### ► Please cite this paper as:

Keyhanmanesh R, Gholamnezhad Z, Boskabady MH. The relaxant effect of *Nigella sativa* on smooth muscles, its possible mechanisms and clinical applications. Iran J Basic Med Sci. 2014; 17:939-949.

### Introduction

*Nigella sativa* (*N. sativa*) is a spice plant belonging to the family from Ranunculaceae (1). Different local names of the plant are; in Farsi, it's commonly known as Siah-Daneh or Fitch in south Asia, it is called Kalonji, its name in Arabic is Habat-ul-Sauda and its English, black cumin or black seeds. It is an annual grassy plant with green-to blue-colored flowers and black seeds.

Traditionally, there is a common Islamic belief that the 'black seed', is a panacea (universal healer) that is a remedy for all ailments, but cannot prevent aging or death (2-3). *N. sativa* has been traditionally used in Iran (4), the Indian subcontinent, Arabian countries (5) and Europe for culinary and medicinal purposes as a natural remedy for a number of illnesses and conditions (6-8). The fixed oil from seeds of *N. sativa* is useful in skin eruptions, paralysis, hemiplegia, back pain, rheumatoid and related inflammatory diseases (9-11). In view of the folkloric, the plant mixture extracts have been used for treatment of diabetes in the Middle East.

The extract of *N. sativa* was effective in reducing blood glucose due to inhibition of hepatic gluconeogenesis and its possible insulinotropic properties (12-18). In addition, different extracts of *N. sativa* and its constituents could lower blood pressure

through blockade of calcium channels (19-24) and showed a potent inhibitory effect on both heart rate and contractility of isolated guinea pig heart which was comparable and even higher than that of diltiazem (25-26). *N. sativa* seeds and its constituent, thymoquinone were also found to reduce the levels of serum cholesterol, triglyceride and glucose as well as leukocyte and platelet counts (16, 23, 27-31). It has also inhibitory effects on arachidonic acid-induced platelet aggregation and blood coagulation (32). In the healthy female volunteers, it was shown that *N. sativa* crushed seeds and its total oil decreased glucose, prolactins, triglycerides and cholesterol level (33).

Moreover, *N. sativa* oil and its constituent, thymoquinone, were found to possess gastroprotective effect against gastric lesions and reduced the volume of gastric acid secretion, free acidity, total acidity and ulcer index (34, 35) which may be related to the conservation of the gastric mucosal redox state (36). Significant increase in mucin content, glutathione level and a significant decrease in mucosal histamine content in the stomach were also shown for the plant seed and thymoquinone (15, 37).

*N. sativa* and its constituents have been traditionally used to alleviate respiratory disorders such as asthma, bronchospasm and chest congestion (38,

\*Corresponding author: Mohammad Hossien Boskabady, Neurogenic Inflammation Research Centre and Department of Physiology, School of Medicine, Mashhad, Post Code 9177948564, Iran. Tel : 098 511 8828565; Fax 0098 511 8828564; email: boskabadyhm@mums.ac.ir, mhboskabady@hotmail.com

39). Nigellone, a carbonyl polymer of thymoquinone, was shown to be an effective prophylactic agent in asthma and bronchitis with higher efficacy in children than in adults (40, 41). Chakravarty (42) concluded that the nigellone inhibit effectively the histamine release from the mast cells, thus showing the basis for its traditional use in asthma. It was also suggested that *N. sativa* volatile oil might be a potentially useful respiratory stimulant (43). The antihistaminic effect of the plant was also reported (44, 45).

The prophylactic effect of boiled extract of *N. sativa* on asthmatic patients as improvement in the symptom score, asthma symptom/week, chest wheeze and pulmonary function test (PFT) values in a three months treatment period was also shown (46). In addition, the beneficial role of *N. sativa* in reducing wheeze associated with lower respiratory tract illness in children (47) and adult asthmatic attacks has been reported (48). The bronchodilatory effect of boiled extract of *N. sativa* on asthmatic patients was also documented by increasing pulmonary function tests and specific airway conductance (sGaw) of the patients after administration of the extract (49).

The antitussive effect of *N. sativa* and its constituent, thymoquinone in guinea pigs has been shown (50,51). Moreover, the preventive effect of hydroethanolic extract of *N. sativa* and its main constituent, thymoquinone, in ovalbumin sensitized guinea pigs and rats, as animal models of asthma was documented (52-56). In a current study, the preventive effect of the hydroethanolic extract of this plant was proved on the tracheal responsiveness of cigarette smoke-exposed guinea pigs (57). In another study, the preventive effect of long term administration of three different concentrations of carvacrol, another constituent of the plant has been shown in asthmatic guinea pigs (58).

With regard to various physiologic effects of smooth muscle relaxant effect such as bronchodilation in obstructive pulmonary diseases, reducing high blood pressure by vasodilation and due to several studies examining the relaxant effect of *N. sativa* and its constituents on smooth muscles, this property of the plant and its possible mechanism(s) as well as clinical application of this effect was reviewed in the present article.

## Methods

Online literature searches were performed using Medline, Pubmed, Iran medex, Scopus, and Google Scholar websites backed to 1959 to identify researches about *N. sativa*, its constituents and relaxant effect.

## Constituents

The seed oil of *N. sativa* was found to be rich in polyphenols and tocopherols (59-60). The plant's seeds contain a yellowish volatile oil; 0.5-1.6% (61) or 0.4-2.5% (62), a fixed oil; 35.6-41.6%, proteins;

22.7% (61), different amino acids (such as arginine, glutamic acid, leucine, lysine, methionine, tyrosine, proline and threonine), carbohydrates; 33.9% and fiber; 5.5% (63). The other compositions of the plant include; fats, crude fibers, minerals; e.g. Fe, Na, Cu, Zn, P, Ca and many vitamins. *N. sativa* seeds yield esters of fatty acids (61), free sterols, steryl esters, steryl glucosides and acylated steryl glucosides (64). Fatty acids were identified in the extract of the plant, which represented about 99.5% of the total fatty acid composition. The fixed oil is composed mainly of fatty acids consisted of four saturated fatty acids (17.0%) and four unsaturated fatty acids (82.5%). The major unsaturated fatty acids were linoleic acid (C18:2, 55.6%) and oleic acid (C18:1, 23.4%), while the main saturated fatty acids was palmitic acid (C16:0, 12.5%) and stearic (C18:0) acids (62, 65,66).  $\beta$ -sitosterol was the dominant sterol (69%); while campesterol and stigmasterol constituted 12% and 19%, of the total sterols respectively (62).

Pharmacologically active constituents of the volatile oil of the plant are thymoquinone; 27.8-57% (61) or 30-48% (67), dithymoquinone, thymohydroquinone and thymol. The content of thymoquinone is highest (57-78%) when water has withheld for 12 days (61). Nigellone is the carbonyl polymer of thymoquinone, isolates from *N. sativa* seeds. This polymer is far less toxic but retains much of the pharmacologic properties of thymoquinone, which is its active principle (42). Other components isolated from the essential oil of the plant included; alpha-pinene (13.75 %), *p*-cymene (7.1-15.5 %), carvacrol (5.8-11.6%), trans-anethole (0.25-2.3% or 38.3%), 4-terpineol (2.0-6.6%), longifoline (1.0-8.0%), limonene (4.3%) and carvone (4.0%), (39, 65, 68-69). Four alkaloids have been reported as constituents of *N. sativa* including; nigellicine (67, 70), nigellidine which contains an indazole nucleus (71-72), nigellimine (73) and isoquinolines (74).

Briefly, thirty-two compounds, constituting 86.7% of the volatile oil, were identified. The oil consisted of six phenyl propanoid compounds (46.1%), nine monoterpenoid hydrocarbons (26.9%), four monoterpenoid ketones (6.0%), eight nonterpenoid hydrocarbons (4.0%), three monoterpenoid alcohols (2.7%) and two sesquiterpenoid hydrocarbons (1.0%). So, the oil of *N. sativa* is characterized by a large amount of phenyl propanoids (65). Table 1 summa-rized the different chemical composition of the plant.

## Relaxant effects on various types of smooth muscles

The relaxant effect of various preparations of *N. sativa* and its constituents on different types of smooth muscles were demonstrated which reviewed in the following sections.

### Tracheal smooth muscle

Reiter and Brandt examined the relaxant effects of 22 plants extracts from 11 different families including *N. sativa* on tracheal and ileal smooth

**Table 1.** Chemical composition of *N. sativa* based on the type of plant preparation

Plant preparation	Constituents	Reference	
Seed oil	Polyphenols and tocopherols	(59-60)	
	Volatile oil	(61-62)	
	Fixed oil	(61)	
	Proteins	(61)	
	Aminoacids (Arg, Glu, Leu, Lys, Met, Tyr, Pro, Thr)	(63)	
	Carbohydrates	(63)	
	Fibers	(63)	
	Minerals ( Fe, Na, Cu, Zn, P, Ca)	(63)	
	Vitamins	(63)	
	Fat: esters of fatty acids, free sterols, steryl esters, steryl glucosides, acylated steryl glucosides	(61, 64)	
	Alkaloids: Nigellidine	(67, 70)	
	Nigellimine	(71-72)	
	Nigellimine	(73)	
	Isoquinolines	(74)	
	Volatile oil	Thymoquinone, Dithymoquinone, Thymohydroquinone or Thymol	(61, 67)
		Nigellone	(42)
p-cymene, Carvacrol, t-anethole, 4-terpineol, limonene, carvone and longifoline		(65, 68-69)	

muscles of the guinea pig in comparison with the relaxant effects of catecholamines and phosphodiesterase inhibitors. All of the oils had relaxant effects on tracheal smooth muscle. However, the extract of *N. sativa* had a higher relaxant effect on tracheal muscle than ileal smooth muscles (75). A previous study also showed that the volatile oil of *N. sativa* could protect guinea pigs against histamine-induced bronchospasm but it did not affect histamine H<sub>1</sub> receptors in isolated tissues (41).

Many studies demonstrated the relaxant effects of a crude extract including aqueous and macerated extracts of *N. sativa* seeds on isolated guinea pig tracheal preparation (76-78). In one study, the effect of three concentrations of aqueous extract of *N. sativa* (0.25, 0.5 and 1 g%) have been evaluated on calcium induced contraction of guinea pig tracheal smooth muscle. The results elucidated the rightward shift in calcium concentration response curves indicating the relaxant effect of this extract on isolated guinea pig tracheal smooth muscle compared to saline (79).

The relaxant effects of hydroethanolic, macerated, aqueous and lipid free macerated aqueous extract of *N. sativa* on tracheal smooth muscle of guinea pigs were also examined. The results showed that mainly water soluble substances of the plant were responsible for the relaxant effect of the plant on airway smooth muscle (80). The relaxant effects of four fractions of *N. sativa* including; n-hexane, dichloromethane, methanol and aqueous fractions, on tracheal smooth muscle of guinea pigs were also examined. These results showed a potent relaxant effect of most fractions of the plant on airway smooth muscle. The most potent relaxant effect was seen for methanol and dichloromethane fractions (81). In a recent study, in order to identify main constituents of the methanolic extract, the relaxant effects of five different methanolic fractions (20%, 40%, 60%, 80%, and 100%) of *N. sativa* on tracheal smooth muscle of guinea pigs were examined. The results of this study showed a potent relaxant effect

of 20% methanolic fractions of this plant (82). Four constituents of 20%-methanolic fraction, consisting of two flavonoids (20-20% and 21-20% fractions) and two polysaccharides (1-20% and 2-20% fractions), were purified by analytical and preparative HPLC and their relaxant effects were examined on methacholine-precontracted guinea pig tracheal smooth muscle. The findings revealed that two flavonoids of 20%-methanolic fraction of *N. sativa* were its main relaxant constituent (83).

The effect of thymoquinone, the main constituent of the volatile oil of *N. sativa*, on the guinea pig isolated tracheal preparation showed that thymoquinone caused a concentration-dependent decrease in the tension of the tracheal smooth muscle (84). However, the results of this study was not in agreement to the findings of other study in which three cumulative concentrations (40, 80, and 120 µM) of thymoquinone did not show any relaxant effect on methacholine or KCl precontracted tracheal smooth muscle of guinea pigs (85).

In addition, the preventive effects of two concentrations of hydroethanolic extract of *N. sativa* on the tracheal responsiveness of ovalbumin sensitized guinea pigs, an animal model of asthma, were examined. The tracheal responses of treated groups with both concentrations of the plant to methacholine and ovalbumin were significantly decreased compared to those of sensitized group. Incubation of the tracheal smooth muscle with the *N. sativa* extract also caused a further decrease in the tracheal responsiveness to methacholine and ovalbumin and a decrease in the contractility response. These results confirmed the relaxant effect of the plant on airway smooth muscle (52). Moreover, the prophylactic effect of short term (56) and long term (54) administration of two doses of thymoquinone, the main constituent of *N. sativa* has also been shown similar results on tracheal smooth muscle. The potent relaxant effect of carvacrol, another constituent of *N. sativa* on guinea pig tracheal smooth muscle was also demonstrated (86-88).

**Table 2.** The relaxant effect of *Nigella sativa* and its constituents on different types of smooth muscle. In most studies, Smooth muscle was contracted, the extract was added and the effect was measured

Type of smooth muscle (SM)	Effect	Reference
Trachea	Relaxant effect	(75-84)
	Protected against histamine-induced bronchospasm	(41)
	Preventive effects on the tracheal responsiveness	(52, 54)
Vascular	Depressant action on heart	(89)
	Vasorelaxation	(21, 90)
Gastrointestinal	Spasmolytic effect	(75, 91-93)
Urogenital	Inhibition of spontaneous movements	(1)

#### Vascular smooth muscle

The essential oil from the seeds of *N. sativa* exhibited a depressant effect on the frog heart and a relaxant property on isolated vascular smooth muscles of rat (89). The vasorelaxation effect of cumulative concentration of the hydroethanolic extract of *N. sativa* (2, 4, 6, 8, 10, and 14 mg/ml) on rat aortic smooth muscle contracted by both KCl and phenylephrine was shown (21). In other study, the effect of thymoquinone on smooth muscle of pulmonary artery was assessed. The finding showed that thymoquinone caused a concentration-dependent decrease in the tension of the pulmonary arterial smooth muscles precontracted by phenylephrine (90).

#### Gastro-intestinal and urogenital smooth muscle

The oils of 16 plants including *N. sativa* inhibited the phasic contractions of the ileal myenteric plexus-longitudinal muscle preparation (75). The volatile oil and ethanol extract of *N. sativa* inhibited spontaneous movements of the rabbit jejunum and prevented contractions induced by high potassium ( $K^+$ ) solution or acetylcholine. This inhibition was dose-dependent, reversible and was not affected by the addition of calcium to the organ bath. These data suggest that the plant has an antispasmodic effect, possibly due to a calcium antagonistic activity (91).

In another study, the inhibitory effect of different doses (10, 20, 40, 80, 100 mM) of thymoquinone on the contractility of guinea pig isolated ileum was evaluated. The findings indicated that the major constituent of *N. sativa*, thymoquinone, inhibit guinea pig isolated ileum contractility, and this effect may be responsible for the smooth muscle relaxant activity of this plant (92). For assessing the spasmolytic activity of thymoquinone, it was added cumulatively after using high  $K^+$  concentration to depolarize the rabbit jejunum and guinea pig ileum smooth muscles in organ bath. In this investigation, sustained contractions were obtained with high  $K^+$  which involves entry of  $Ca^{2+}$  into the cells through voltage-operated  $Ca^{2+}$  channels. Thymoquinone showed a concentration-dependent relaxant activity on both intestinal smooth muscle preparations. According to the results of this study, it has been concluded that thymoquinone-induced smooth muscle relaxation occurred via blockade of voltage-operated  $Ca^{2+}$  channels in intestinal jejunal and ileal tissues (93). The volatile oil of *N. sativa* inhibited the spontaneous movements of rat and guinea pig

uterine smooth muscle and also the contractions induced by oxytocin, showing its antioxytotic potential (1).

The relaxant effect of the plant on different types of smooth muscles was summarized in Table 2.

#### Possible mechanisms of the relaxant effect of *N. sativa* on smooth muscles

The relaxant effects seen for *N. sativa* on smooth muscles in various studies might be produced due to several different mechanisms which are described in this part of article.

#### Calcium channel blocking effect

Gilani and his colleagues studied the effects of extract of *N. sativa* seeds on isolated tracheal and ileal smooth muscle and showed that the aqueous-methanolic extract of this plant had spasmolytic effect mediated through calcium channel blocking activity. This study showed that the extract antagonized the contractions induced by histamine, carbachol and KCl. The relaxant effect of the plant on smooth muscles was suggested to be mediated via calcium channels. The above pharmacological activities of the petroleum ether fraction of the extract were about 10 times higher than those of the crude extract (76). The calcium-antagonistic effects of three increment concentrations (0.25, 0.5 and 1 g%) of the aqueous extract of *N. sativa* were also tested by measuring the cumulative concentration response curves of  $CaCl_2$ -induced contractions of isolated guinea pig tracheal smooth muscle in the presence of diltiazem, a calcium channel blocker and the extract. The results showed rightward shift in  $CaCl_2$ -concentration response curve in the presence of the extract similar to the effect of diltiazem which suggest a calcium antagonistic effect of *N. sativa* in isolated tracheal smooth muscle (79). However, the absence of inhibitory effects of the extracts on KCl induced contraction of tracheal smooth muscle suggested that the calcium channel blocking effect of this plant did not contribute to the relaxant effect of this plant on the tracheal smooth muscle of guinea pigs (78).

The effect of *N. sativa* aqueous extract on calcium channels was also examined more precisely by its effects on heart rate and contractility of isolated heart in the presence of calcium free Krebs solution. The findings indicated that the extract of this plant almost completely inhibited calcium channels of

isolated guinea pig heart (26). Previous experiment implied that *N. sativa* seed is able to relax the aorta and by inhibition of inositol triphosphate (IP<sub>3</sub>) and/or ryanodine receptor-dependent release of intracellular Ca<sup>2+</sup> (21). It was also shown that the voltage-dependent Ca<sup>2+</sup> channels are involved in KCl-induced contraction. Therefore, voltage-dependent Ca<sup>2+</sup> channels blockade is the suggested mechanism of vasorelaxant substances which affect KCl induced contractions (21, 94). The dose-dependent inhibitory effect of volatile oil of *N. sativa* on contractions of the rabbit jejunum induced by high potassium (K<sup>+</sup>) solution or acetylcholine was also shown which was not affected by the addition of calcium to the organ bath. These results also suggest that the antispasmodic effect of the plant is possibly due to a calcium antagonistic activity (91). The calcium channel blocking effect of *N. sativa* was also confirmed by various other studies (20, 22-24)

In other studies, the relaxant activity of thymoquinone, a bioactive component of *N. sativa* was evaluated on rabbit and guinea pig cardiac, intestinal and tracheal smooth muscles. The data of these studies suggested that its relaxant effect on smooth and cardiac muscles partially mediated via blockade of voltage-operated Ca<sup>2+</sup> channels (92-93).

The inhibited spontaneous movements of the rabbit jejunum and the preventive effect of volatile oil and ethanol extract of *N. sativa* on contractions induced by high potassium (K<sup>+</sup>) solution or acetylcholine which was not affected by the addition of calcium to the organ bath also suggested a calcium antagonistic activity of the plant (91). Concentration-dependent relaxant effect of thymoquinone on intestinal smooth muscle preparations contracted with high K<sup>+</sup> which involves entry of Ca<sup>2+</sup> into the cells through voltage-operated Ca<sup>2+</sup> channels also showed involvement of voltage-operated Ca<sup>2+</sup> channels blockade in this effect (93).

#### *Potassium channel opening effect*

The inhibitory effect of aqueous and macerated extracts of *N. sativa* on contraction of tracheal smooth muscle in the presence of both ordinary and calcium free Krebs solutions have been examined previously (78). The extracts of *N. sativa* did not show any relaxant effect on KCl induced tracheal smooth muscle contraction, but they had a relaxant effect when the muscle was contracted by methacholine. These results may indicate an opening effect of the plant on potassium channels because, if a compound had a potassium channel opening effect, it would not have relaxant effects on smooth muscle contracted by KCl, while it could show a relaxant effect on methacholine induced muscle contraction. In addition, the bronchodilatory effect of potassium channel openers has been demonstrated previously (95) which support this suggestion.

On the other hand, membrane hyperpolarization is the result of an efflux of K<sup>+</sup> rises due to the opening of the K<sup>+</sup> channels in the vascular smooth muscle cells. This effect is followed by the closure of voltage-dependent Ca<sup>2+</sup> channels, leading to the reduction in Ca<sup>2+</sup> entry and vasodilation (96). Vascular smooth muscle cells express both K<sub>ATP</sub> and nonselective K<sup>+</sup> channel (97-98). A previous study indicated the nonselective blockade of the K<sub>ATP</sub> channel by *N. sativa* extract in rat aorta (21). The relaxant effect of different concentrations (50, 100, 200, 500 and 1000 μM) of thymoquinone has also shown in the isolated rat pulmonary artery. In this study, blocking of ATP-sensitive K<sup>+</sup> channel by glibenclamide significantly reduced thymoquinone-induced relaxation of pulmonary arterial smooth muscle. These findings also suggested that relaxation of pulmonary arterial smooth muscle by thymoquinone could be through K<sup>+</sup> channel activation (90).

#### *The anticholinergic and muscarinic receptor inhibitory effects of the plant*

The anticholinergic effect of aqueous and macerated extracts of *N. sativa* has been demonstrated by performing concentration response curve to methacholine, a muscarinic agonist in the presence of the extracts and atropine (77). The extracts of *N. sativa* and atropine showed rightward shifts in methacholine concentration response curve but the maximum response to methacholine was not achieved in the presence of macerated extract. These results indicated a competitive antagonistic effect of aqueous extract and a non-competitive antagonistic effect of macerated extract of *N. sativa* on muscarinic receptors (77). In addition, the existence of α-pinene in essential oil of this plant was demonstrated which showed anticholinergic activity (99).

The contribution of muscarinic receptors blocking effect on the relaxant effect of extracts of *N. sativa* was examined on incubated tracheal preparations with atropine. The results indicated a functional antagonism effect of aqueous extract from this plant at muscarinic receptors (78).

The inhibitory effect of carvacrol, a constituent of *N. sativa* on muscarinic receptors of tracheal smooth muscle as one possible mechanism responsible for its relaxant effect was also had been shown (87). The slopes of the Schild plot in the experimental groups were supported the notion of competitive antagonism of carvacrol on muscarinic receptors of guinea pig tracheal smooth muscle. These findings indicated a competitive antagonistic effect of carvacrol at muscarinic receptors of tracheal smooth muscle (87).

#### *Histaminic (H1 receptor) antagonistic activity*

Mahfous and EL-Dakhakhny reported that the volatile oil from *N. sativa* protected guinea pigs against bronchospasm induced by histamine, but in

**Table 3.** Possible mechanisms of the relaxant effect of *Nigella sativa* and its constituents

Possible mechanisms	Type of muscle	Reference
Calcium antagonistic effect	Trachea	(76, 78-79, 92-93)
	Vascular	(21, 26, 92-94)
	Gastrointestinal	(92-93)
Potassium channel opening effect	Trachea	(95)
	Vascular	(21, 90)
Anticholinergic and anti-muscarinic effect	Trachea	(77-78, 87-88, 99)
Histaminic antagonistic activity	Trachea	(41, 77, 101-102)
Stimulatory effect on $\beta$ 2-adrenoceptors	Trachea	(103-105)

isolated tissues, it did not affect histamine H<sub>1</sub> receptors (41). However, in an *in vivo* study, increasing respiratory rate and intra-tracheal pressure of guinea pigs due to intravenous administration of volatile oil from *N. sativa* has been demonstrated (100). The reason(s) of this discrepancy is unclear to us.

The concentration response curves of histamine induced contraction of smooth muscle were performed in the presence of saline, the plant extracts and chlorpheniramine. The results showed a parallel rightward shift in the presence of macerated extract compared to the curve obtained in the presence of saline similar to that of chlorpheniramine and the maximum response to histamine was also achieved. The values of concentration response - 1 (CR - 1) produced by the macerated extract was also significantly higher than that of chlorpheniramine. These results indicated a competitive antagonistic effect of macerated extract from *N. sativa* on histamine H<sub>1</sub> receptors of tracheal smooth muscles which was greater than that of chlorpheniramine at concentrations used (101). In fact, in a previous study, the relaxant effect of H<sub>1</sub> receptors antagonist, chlorpheniramine on smooth muscle was shown which supported the contribution of H<sub>1</sub> receptors inhibition in the relaxant effect of *N. sativa* on smooth muscles (102).

The inhibitory effect of carvacrol, a constituent of *N. sativa*, on histamine H<sub>1</sub> receptors was also investigated and the results showed parallel and concentration dependent rightward shifts in histamine concentration response curve and maximum response to histamine was achieved in the presence of carvacrol. This finding showed a competitive antagonistic effect for carvacrol on H<sub>1</sub> receptors of tracheal smooth muscles (103).

#### Stimulatory effect on $\beta$ 2- adrenoceptors

The stimulatory effect of aqueous and macerated extracts from *N. sativa* on  $\beta$ 2- adrenoceptors was examined by performing isoprenaline concentration response curves inducing relaxation of precontracted tracheal smooth muscle in the presence of saline, extracts and propranolol (103, 104). While propranolol caused a parallel rightward shift as expected, both extract lead to parallel leftward shift of the curves. These results may suggest a stimulatory effect of both extract from *N.*

*sativa* on  $\beta$ 2- adrenoceptors. The stimulatory effect of carvacrol, the constituent of the plant on  $\beta$ -adrenergic receptors was also evaluated similarly and the results showed concentration dependent leftward shift in isoprenaline concentration response curve obtained in the presence of carvacrol which suggested stimulatory effect of this compound on  $\beta$ -adrenergic receptors (105).

The vasorelaxation effect of cumulative concentration of the hydroethanolic extract of *N. sativa* on rat aortic smooth muscle contracted by both KCl and phenylephrine, a selective  $\alpha_1$ -adrenergic receptor agonist (21) and the relaxant effect of thymoquinone on smooth muscle of pulmonary artery precontracted by phenylephrine (90) indicated the involvement of  $\alpha_1$ -adrenergic receptor blockade in this effect.

#### Other possible mechanisms of the relaxant effect of *N. sativa* on smooth muscles

The other possible mechanisms responsible for relaxant effect of *N. sativa* and its constituents included; stimulation of inhibitory non-adrenergic non-cholinergic nervous system (NANC) or inhibition of stimulatory NANC (106), methylxanthine activity of the plant (107), inhibition of phosphodiesterase (108), nitric oxide production because it was shown that nitric oxide induces airway smooth muscle cell relaxation (109). However, the extract of *N. sativa* caused a dose-dependent decrease in NO production (110). These results indicated that this mechanism cannot contribute in the relaxant effect of the plant on smooth muscles. Non-competitive blocking of serotonin,  $\alpha_1$  and endothelin receptors (90), inhibition of eicosanoid generation, namely thromboxane B<sub>2</sub> and leukotriene B<sub>4</sub> (111), stabilizing of mast-cells and inhibiting the release of histamine (61), inhibition of PGD<sub>2</sub> synthesis and Th<sub>2</sub>-driven immune response (112) also could contribute in the relaxant effect of the plant on smooth muscle. However, the contribution of these mechanisms in the relaxant effect of *N. sativa* should be examined in further studies. Table 3 summarized the possible mechanisms of *N. sativa* and its constituents.

#### Clinical applications

The relaxant effect of *N. sativa* on tracheal smooth muscle could be of therapeutic value in treatment of airway obstruction in obstructive pulmonary diseases such as asthma and chronic pulmonary

**Table 4.** Possible clinical applications of the relaxant effect of *Nigella sativa* and its constituents on different types of smooth muscle

Method	Possible effect	Reference
Respiratory system	Bronchodilatory effects	(46, 49, 84, 113)
	Elevation in FEV1 and FVC values in asthmatic patients	(48, 49)
Cardiovascular system	Antitussive activity	(50, 51)
	Vasodilatory effect	(22-23)
Gastrointestinal tract	Lowering blood pressure and improving of lipid profile	(23)
	Spasmolytic effect	(75-76, 92, 114)
Urogenital tracts	Gastroprotective effect	(35-36)
		(115)

diseases (COPD). In fact, the bronchodilatory effects of the boiled extract of the plant on asthmatic patients (49) as well as its effect on airway of these patients (46) and chemical war victims who had airway obstruction (113) were demonstrated. In one of these studies (49), the boiled extract of *N. sativa* was administered orally and values of pulmonary function tests were measured repeatedly in 30 min interval until 180 min. The results showed increase in all pulmonary function tests values which was similar to the effect of oral theophylline and inhaled salbutamol both from the magnitude and duration of point of view (49). The comparative study also showed that the plant significantly increased the values of forced expiratory volume in one second (FEV1%) and forced vital capacity (FVC) with marked reduction in asthmatic attacks (48). Antitussive activity of this plant and its constituents has also shown in guinea pigs (50, 51). These studies further support the traditional use of black seeds either alone or in combination with honey to treat bronchial asthma (84).

Relaxant effect of black seed on vascular smooth muscle leads to vasodilatory effect which could indicate the therapeutic application of the plant in hypertensive patients (20, 22, 23). In fact, administration of *N. sativa* seed extract for 8 weeks reduced systolic and diastolic blood pressure in a dose-dependent manner and lowered the levels of total and LDL cholesterol in patients with mild hypertension (23).

The relaxant effect of the plant on gastrointestinal smooth muscle was also had been shown (75, 91-93). The aqueous and methanolic extract of *N. sativa* seeds and its constituent, thymoquinone, was showed to have spasmolytic effect which provides a scientific basis for its traditional use in diarrhea (76) and colitis (114). Moreover, both *N. sativa* and thymoquinone can partially protect gastric mucosa from acute alcohol-induced mucosal injury (35) and ischaemia/reperfusion induced gastric lesion (36).

With regard to the relaxant effect of *N. sativa* on urogenital smooth muscle, the plant also may have therapeutic values in the disorders of these systems associated with the obstruction of their tubes. In fact the protective effect of *N. sativa* on the kidney function in rats has been shown (115). Table 4 summarized the possible clinical application of *N. sativa*.

## Conclusion

In conclusion the relaxant effects of *N. sativa* and some of its constituents on trachea, vascular, gastrointestinal and urogenital smooth muscles were discussed in the present review article. The possible mechanisms of the relaxant effect of the plant and its constituents on smooth muscle as well as the clinical applications of this effect were also indicated. However, further clinical studies were needed to demonstrate responsible constituents of the plant, the exact mechanism of action and the clinical applications of this effect more clearly.

## References

1. Aqel M, Shaheen R. Effect of volatile oil of *Nigella sativa* seeds on the uterine smooth muscle of rat and guinea pig. *J Ethnopharmacol* 1996; 52:23-26.
2. Worthen DR, Ghosheh OA, Crooks PA. The *in vitro* anti-tumor activity of some crude and purified components of black seed, *Nigella sativa* L. *Anticancer Res* 1998; 18:1527-1532.
3. Khan MA. Chemical composition and medicinal properties of *Nigella sativa* Linn. *Inflam Pharmacol* 1999; 7:15-35.
4. Ave-Sina. Law in Medicine. Tehran: Ministry of Guidance Publication; 1990.
5. Sayed MD. Traditional medicine in health care. *J Ethnopharmacol* 1980; 2:19-22.
6. Ali BH, Blunden G. Pharmacological and toxicological properties of *Nigella sativa*. *Phytother Res* 2003; 17:299-305.
7. Ramadan MF. Nutritional value, functional properties and nutraceutical applications of black cumin (*Nigella sativa* L.): an overview. *Int J Food Sci Tech* 2007; 42:1208-1218.
8. AL Ameen NM, Altubaigy F, Jahangir T, Mahday IA, Mohammed EA, Musa OA. Effect of *Nigella sativa* and bee honey on pulmonary, hepatic and renal function in Sudanese in Khartoum state. *J Med Plants Res* 2011; 5:6857-6863.
9. Al-Ghamdi MS. The anti-inflammatory, analgesic and antipyretic activity of *Nigella sativa*. *J Ethnopharmacol* 2001; 76:45-48.
10. Mutabagani A, El-Mahdi SM. A study of the anti-inflammatory activity of *Nigella sativa* and thymoquinone in rats. *Saudi Pharmacol J* 1997; 5:110-113.
11. Tekeoglu I, Dogan A, Ediz L, Budancamanak M, Demirel A. Effects of Thymoquinone (volatile oil of black cumin) on rheumatoid arthritis in rat models. *Phytother Res* 2007; 21:895-897.

12. Al-Hader A, Aqel M, Hasan Z. Hypoglycemic Effects of the Volatile Oil of *Nigella sativa* Seeds. *Pharm Biol* 1993; 31:96-100.
13. Meral I, Yener Z, Kahraman T, Mert N. Effect of *Nigella sativa* on glucose concentration, lipid peroxidation, anti-oxidant defence system and liver damage in experimentally-induced diabetic rabbits. *J Vet Med A Physiol Pathol Clin Med* 2001; 48:593-599.
14. Fararh KM, Atoji Y, Shimizu Y, Takewaki T. Insulinotropic properties of *Nigella sativa* oil in streptozotocin plus nicotinamide diabetic hamster. *Res Vet Sci* 2002; 73:279-282.
15. El- Dakhakhny M, Barakat M, El-Halim M, Aly SM. Effects of *Nigella sativa* oil on gastric secretion and ethanol-induced ulcer in rats. *J Ethnopharmacol* 2000; 72:299-304.
16. Hawsawi ZA, Ali BA, Bamosa AO. Effect of *Nigella sativa* (black seed) and thymoquinone on blood glucose in albino rats. *Ann Saudi Med* 2001; 21:242-244.
17. Sultan MT, Butt MS, Karim R, Iqbal SZ, Ahmad S, Zia-Ul-Haq M, et al. Effect of *Nigella sativa* fixed and essential oils on antioxidant status, hepatic enzymes, and immunity in streptozotocin induced diabetes mellitus. *BMC Complement Altern Med* 2014; 14:193.
18. Alimohammadi S, Hobbenaghi R, Javanbakht J, Kheradmand D, Mortezaee R, Tavakoli M, et al. Protective and antidiabetic effects of extract from *Nigella sativa* on blood glucose concentrations against streptozotocin (STZ)-induced diabetic in rats: an experimental study with histopathological evaluation. *Diagn Pathol* 2013; 8:137.
19. Gilani AH, Shaheen F, Shakir T. Thymol lowers blood pressure through blockade of calcium channels. *Fundam Clin Pharmacol* 2001; 15:163-????.
20. Zaoui A, Cherrah Y, Lacaille-Dubois M A, Settaf A, Amarouch H, Hassar M. Diuretic and hypotensive effects of *Nigella sativa* in the spontaneously hypertensive rat. *Therapie* 2000; 55:379-382.
21. Niazmand S, Fereidouni E, Mahmoudabady M, Mousavi SM. Endothelium-Independent vasorelaxant effects of hydroalcoholic extract from *Nigella sativa* seed in rat aorta: The Roles of Ca<sup>2+</sup> and K<sup>+</sup> Channels. *BioMed Res Int* 2014; 2014:247054
22. Najmi A, Nasiruddin M, Khan RA, Haque SF. Indigenous herbal product *Nigella sativa* proved effective as an antihypertensive in metabolic syndrome. *Asian J Pharmaceu Clin Res* 2013; 6:61-64.
23. Dehkordi FR, Kamkhab AF. Antihypertensive effect of *Nigella sativa* seed extract in patients with mild hypertension. *Fundam Clin Pharmacol* 2008; 22:447-452.
24. Peixoto-Nevesa D, Silva-Alvesa KS, Gomesa MDM, Lima FC, Lahloua S, Magalhaes PJC, et al. Vasorelaxant effects of the monoterpenic phenol isomers, carvacrol and thymol, on rat isolated aorta. *Fundam Clin Pharmacol* 2010; 24:341-350.
25. Boskabady MH, Shafei MN, Parsaee H. Effects of aqueous and macerated extracts from *Nigella sativa* on guinea pig isolated heart activity. *Pharmazie* 2005; 60:943-948.
26. Shafei MN, Boskabady MH, Parsaee H. Effect of aqueous extracts from *Nigella sativa* on guinea pig isolated heart. *Ind J Exper Biol* 2005; 43:635-639.
27. Hassanin NI, Hassan FM. A preliminary study on the effect of *Nigella sativa* seeds on hypoglycemia. *Vet Med J Giza* 1996; 44:699-708.
28. Zaoui A, Cherrah Y, Mahassini N, Alaoui K, Amarouch H, Hassar M. Acute and chronic toxicity of *Nigella sativa* fixed oil. *Phytomedicine* 2002; 9:69-74.
29. Ahmad S, Beg ZH. Elucidation of mechanisms of actions of thymoquinone-enriched methanolic and volatile oil extracts from *Nigella sativa* against cardiovascular risk parameters in experimental hyperlipidemia. *Lipids Health Dis* 2013; 12:86-????.
30. Bamosa AO, Ali BA, Sowayan SA. Effect of oral ingestion of *Nigella sativa* seeds on some blood parameters. *Saudi Pharm J* 1997; 5:126-129.
31. Farzaneh E, Rahmani Nia F, Mehrtash M, Mirmoeini FS, Jalilvand M. The effects of 8-week *Nigella sativa* supplementation and aerobic training on lipid profile and VO<sub>2</sub> max in sedentary overweight females. *Int J Prev Med* 2014; 5:210-216.
32. Enomoto S, Asano R, Iwahori Y, Narui T, Okada Y, Singab AN, et al. Hematologic studies on black cumin oil from the seeds of *Nigella sativa*. *Biol Pharm Bull* 2001; 24:307-310.
33. Ibraheim ZZ. Effects of *Nigella sativa* seeds and total oil on some blood parameters in female doxorubicin volunteers. *Saudi Pharm. J.* 2002; 10:54-59.
34. Raj Kapoor B, Anandan R, Jayakar B. Anti-ulcer effect of *Nigella sativa* against gastric ulcers in rats. *Current Science* 2002; 82:177-185.
35. Kanter M, Demir H, Karakaya C, Ozbek H. Gastroprotective activity of *Nigella sativa* L oil and its constituent, thymoquinone against acute alcohol-induced gastric mucosal injury in rats. *World J Gastroenterol* 2005; 11:6662-6666.
36. El-Abhar HS, Abdallah DM, Saleh S. Gastroprotective activity of *Nigella sativa* oil and its constituent, thymoquinone, against gastric mucosal injury induced by ischemia/reperfusion in rats. *J Ethnopharmacol* 2003; 84:251-258.
37. Kanter M, Coskun O, Uysal H. The antioxidative and antihistaminic effect of *Nigella sativa* and its major constituent, thymoquinone on ethanol-induced gastric mucosal damage. *Arch Toxicol* 2006; 80:217-224.
38. El-Sayed MM, El-Banna HA, Fathy FA. The use of *N. Sativa* oil as a natural preservative agent in processed cheese spread. *Egypt J Food Sci* 1994; 22:381-396.
39. Toma C, Simu GM, Hanganu D, Olah N, Vata FM G, Hammami C, et al. Chemical composition of the tunisian *Nigella sativa*. Note I. profile on essential oil. *Farmacia* 2010; 58:458-464.
40. Badar El-Din MK. Anti-asthmatic activity of the active principle of *Nigella sativa* "Nigellone". *Gazette Egypt Paediatr Assoc.* 1960; 8:864-867.
41. Mahfouz M, El-Dakhakhany M. Some chemical and pharmacological properties of the new antiasthmatic drug "Nigellone". *Egypt Pharmacol Bull* 1960; 6:357-360.
42. Chakravarty N. Inhibition of histamine release from mast cells by nigellone. *Ann Allergy* 1993; 70:237-242.
43. El-Tahir KE, Ashour MM, Al-Harbi MM. The cardiovascular actions of the volatile oil of the black seed (*Nigella sativa*) in rats: elucidation of the mechanism of action. *Gen Pharmacol* 1993; 24:1123-1131.
44. EL-Dakhakhany M. Some pharmacological properties of some constituents of *Nigella sativa* L seeds: The carbonyl fraction of essential oil. *Proceeding of the 2nd International conference on Islamic Medicine.* Kuwait 1982; 426-431.



45. Chakarvarti N. Inhibition of histamine release from mast cells by nigellone. *Ann Allergy* 1993; 70:237-242.
46. Boskabady MH, Javan H, Sajadi M, Rakhshandah H. The possible prophylactic effect of *Nigella sativa* seed extract in asthmatic patients. *Fundam Clin Pharmacol* 2007; 21:559-566.
47. Ahmad J, Ali Khan R, Malik MA. A study of *Nigella sativa* oil in the management of wheeze associated lower respiratory tract illness in children. *African J Pharmacy Pharmacol* 2010; 4:436-439.
48. Al-Jawad FH, Al-Razzuqi RAM, Hashim HM, Ismael AH. Broncho-relaxant activity of *Nigella sativa* versus anthemisnobilis in chronic bronchial asthma; a comparative study of efficacy. *IOSR J Pharmacol* 2012; 2:81-83.
49. Boskabady MH, Mohsenpoor N, Takaloo L. Antiasthmatic effect of *Nigella sativa* in airways of asthmatic patients. *Phytomedicine* 2010; 17:707-713.
50. Boskabady MH, Kiani S, Jandaghi P, Ziaei T, Zarei A. Antitussive effect of *Nigella sativa* in guinea pigs. *Pak J Med Sci* 2004; 20:224-228.
51. Hosseinzadeh H, Eskandari M, Ziaee T. Antitussive effect of thymoquinone, a constituent of *Nigella sativa* seeds, in guinea pigs. *Pharmacologyonline* 2008; 2:480-484.
52. Boskabady MH, Keyhanmanesh R, Khamneh S, Ebrahimi Saadatloo MA. The effect of *Nigella sativa* extract on tracheal responsiveness and lung inflammation in ovalbumin sensitized guinea pigs. *Clinics* 2011; 66:879-887.
53. Boskabady MH, Keyhanmanesh R, Khamneh S, Doostar Y, Khakzad MR. Potential immunomodulation effect of the extract of *Nigella sativa* on ovalbumin sensitized guinea pigs. *J Zhejiang Univ-Sci B (Biomed & Biotechnol)* 2011; 12:201-209.
54. Keyhanmanesh R, Boskabady MH, Khamneh S, Ebrahimi MA. The effect of thymoquinone, the main constituent of *Nigella sativa* on tracheal responsiveness and WBC count in lung lavage of sensitized guinea-pigs. *Planta Med* 2010; 76:218-222.
55. Keyhanmanesh R, Boskabady MH, Khamneh S, Doostar Y. Effect of thymoquinone on lung pathology and cytokine levels of ovalbumin-sensitized guinea-pigs. *Pharmacol Rep* 2010; 62:910-916.
56. Abd El Aziz AE, El Sayed NS, Mahran LG. Anti-asthmatic and anti-allergic effects of thymoquinone on airway-induced hypersensitivity in experimental animals. *J Appl Pharm Sci* 2011; 1:109-117.
57. Keyhanmanesh R, Nazemiyeh H, Mazouchian H, Bagheri Asl MM, ShoarMK, Alipour MR, et al. *Nigella sativa* pretreatment in guinea pigs exposed to cigarette smoke modulates in vitro tracheal responsiveness. *Iran Red Crescent Med J* 2014; 16:e10421.
58. Boskabady MH, Jalali S. Effect of carvacrol on tracheal responsiveness, inflammatory mediators, total and differential WBC count in blood of sensitized guinea pigs. *Exp Biol Med* 2013; 238:200-208.
59. Al-Naqeeb G, Ismail M, Al-Zubairi AS. Fatty acid profile, tocopherol content and total antioxidant activity of oil extracted from *Nigella sativa* seeds. *Int J Pharmacol* 2009; 5:244-250.
60. Mezit A, Meziti H, Boudiaf K, Mustapha B, Bouriche H. Polyphenolic profile and antioxidant activities of *Nigella sativa* seed extracts in vitro and in vivo. *World Academy of Science, Engin Tech* 2012; 64:24-32.
61. Gilani AH, Jabeen Q, Ullah Khan MA. A review of medicinal uses and pharmacological activities of *Nigella sativa*. *Pak J Biol Sci* 2004; 7:441-451.
62. Nergiz C, Otles S. Chemical composition of *Nigella sativa* L. seeds. *Food Chem* 1993; 48:259-261.
63. Randhawa MA, Al-Ghamdi MS. A review of the pharmaco-therapeutic effects of *Nigella sativa*. *Pak J Med Res.* 2002; 41:77-83.
64. Menounos P, Staphylakis K, Gegiou D. The sterols of *nigella sativa* seed oil. *Phytochemistry* 1986; 25:761-763.
65. Nickavar B, Mojab F, Javidnia K, Roodgar Amolia MA. Chemical composition of the fixed and volatile oils of *Nigella sativa* L. from Iran. *Z Naturforsch C* 2003; 58:629-631.
66. Cheikh-Rouhou S, Besbes S, Hentati B, Blecker C, Deroanne C, Attia H. *Nigella sativa* L.: Chemical composition and physicochemical characteristics of lipid fraction. *Food Chem* 2007; 101:673-681.
67. Ghosheh OA, Houdi AA, Crooks PA. High performance liquid chromatographic analysis of the pharmacologically active quinones and related compounds in the oil of the black seed (*Nigella sativa* L.). *J Pharm Biomed Anal* 1999; 19:757-762.
68. Hosseinzadeh H, Parvardeh S. Anticonvulsant effects of thymoquinone, the major constituent of *Nigella sativa* seeds in mice. *Phytomedicine* 2004; 11:56-64.
69. Burits M, Bucar F. Antioxidant activity of *Nigella sativa* essential oil. *Phytother Res* 2000; 14:323-328.
70. Ur-Rahman A, Malik S, Cun-heng H, Clardy J. Isolation and structure determination of nigellidine, a novel alkaloid from the seeds of *Nigella sativa*. *Tetrahedron Lett* 1985; 23:2759-2762.
71. Ur-Rahman A, Malik S, Hasan SS, Choudhary MI, Ni C-Z, Clardy J. Nigellidine, a new indazole alkaloid from the seeds of *Nigella sativa*. *Tetrahedron Lett* 1995; 36:1993-1996.
72. Merfort I, Wray V, Barakat HH, Hussein SAM, Nawwar MAM, Willuhn G. Flavonol triglycosides from seeds of *Nigella sativa*. *Phytochemistry* 1997; 46:359-363.
73. Malik S, Zaman K. Nigellimine, a new isoquinoline alkaloid from the seeds of *Nigella sativa*. *J Nat Prod* 1992; 55:676-678.
74. Ur-Rahman A, Malik S, Ahmed S, Choudhary MI, Ur-Rehman H. Nigellimine-N-oxide- a new isoquinoline alkaloid from the seeds of *Nigella sativa*. *Heterocycles* 1985a; 23:953-955.
75. Reiter M, Brandt W. Relaxant effects on tracheal and ileal smooth muscles of the guinea-pig. *Arzneimittelforschung* 1985; 35:408-414.
76. Gilani AH, Aziz N, Khurram IM, Chaudhary KS, Iqbal A. Bronchodilator, spasmolytic and calcium antagonist activities of *Nigella sativa* seeds (Kalonji): a traditional herbal product with multiple medicinal uses. *J Pak Med Assoc* 2001; 51:115-120.
77. Boskabady MH, Shahabi M. Bronchodilatory and anticholinergic effects of *Nigella sativa* on isolated guinea-pig tracheal chains. *Iran J Med Sci* 1997; 22:127-133.
78. Boskabady MH, Shirmohammadi B, Jandaghi P, Kiani S. Possible mechanism(s) for relaxant effect of aqueous and macerated extracts from *Nigella sativa* on tracheal chains of guinea pig. *BMC Pharmacol* 2004; 4:3.

79. Boskabady MH, Shirmohammadi B. Effect of *Nigella sativa* on isolated guinea pig tracheal chains. Arch Iran Med 2002; 5:103-107.
80. Boskabady MH, Keyhanmanesh R, Ebrahimi Saadatloo MA, Khamnei S. The contribution of water and lipid soluble substances in the relaxant effects of *Nigella sativa* on guinea pig tracheal chains. Iran J Basic Med Sci 2007; 10:154-161.
81. Boskabady MH, Keyhanmanesh R, Ebrahimi Saadatloo MA. Relaxant effects of different fractions from *Nigella sativa* on guinea pig tracheal chains and its possible mechanism(s). Ind J Exp Biol 2008; 46:805-810.
82. Keyhanmanesh R, Bagban H, Nazemiyeh H, Babil FM, Alipour MR, Ahmady M. The Relaxant effects of different methanolic fractions of *Nigella sativa* on guinea pig tracheal chains. Iran J Basic Med Sci 2013; 16:123-128.
83. Keyhanmanesh R, Bagban H, Nazemieh H, Babil FM, Alipour MR. The main relaxant constituents of *Nigella sativa* methanolic fraction on guinea pig tracheal chains. Iran J Allergy Asthma Immunol 2013; 12:136-143.
84. Al-Majed AA, Daba MH, Asiri YA, Al-Shabanah OA, Mostafa AA, El-Kashef HA. Thymoquinone-induced relaxation of guinea-pig isolated trachea Res Commun Mol Pathol Pharmacol 2000; 110:333-345.
85. Boskabady MH, Aslani M R. Possible relaxant effects of thymoquinone on guinea pig tracheal chains. Iran Biomed J 2005; 9:123-128.
86. Silva YMS, Silva MTA, Sampaio PA, Quintans JSS, Quintans-Junior LJ, Ribeiro LAA. Relaxant effect of carvacrol, citronellal and p-cymene, monoterpenes present in thymus and cymbopogon species, in guinea pig trachea: A comparative study. J Med Plant Res 2014; 8:881-888.
87. Boskabady MH, Jafari Z, Pouraboli I. The effect of carvacrol on muscarinic receptors of guinea-pig tracheal chains. Phytother Res 2011; 25:530-535.
88. Boskabady MH, Jandaghi P. Relaxant effects of carvacrol on guinea pig tracheal chains and its possible mechanisms. Pharmazie 2003; 58:661-663.
89. Agarwal R, Kharya MD, Shrivastava R. pharmacological studies of essential oil and unsaponifiable matter of seeds of *Nigella sativa*. Indian J Pharm Sci 1979; 41:248.
90. Suddek GM. Thymoquinone-induced relaxation of isolated rat pulmonary artery. J Ethnopharmacol 2010; 127:210-214.
91. Aqel MB. Effect of *Nigella sativa* seeds on intestinal smooth muscles. Int J Pharmacog 1993; 31:55-60.
92. Parvardeh S, Fatehi M. Inhibitory effects of thymoquinone, the major component of *Nigella sativa* L. seeds, on spontaneous and evoked contractions of guinea pig isolated ileum. J Med Plants 2007; 6:29-39.
93. Ghayur MN, Gilani AH, Janssen L J. Intestinal, airway, and cardiovascular relaxant activities of thymoquinone. Evid Based Complement Alternat Med 2012; 2012:305319.
94. Lohn M, Furstenu M, Sagach V, Elger M, Schulze W, Luft FC, et al. Ignition of calcium sparks in arterial and cardiac muscle through caveolae. Circul Res 2000; 87:1034-1039.
95. Buckle DR, Arch JRS, Boering N E, Foster KA, Taylor JF, Taylor SG, et al. Relaxation effect of potassium channel activators BRL 38227 and pinacidil on guinea pig and human airway smooth muscle, and blockade of their effects by glibenclamide and BRL. Pulmon Pharmacol 1993; 6:77-86.
96. Nelson MT, Quayle J M. Physiological roles and properties of potassium channels in arterial smooth muscle. American J Physiol: Cell Physiol 1995; 268:799-822.
97. Cortes SF, Rezende BA, Corriu C, Medeiros IA, Teixeira MM, Lopes MJ, et al. Pharmacological evidence for the activation of potassium channels as the mechanism involved in the hypotensive and vasorelaxant effect of dioclein in rat small resistance arteries. Br J Pharmacol 2001; 133:849-858.
98. Jackson WF. Ion channels and vascular tone. Hypertension 2000; 35:173-178.
99. Bogats LN, Epshtein MM. Effect of  $\alpha$ -pinene on the chemoreceptor function of the peripheral nerves. Fiziol. Zhur Akad Nauk Rse 1959; 5:659-661.
100. El-Tahir KEH, Ashour MM, Al-Harbi MM. The respiratory effect of the volatile oil of the black seed (*Nigella sativa*) in guinea-pig: elucidation of the mechanism(s) of action. Gen Pharmacol 1999; 24:1115-1122.
101. Boskabady MH, Shiravi N. Inhibitory effect of *Nigella sativa* on histamine (H1) receptors of isolated guinea pig tracheal chains. Pharm Biol 2002; 40:596-602.
102. Popa VT. Bronchodilatory activity of an H1 blocker chlorpheniramine. J Allergy Clin Immunol 1977; 59:54-63.
103. Boskabady MH, Kiani S, Jandaghi P. Stimulatory effect of *Nigella sativa* on  $\beta$ 2-adrenoceptors of guinea pig tracheal chains. Medical J Islam Repub Iran 2007; 18:153-158.
104. Boskabady MH, Tabanfar H, Gholamnezhad Z, Sadeghnia HR. Inhibitory effect of *Zataria multiflora* Boiss and carvacrol on histamine (H1) receptors of guinea-pig tracheal chains. Fundam Clin Pharmacol 2012; 26:609-620.
105. Boskabady MH, Kaveh M, Eftekhar N, Nemati A. *Zataria multiflora* Boiss and Carvacrol Affect  $\beta$ 2-Adrenoceptors of Guinea Pig Trachea. Evid Based Complement Alternat Med 2011; 2011:857124.
106. Linden A, Lofdahl CG, Ullman A, Skoogh BE. Non-adrenergic, noncholinergic responses stabilize smooth muscle tone with and without parasympathetic activation in guinea pig isolated airways. Eur Respir J 1993; 6:425-433.
107. Meini S, Ballati L, Evangelista S, Manzini S. Isbufulline, a xanthine derivative, inhibits bronchoconstrictor responses produced by stimulation of capsaicin sensitive nerves in guinea pig, *in vitro* and *in vivo* evidence. Pulm Pharmacol 1993; 6:279-286.
108. Van Amsterdam RGM, Meurs H, Brouwer F, Postema JB, Timmermans A, Zaagsma J. Role of phosphoinositol metabolism in functional antagonism of airway smooth muscle contraction by beta-adrenoceptor agonist. Eur J Pharmacol 1989; 172:175-183.
109. Perez-Zoghbi JF, Bai Y, Sanderson MJ. Nitric oxide induces airway smooth muscle cell relaxation by decreasing the frequency of agonist-induced  $Ca^{2+}$  oscillations. J Gen Physiol 2010; 135:247-259.
110. Mahmood MS, Gilani AH, Khwaja A, Rashid A, Ashfaq MK. The *in vitro* effect of aqueous extract of

*Nigella sativa* seeds on nitric oxide production. *Phytother Res* 2003; 117:921-924.

111. Houghton PJ, Zarka R, de las Heras B, Hault JR. Fixed oil of *Nigella sativa* and derived thymoquinone inhibit eicosanoid generation in leukocytes and membrane peroxidation. *Planta Medica* 1995; 61:33-36.

112. El-Mezayen R, El-Gazzar M, Nicolls MR, Marecki JC, Dreskin SC, Nomiya H. Effect of thymoquinone on cyclooxygenase expression and prostaglandin production in a mouse model of allergic airway inflammation. *Immunol Lett* 2006; 106:72-81.

113. Boskabady MH, Farhadi F. The possible prophylactic effect of *Nigella sativa* seed extract on respiratory symptoms, pulmonary function tests on chemical war victims. *J Alt Comple Med* 2008; 14:1137-1144.

114. Mahgoub AA. Thymoquinone protects against experimental colitis in rats. *Toxicol Lett* 2003; 143:133-143.

115. Dollah MA, Parhizkar S, Izwan M. Effect of *Nigella sativa* on the kidney function in rats. *Avicenna J Phytomed* 2013; 3:152-158.

Archive of SID