

Screening of DFNB3 in Iranian families with autosomal recessive non-syndromic hearing loss reveals a novel pathogenic mutation in the MyTh4 domain of the *MYO15A* gene in a linked family

Somayeh Reisi¹, Mohammad Amin Tabatabaiefar², Mohammad Hosein Sanati³, Morteza Hashemzadeh Chaleshtori^{4*}

¹ Department of Genetics, Faculty of Basic Sciences, University of Shahrekord, Shahrekord, Iran

² Medical Genetics Dept., Isfahan University of Medical Sciences, Isfahan, Iran

³ Medical Genetics Dept., National Institute of Genetic Engineering and Biotechnology (NIGEB), Tehran, Iran

⁴ Cellular and Molecular Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

ARTICLE INFO

Article type:

Original article

Article history:

Received: Nov 27, 2015

Accepted: Apr 28, 2016

Keywords:

Hearing loss

Iran

Linkage analysis

MYO15A

mutation

ABSTRACT

Objective(s): Non-syndromic sensorineural hearing loss (NSHL) is a common disorder affecting approximately 1 in 500 newborns. This type of hearing loss is extremely heterogeneous and includes over 100 loci. Mutations in the *GJB2* gene have been implicated in about half of autosomal recessive non-syndromic hearing loss (ARNSHL) cases, making this the most common cause of ARNSHL. For the latter form of deafness, most frequent genes proposed include *GJB2*, *SLC26A4*, *MYO15A*, *OTOF*, and *CDH23* worldwide.

Materials and Methods: The aim of the present study was to define the role and frequency of *MYO15A* gene mutation in Iranian families. In this study 30 Iranian families were enrolled with over three deaf children and negative for *GJB2*. Then linkage analysis was performed by six DFNB3 short tandem repeat markers. Following that, mutation detection accomplished using DNA sequencing.

Results: One family (3.33%) showed linkage to DFNB3 and a novel mutation was identified in the *MYO15A* gene (c.6442T>A) as the disease-causing mutation. Mutation co-segregated with hearing loss in the family but was not present in the 100 ethnicity-matched controls.

Conclusion: Our results confirmed that the hearing loss of the linked Iranian family was caused by a novel missense mutation in the *MYO15A* gene. This mutation is the first to be reported in the world and affects the first MyTH4 domain of the protein.

► Please cite this article as:

Reisi S, Tabatabaiefar MA, Sanati MH, Hashemzadeh Chaleshtori M. Screening of DFNB3 in Iranian families with autosomal recessive non-syndromic hearing loss reveals a novel pathogenic mutation in the MyTh4 domain of the *MYO15A* gene in a linked family. Iran J Basic Med Sci 2016; 19:772-778.

Introduction

Hearing loss (HL) is the most common sensory disorder estimated to affect 70 million people worldwide. This disorder is mostly a birth defect that affects 1 in 500 newborns (1). HL is extremely heterogeneous and can result from genetic or environmental factors or both (2). More than 50% of hearing loss cases are caused by genetic factors, of which 70% are non-syndromic (NSHL) and the remaining 30% are attributed to syndromic forms (3). Different loci have been identified for NSHL and have been named according to their mode of inheritance (DFNB for autosomal recessive, DFNA for autosomal dominant, and DFN for X-linked) and the time of their identification (ex. DFNB1-DFNB95) (4).

Over 60 genes have been discovered causing NSHL (Hereditary hearing loss homepage) (5). Studies have indicated that *GJB2* gene mutations are a major cause of ARNSHL. The contribution of other loci to the ARNSHL is much lower. The extreme heterogeneity of HL makes it cumbersome to identify the genetic cause of this disease in single families leading to difficulties in their genetic counseling and testing. Despite previous intensive GLA and candi-date gene screening, a large proportion of ARNSHL remains genetically unexplained (5, 6). Due to the high heterogeneity of HL, researchers have suggested studying of large families in populations like middle east in which consanguineous marriage rate is high (7). Iran, with specific population characteristics

*Corresponding author: Morteza Hashemzadeh Chaleshtori, Cellular and Molecular Research Center, School of Medicine, Shahrekord University of Medical Sciences, Shahrekord, Iran. Tel: +98 381 3335654, Fax: +98 381 3330709, email: mchalesh@yahoo.com

such as high consanguineousmarri-age rate and heterogeneous population, offers a good opportunity to study rare autosomal recessive disorders including ARNSHL. Studies in Iran showed that contribution proportion of *GJB2* mutations in HL in different populations varies ranging from approximately 27%–38% in the north to 0%–0.4% in the southeast(8-10). So far, the contribution of mutations in other genes associated with HL has been identified only in a limited number of Iranian families and it seems in order to determine a more exact contribution of each one of these genes in Iranian families, more extensive studies are required to be done. For ARNSHL, the most frequent causative genes in order of frequency are *GJB2*, *SLC26A4*, *MYO15A*, *OTOF*, and *CDH23*. Mutations in the *SLC26A4* gene, which is located at the *DFNB4* locus are the second most frequent cause of ARNSHL and in order of frequency after *GJB2* and *SLC26A4*, mutations in *MYO15A* is the third most frequent cause of ARNSHL in the world (11). Forty-three mutations have been reported in *MYO15A*, most of which have been found using GLA in consanguineous families from specific countries (12-15).

MYO15A encodes an unconventional myosin XVa, which has a role in stereocilia formation (16). Myosins are molecular motor proteins that drive the moment of actin filaments via ATP hydrolysis to facilitate muscle contraction, organelle trafficking, cell movement, cytokinesis, and signal transduction (17). The protein is unique among unconventional myosins in that it includes a long N-terminal domain (coded by exon 2) that is alternatively spliced to generate distinct class 1 and class 2 protein isoforms. The N-terminal domain is required for normal hearing, as premature stop mutations that result in loss of this domain cause *DFNB3* hearing loss (18, 19). Myosin XVa also contains domains that are conserved within the myosin protein family, including the motor domain, IQ motifs(calmodulin/myosin light chain binding), MyTh4 domains (myosin tail like homology region 4), FERM motifs (4.1 protein, Ezrin, Radixin, and Moesin), SH3 domain (Src homology 3), and the PDZ ligand domain. Many missense and nonsense mutations in coding exons of the motor domain, FERM, and MyTh4 domain have been identified which cause profound HL in different populations (20-22).

In this study, we have characterized an ARNSHL family from the west of Iran and identified a novel mutation in the *MYO15A* gene. This novel mutation has been identified in the highly conserved MyTh4 domain in this family.

Materials and Methods

Sampling and genomic DNA extraction

In this descriptive laboratory study, 30 Iranian families with at least 2 affected children were

selected, informational questionnaires were filled out by family members, and clinical assessments were performed, also consent was obtained from all family members. This research has been approved by the National Institute of Genetic Engineering and Biotechnology and Cellular and Molecular Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran. All cases of syndromic hearing loss caused by environmental factors such as ototoxic medications, hepatitis, head trauma, or meningitis were excluded. All families were informed and written consent was taken and 5 ml peripheral blood, in tubes containing 0.5 M EDTA from all available members of all families was obtained. Genomic DNA was extracted from blood samples using the phenol-chloroform standard method. DNA quality (purity and DNA concentration) was checked by spectroscopy (UNICO 2100, USA) (23, 24).

GJB2 mutation screening

At least one affected member from every pedigree was selected for detection of *GJB2* mutations and reaction was carried out in accordance with previous protocols (25). PCR product was run in an 8% polyacrylamide gel electrophoresis (PAGE) at 45 mA for 1.5 hr. DNA bands were visualized by silver staining. A single PCR product of 809 bp was obtained and DNA sequencing of the PCR product was carried out in order to detect any change in the gene.

S-LINK analysis, *DFNB3* STR marker, and linkage analysis

For S-Link and LOD score calculation, we used the Easy linkage plus (ver. 5.05) genetic software (26). For S-LINK calculation we used FastSlink version 2.51. Two-point and multipoint parametric LOD scores were calculated using Superlink (ver. 1.6) and GeneHunter (ver. 2.91), respectively. For LOD score calculations using these software packages, inheritance pattern of autosomal recessive, complete penetrance, and disease allele frequency of 0.001 were assumed. HaploPainter version 029.5 software package was used for reconstruction of haplotypes (27). Negative subjects for *GJB2* mutations were selected for linkage analysis. For GLA, 6 different STR markers were used. Upon encountering an uninformative marker, further markers were examined. Table 1 summarizes the markers used in the study and their general characteristics. The criteria for selecting these markers are as following: greater heterogeneity values, shorter fragments in amplicon length and lying close to the known locus. STR markers were selected based on their physical distance found at NCBI UniSTS, and Touchdown program was used for markers. Thermal cycling conditions for amplifying markers were in accordance with previous protocols (25).

Table 1. Genetic STR markers used in this study and their characteristics

Heterozygosity	Size (bp)	Reverse primer	Forward primer	STR
0.72	169-185	GGCCACCATAATCATGTCAGACAAT	GGCCACCATAATCATGTCAGACAAT	D17S921
0.70	159-203	GAGAATCACCTGAACCCG	AATTCAAAGGCTAAAAGCAAAC	D17S1843
0.76	119-131	AAGGGCTTGCTTTGAC	ACTATCCGCCCAATACA	D17S953
0.62	177-187	TGCCTAAACTGCTTTCAGGTGAG	TGCACAGGCCAATTCCTTAC	D17S1857
0.71	103-151	TACATTTAATGCAGGATGCC	CTCTTTGTGCTTGGCAGGTT	D17S740
0.81	139-163	ATATTTCAATATTGTAACCAGTCCC	CCAACATCTAGAATTAATCAGAATC	D17S2196

Mutation screening for the MYO15A gene

Sixty-six exons of the MYO15A gene [NM_016239.3] and flanking regions were amplified using designed primers by the Oligo (ver. 7.05) software. Reaction conditions for amplifying exons in 50 µl was as follows: 2 µl MgCl₂ (50 mM), 2.5 µl PCR buffer (10X), 0.5 µl of each of forward and reverse primers (50 PM), 0.5 U Taq DNA polymerase (5U/ul), 1 µl dNTP mix (10 mM) and 1 µl genomic DNA (about 100 ng). PCR was done as follows: 95 °C for primary denaturation for 5 min; 35 cycles of 94 °C denaturation for 1 min, 59 °C annealing temperature for 1 min, 72 °C extension for 1 min, and a final extension at 72 °C for 5 min. In order to detect any variant in the gene sequence, direct sequencing of amplified exons was performed bi-directionally using ABI 3730XL.

Pathogenicity investigation of novel variant


Co-segregation study in family members was accomplished and investigation of conservative amino acid coding of novel variant confirmed the presence of mutation. The absence of new variation in 100 healthy control samples also confirmed the mutation performed by restriction fragment length polymorphism (RFLP) using *MscI* restriction enzyme. In the current study, we used the following primers: F-5'ATTTATATGGGCAGGGGCAC3' and R-5'

AGGCGGCCAGCAGTGGCC 3' for the amplification.

Results

Most of the subjects in this study exhibited severe to profound bilateral sensorineural hearing loss. 70% of families in this study were consanguineous families. According to the information in the pedigree, the type of hearing loss was autosomal recessive non-syndromic.

Mutation 35delG was identified in 5 families out of 30 families by sequencing of the coding region of the *GJB2* gene. These families were homozygous for this mutation and were eliminated from further analysis. The remaining families were further analyzed for the linkage to DFNB1 using 3 informative markers. However, linkage analysis could not find any other family linked to the DFNB1 locus. For linkage analysis to other loci, first S-LINK values were calculated for the remaining families. 14 families had S-LINK values ≥ 3 , 5 families were of S-LINK values < 2 , and the rest of families had S-LINK values 2 to 3. GLA for the DFNB3 locus was carried out by 6 STR markers. By haplotype analysis, out of the 25 remaining families, one family was found to be linked to this locus and this linked family displayed complete pattern of linkage (Figure 1).

**Figure 1.** Pedigree and haplotypes of the family linked to DFNB3. The order of markers is based on the Marshfield map




Figure 2. Audiograms for a proband member (18 in Figure 1) of a family linked to DFNB3. “O” indicates air conduction for the right ear, while “X” indicates air conduction for the left ear. Audiograms of the affected individual had shown severe to profound hearing loss at two different time intervals

Audiogram analysis of this family confirmed HL to be profound in all affected subjects of the family (Figure 2). Calculated multipoint LOD score for this family was 3.15, which confirmed the linkage to the locus. DNA sequencing of 66 coding exons of the *MYO15A* gene demonstrated a new variant in the homozygous state in exon 30. We first analyzed all polymorphisms of the gene from all data banks. This novel variant was a missense substitution and caused the substitution of nucleotide T to A in the location 6442 of the coding region of *MYO15A* (c.6442T>A) and substitution of tryptophan for

arginine at residue 2148 (p.Trp2148Arg). The ConSurf web server calculated the conservative score for the residue to be 8, which implies it as highly conserved among diverse species (Figure 3). Co-segregation showed that variant in affected subjects and in parents was in homozygous and heterozygous states, respectively. One normal individual was homozygous for the normal allele (Figure 4). We analyzed the variant in 100 normal controls and no case was found to be positive for it. Thus, the observations would confirm the pathogenicity of the novel variant.



Figure 3. conservity of altered amino acids in different species. Calculated score for amino acid was 8




Figure 4. Chromatogram results of novel *MYO15A* variants in an Iranian family
 A: normal alleles. B: A normal subject with heterozygous c.6442T>A allele. C: A patient with homozygous c.6442T>A allele

Discussion

In the present study, we analyzed 30 families with ARNSHL and found one family (~3%) to be linked to DFNB3 by genetic linkage analysis (GLA). After DNA sequencing of the *MYO15A* gene, a novel variant (c.6442T>A, p.Trp2148Arg) was identified in the first MyTH4 domain. *MYO15A* with 66 coding exons encodes an unconventional myosin (myosin XV) that is expressed in the cochlea (19). This protein has important roles in the differentiation and elongation of the inner ear hair cell stereocilia, and it is also necessary for actin organization in hair cells (28). Mutations that cause hearing loss were first identified at the DFNB3 locus, in residents of a village in Indonesia. It was estimated that mutation frequencies were 9% among the inbred population. Since then, many mutations have been reported from different countries such as Pakistan, India, Turkey, Indonesia, and Brazil, interestingly, all of which belong to non-Caucasian populations (13, 14). Recently two compound heterozygote variants have been reported in Korean and Chinese families (15, 29). In addition, a novel homozygous mutation (c.9316dupC) in second MyTH4 of the *MYO15A* gene was identified by exome sequencing and Sanger sequencing in a Chinese family with ARNSHL (30). The novel mutation, found in this study, is located in the first domain of myosin tail homology4 (MTH4). To date, 7 mutations have been identified in this domain that are missense mutations. The MyTH4 domain of myosin has some roles in microtubule binding, as well as in actin binding at the plasma membrane (15, 31). Some studies suggest that MyTH4/FERM domain in MYOXVA is essential for its localization to stereocilia tips, although, their specific function is not been clearly known (18, 32, 33). MyTH4 domain contains a high percentage of

conserved amino acids relative to other domains of the protein(33). Therefore, mutations in this domain can have a significant effect on proper function and structure of the protein. The c.6442T>A mutation leads to substitution of tryptophan for arginine amino acid at codon 2148. Tryptophan is a highly hydrophobic amino acid with aromatic structure, while arginine is a hydrophilic and positive charged amino acid. This amino acid substitution interferes with protein/microtubules interaction and actin and consequently disrupts the normal auditory function. MyTH4 domain interacts with PDZ domain of whirlin protein which is a cytoskeletal scaffold protein and is essential for normal auditory function. The colocalization of MYOXVA and whirlin proteins is essential for the assembly of actin microfilaments at stereocilia tips. In addition to the change in protein structure, this mutation interferes with the interaction between these two proteins and prevents the formation of actin microfilaments that is required for normal hearing. Different studies on Iranian populations show that mutations in *GJB2* and *SLC26A4* are the first and second hearing loss causing mutations, respectively(34). To date, up to 10 mutations in the *MYO15A* gene have been identified in the Iranian population. Thus, according to the current studies the DFNB3 locus in hearing loss might rank third after *GJB2* and *SLC26A4*.

Conclusion

We have identified a novel mutation in the *MYO15A* gene in an Iranian family. This mutation affects the first MyTH4 domain of the protein. This region is highly conserved and should be investigated in future studies.

Acknowledgment

We take this opportunity to express our special thanks and appreciation to all staff members of the National Institute of Genetic Engineering and Biotechnology, Tehran, Iran, as well as Shahrekord Medical University, Shahrekord, Iran. We also would like to sincerely thank the kind participants and their families for their cooperation in this research.

References

- Morton CC, Nance WE. Newborn hearing screening—a silent revolution. *N Engl J Med* 2006; 354:2151-2164.
- Collin RW, Kalay E, Oostrik J, Caylan R, Wollnik B, Arslan S, *et al.* Involvement of DFNBS9 mutations in autosomal recessive nonsyndromic hearing impairment. *Hum Mutat* 2007; 28:718-723.
- Van Camp G, Willems PJ, Smith R. Nonsyndromic hearing impairment: unparalleled heterogeneity. *Am J Hum Genet* 1997; 60:58.
- Willems PJ. Genetic causes of hearing loss. *N Engl J Med* 2000; 342:1101-1109.
- Kenneson A, Van Naarden Braun K, Boyle C. GJB2 (connexin 26) variants and nonsyndromic sensorineural hearing loss: a HuGE review. *Genet Med* 2002; 4:258-274.
- Primignani P, Trotta L, Castorina P, Lalatta F, Sironi F, Radaelli C, *et al.* Analysis of the GJB2 and GJB6 genes in Italian patients with nonsyndromic hearing loss: frequencies, novel mutations, genotypes, and degree of hearing loss. *Genet Test Mol Biomarkers* 2009; 13:209-217.
- Strachan T, Read A. *Human molecular genetics*. New York: Garland science; 2011.
- Chaleshtori MH, Farhud D, Patton M. Congratulation to Margaret Chan Familial and Sporadic GJB2-related deafness in Iran: review of gene mutations. *Iran J Public Health*, 2007; 36:1-14.
- Mahdieh N, Rabbani B, Shirkavand A, Bagherian H, Movahed ZS, Fouladi P, *et al.* Impact of consanguineous marriages in GJB2-related hearing loss in the Iranian population: a report of a novel variant. *Genet Test Mol Biomarkers* 2011; 15:489-493.
- Najmabadi H, Nishimura C, Kahrizi K, Riazalhosseini Y, Malekpour M, Daneshi A, *et al.* GJB2 mutations: passage through Iran. *Am J Med Genet Part A* 2005; 133:132-137.
- Hilgert N, Smith RJ, Van Camp G. Forty-six genes causing nonsyndromic hearing impairment: which ones should be analyzed in DNA diagnostics? *Mutat Res Rev Mutat Res* 2009; 681:189-196.
- Bashir R, Fatima A, Naz S. Prioritized sequencing of the second exon of *MYO15A* reveals a new mutation segregating in a Pakistani family with moderate to severe hearing loss. *Eur J Med Genet* 2012; 55:99-102.
- Cengiz FB, Duman D, Sirmaci A, Tokgöz-Yilmaz S, Erbek S, Oztürkmen-Akay H, *et al.* Recurrent and private MYO15A mutations are associated with deafness in the Turkish population. *Genet Test Mol Biomarkers* 2010; 14:543-550.
- Fattahi Z, Shearer AE, Babanejad M, Bazazzadegan N, Almadani SN, Nikzat N, *et al.* Screening for MYO15A gene mutations in autosomal recessive nonsyndromic, GJB2 negative Iranian deaf population. *Am J Med Genet Part A* 2012; 158:1857-1864.
- Woo H-M, Park HJ, Baek JI, Park MH, Kim UK, Sagong B, *et al.* Whole-exome sequencing identifies MYO15A mutations as a cause of autosomal recessive nonsyndromic hearing loss in Korean families. *BMC Med Genet* 2013; 14:72.
- Anderson DW, Probst FJ, Belyantseva IA, Fridell RA, Beyer L, Martin DM, *et al.* The motor and tail regions of myosin XV are critical for normal structure and function of auditory and vestibular hair cells. *Hum Mol Genet* 2000; 9:1729-1738.
- Friedman TB, Liang Y, Weber JL, Hinnant JT, Barber TD, Winata S, *et al.* A gene for congenital, recessive deafness DFNB3 maps to the pericentromeric region of chromosome 17. *Nat Genet* 1995; 9:86-91.
- Belyantseva IA, Boger ET, Friedman TB. Myosin XVa localizes to the tips of inner ear sensory cell stereocilia and is essential for staircase formation of the hair bundle. *Proc Natl Acad Sci* 2003; 100:13958-13963.
- Liang Y, Wang A, Belyantseva IA, Anderson DW, Probst FJ, Barber TD, *et al.* Characterization of the Human and Mouse Unconventional Myosin XV Genes Responsible for Hereditary Deafness DFNB3 and Shaker 2. *Genomics* 1999; 61: 243-258.
- Belguith H, Aifa-Hmani M, Dhoub H, Said MB, Mosrati MA, Lahmar I, *et al.* Screening of the DFNB3 locus: identification of three novel mutations of MYO15A associated with hearing loss and further suggestion for two distinctive genes on this locus. *Genet Test Mol Biomarkers* 2009; 13:147-151.
- Liburd N, Ghosh M, Riazuddin S, Naz S, Khan S, Ahmed Z, *et al.* Novel mutations of MYO15A associated with profound deafness in consanguineous families and moderately severe hearing loss in a patient with Smith-Magenis syndrome. *Hum Genet* 2001; 109:535-541.
- Wang A, Liang Y, Fridell RA, Probst FJ, Wilcox ER, Touchman JW, *et al.* Association of unconventional myosin MYO15 mutations with human nonsyndromic deafness DFNB3. *Science* 1998; 29:1447-1451.
- Grimberg J, Nawoschik S, Belluscio L, Mckee R, Turck A, Eisenberg A, *et al.* A simple and efficient non-organic procedure for the isolation of genomic DNA from blood. *Nucleic Acids Res* 1989; 17:8390-8390.
- Kleihues P, Schäuble B, zur Hausen A, Estève J, Ohgaki H. Tumors associated with p53 germline mutations: a synopsis of 91 families. *Am J Pathol* 1997; 150:1.
- Tabatabaiefar MA, Alasti F, Zohour MM, Shariati L, Farrokhi E, Farhud D, *et al.* Genetic linkage Analysis of 15 DFNB Loci in A group of Iranian families with Autosomal recessive hearing loss. *Iran J Public Health* 2011; 40:34-38.
- Lindner TH, Hoffmann K. easyLINKAGE: a PERL script for easy and automated two-/multi-point linkage analyses. *Bioinformatics* 2005; 21:405-407.
- Thiele H, Nürnberg P. HaploPainter: a tool for drawing pedigrees with complex haplotypes. *Bioinformatics* 2005; 21:1730-1732.

28. Berg JS, Powell BS, Cheney RE. *A millennial myosin census*. Mol Biol Cell 2001; 12:780-794.
29. Gao X, Zhu QY, Song YS, Wang GJ, Yuan YY, Xin F, et al. Novel compound heterozygous mutations in the MYO15A gene in autosomal recessive hearing loss identified by whole-exome sequencing. J Transl Med 2013; 11: 284-291.
30. Xia H, Huang X, Guo Y, Hu P, He G, Deng X, et al. Identification of a novel MYO15A mutation in a Chinese family with autosomal recessive nonsyndromic hearing loss. PLoS ONE 2015; 10: 1-9.
31. Weber KL, Sokac AM, Berg JS, Cheney RE, Bement WM. A microtubule-binding myosin required for nuclear anchoring and spindle assembly. Nature 2004; 431:325-329.
32. Mburu P, Mustapha M, Varela A, Weil D, El-Amraoui A, Holme RH, et al. Defects in whirlin, a PDZ domain molecule involved in stereocilia elongation, cause deafness in the whirler mouse and families with DFNB31. Nat Genet 2003; 34:421-428.
33. Shearer AE, Hildebrand MS, Webster JA, Kahrizi K, Meyer NC, Jalalvand K, et al. Mutations in the first MyTH4 domain of MYO15A are a common cause of DFNB3 hearing loss. Laryngoscope 2009; 119:727-733.
34. Reiisi S, Sanati MH, Tabatabaiefar MA, Ahmadian S, Reiisi S, Parchami S, et al. The study of SLC26A4 gene causing autosomal recessive hearing loss by linkage analysis in a Cohort of Iranian Populations. Int J Mol Cell Med 2014; 3:176.

Archive of SID