

## Anti-melanogenic activity of *Viola odorata* different extracts on B16F10 murine melanoma cells

Vafa Baradaran Rahimi<sup>1</sup>, Vahid Reza Askari<sup>1</sup>, Seyed Ahmad Emami<sup>2</sup>, Zahra Tayarani-Najaran<sup>3\*</sup>

<sup>1</sup> Student Research Committee, Department of Pharmacology, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

<sup>2</sup> Department of Pharmacognosy, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

<sup>3</sup> Department of Pharmacodynamics and Toxicology, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

### ARTICLE INFO

#### Article type:

Original article

#### Article history:

Received: Jun 9, 2016

Accepted: Jan 12, 2017

#### Keywords:

B16F10 cell line

Melanin

ROS

Tyrosinase

Violaceae

*Viola odorata* L.

### ABSTRACT

**Objective(s):** In previous studies, antioxidant activity of *Viola odorata* L. has been demonstrated. In this study, we have investigated the anti-melanogenic effect of extract and fractions of the plant in B16F10 cell line.

**Materials and Methods:** Impact of different increasing concentrations of extract and fractions of *V. odorata* was evaluated on cell viability, cellular tyrosinase, melanin content and mushroom tyrosinase as well as ROS production in B16F10 murine melanoma cell line.

**Results:** *Viola odorata* had no cytotoxicity on B16F10 cells compared to control group. Kojic acid as positive control had significant decreasing effects on cellular and mushroom tyrosinase activity, melanin content and ROS production ( $P < 0.001$ , for all cases). *V. odorata* (1-20  $\mu\text{g/ml}$ ) decreased all measured parameters including cellular tyrosinase and melanin content as well as ROS production and among all extract and fractions ethyl acetate fraction had the best effect ( $P < 0.05$ ).

**Conclusion:** *Viola odorata* had promising anti-melanogenic activity through inhibition of cellular tyrosinase activity and ROS production as well as melanin content. More basic and clinical studies need to aver its impact.

#### ► Please cite this article as:

Baradaran Rahimi V, Askari VR, Emami SA, Tayarani-Najaran Z. Anti-melanogenic activity of *Viola odorata* different extracts on B16F10 murine melanoma cells. Iran J Basic Med Sci 2017; 20:242-249; <http://dx.doi.org/10.22038/ijbms.2017.8350>

### Introduction

Skin is the vastest organ covering about 16% of total body weight of the adult (1). Furthermore, the skin is a multi-functional barrier protecting internal organs from physical and chemical injuries, pathogens, dehydration and ultraviolet (UV) irradiation as well as controlling body temperature, excretion of wastes, production of vitamin D and playing important role in psychosocial relations (2, 3). Skin tone is an individual characteristic that is most variable in range of pale to nigra in different racial origin and dependent to polymeric and non-protein compound named melanin, which found in basal epidermis layer (4).

Melanin is the main pigment in primates, synthesized in melanosomes in melanogenesis process and transferred to the other part of epidermis. Melanin cares against harmful impact of oxidative stress and DNA damage, and plays scavenging role in both internal and external sources of free radical (3, 5). Melanogenesis is started with tyrosinase (EC: 1.14.15.1), which oxidize the tyrosine to dopaquinone. In the presence of glutathione or cysteine (thiols moiety), dopaquinone in consecutive stages is converted to red pigment called pheomelanin,

while in the absence of glutathione or cysteine, black pigment named eumelanin is made (6). Tyrosinase is the key and rate-limiting enzyme in melanogenesis process and has a binuclear copper active site which catalyzes the hydroxylation of phenols to catechols and consecutively oxidized catechols to quinones (6, 7). Despite the protective role of melanin, excessive accumulations of melanin cause hyperpigmentation areas in the skin which named spot (8).

*Viola odorata* L. is a grassy and perennial plant (Figure 1) that belongs to Violaceae (9). In previous studies, many pharmacological properties of *V. odorata* has been demonstrated including: anti-bacterial (10), anti-inflammatory (11), anti-cancer (12, 13), anti-pyretic (14), hypnotic and sedative (9, 15), blood pressure lowering (14), diuretic and laxative (16), as well as anti-oxidant activity (17).

Pigmentation disorders is the third cause of skin consultation mainly in patients in range of 15-30 and 40-54 years old (18). So, in the present study we extracted the aerial parts of *V. odorata* with methanol. The extract then fractionated with ascending polarity solvents including *n*-hexane,

\*Corresponding author: Zahra Tayarani-Najaran. Department of Pharmacodynamics and Toxicology, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran. Tel: +98-51-38823255; email: tayaraninz@mums.ac.ir



**Figure 1.** The picture shows the leaves and flowers of *V. odorata*. (Ref: [http://americanvioletsociety.org/Violet\\_Keys/Viola\\_odorata\\_002\\_L.htm](http://americanvioletsociety.org/Violet_Keys/Viola_odorata_002_L.htm); photographed by Susan Polan)

dichloromethane ( $\text{CH}_2\text{Cl}_2$ ), ethyl acetate (EtOAc), *n*-butanol (BuOH), and finally water ( $\text{H}_2\text{O}$ ) to investigate the anti-melanogenic activity of the plant on B16F10 murine melanoma cell line.

## Materials and Methods

### Preparation of plant extracts

*Viola odorata* was collected from Mashhad, Razavi Khorasan Province, Northeast of Iran in the flowering stage in month of July 2013. The plant was identified by Mrs. Suzani and voucher sample has been deposited at the herbarium of School of Pharmacy, Mashhad University of Medical Sciences, Iran (No. 12855).


The leaves were separated, dried in shadow and powdered. Then, 327 g of leaves powder were macerated with 95% methanol for 24 hr at controlled room temperature (23-27 °C). The mixture was percolated as described previously (19). The extract was concentrated via a rotary evaporator and subsequently freeze dried. This yielded about 15.3% (50 g) extract.

Thereafter, the methanol extract was re-dissolved in 95% of methanol, transferred to a decantation funnel and then fractionated with following solvents: *n*-hexane,  $\text{CH}_2\text{Cl}_2$ , EtOAc, BuOH and finally  $\text{H}_2\text{O}$  as schemed in Figure 2 (20).

Dried extract/fractions were dissolved in final concentration of DMSO lesser than 1% (v/v) plus completed RPMI-1640 medium. To evaluate the DMSO at this concentration had no significant effect on experiments; the same concentration was examined in separated group in each tests.

### Cell culture

B16F10 murine melanocyte cell line was obtained from Pasture Institute, Tehran, Iran. B16F10 cells were cultured in completed RPMI-1640 medium (10% FBS, 100 U/ml penicillin, 100  $\mu\text{g}/\text{ml}$  streptomycin, 2 mM L-glutamine, and 1 mM sodium pyruvate and HEPES buffer) at 37 °C in 5%  $\text{CO}_2$ . Cells



**Figure 2.** Extraction and fractionation of *Viola odorata* aerial parts

were passaged every 3 days until achieved confluent monolayers melanocytes, and then harvested using 0.05% trypsin and 0.54 mM ethylenediaminetetraacetic acid (EDTA). Kojic acid (2 mM) and medium containing DMSO were used as negative and positive control, respectively in all experiments.

### Cell viability assay

Cell viability was examined using alamarBlue® dye. Upon entering this dye to viable cells, blue and non-fluorescent resazurin is converted to purple and detectable fluorescent dye (21).

About  $10^4$  B16F10 cells/100  $\mu\text{l}$  were cultured in flat-96 well plate overnight. Next, a range of 1 to 32  $\mu\text{g}/\text{ml}$  of *V. odorata* extract and fractions were added to B16F10 cells. After 48 hr, 15  $\mu\text{l}$  of alamarBlue® solution (0.1 mg/ml) was added to each well and incubated at 37 °C for 3 hr. The absorbance was read using micro-plate reader (awareness Inc., US) at 570 and 600 nm. The assay was performed in triplicate and repeated four times for each sample. The cell viability was expressed as percent of live cells.

### Analysis of mushroom tyrosinase function

The activity of mushroom tyrosinase as the enzyme which is responsible for converting levodopa (L-dopa) to dopamine was examined using spectrophotometric method as previously described (22, 23). Briefly, 160  $\mu\text{l}$  of L-Dopa at the concentration of 5 mM with extract and fractions of *V. odorata* (1-64  $\mu\text{g}/\text{ml}$ ) were added to flat-96 well plate and incubated with 20  $\mu\text{l}$  of mushroom tyrosinase (200 unit/ml) at 37 °C for 30 min. The absorbance was measured at 475 nm.

### Measurement of cellular tyrosinase activity

Cellular tyrosinase activity was performed using spectrophotometric method with measurement of

oxidation of Dopa to Dopachrome (22, 23). In brief, B16F10 melanoma cells were cultured at density of  $10^6$  cells/well in flat-6 well plate. Cells were mixed with or without different concentrations (1, 5, 10 and 20  $\mu\text{g/ml}$ ) of extract of *V. odorata* as well as kojic acid (2 mM) and incubated for 24 hr. The supernatant was removed and rinsed cells with phosphate buffer saline (PBS) twice. Next, cells were detached, transferred to 1.5 ml micro-tube and centrifuged at 1100 rpm for 7 min. 100  $\mu\text{l}$  of lysis buffer contains 100 mM  $\text{Na}_3\text{PO}_4$  (pH=6.8) was mixed with cells for 30 min on ice. After centrifugation at 10000 rpm at 4 °C for 20 min, 100  $\mu\text{l}$  of 5 mM L-dopa was incubated with 100  $\mu\text{l}$  of supernatants (each contain 100  $\mu\text{g}$  the protein) at 37 °C for 2 hr. Cellular tyrosinase activity was measured by measuring the absorbance of each wells compared to control group at 475 nm.

#### Assessment of melanin content

B16F10 cells were cultured at density of  $10^5$  cells per well of flat-six well plate overnight. *Viola odorata* extract and fractions (1, 5 and 10  $\mu\text{g/ml}$ ) were added to well, and incubated for 24 hr. Then, the supernatants were removed and wells rinsed with PBS twice. Cells were detached and centrifuged at 1100 rpm for 7 min. 100  $\mu\text{l}$  of 2 M NaOH was added to cells, incubated at 100 °C for 30 min and optical density measured using micro-plate reader (Awareness Inc., US) at 405 nm (22).

#### Evaluation of in vitro reactive oxygen species (ROS) level

Dichloro-dihydro-fluorescein diacetate (DCFH-DA) assay is a fluorimetric probe for detection of oxidative stress (24). Reactive oxygen species level was measured as described previously with minor modifications (25). Twenty thousands of B16F10 melanoma cells were cultured in a flat-96 well plate and incubated overnight. Different concentrations (1, 5 and 10  $\mu\text{g/ml}$ ) of *V. odorata* extract and fractions were incubated with cells for 24 hr. Next,  $\text{H}_2\text{O}_2$  at

concentration of 24 mM was added to each well and incubated at 37 °C for 30 min. Then, 2.5  $\mu\text{l}$  of DCFH-DA was added to the wells, and incubated for 30 min. The intensities were measured using ELISA reader (Awareness Inc., US) at excitation wavelength of 504 nm and emission wavelength of 524 nm.

#### Statistical analysis

The results were expressed as means $\pm$ SEM. Group-data in comparison to un-treated group were performed using one way analysis of variance (ANOVA) with Dunnett's *post hoc* test by GraphPad Prism @ 6 (GraphPad Software, San Diego, CA) software. Significance was considered at  $P<0.05$ .

## Results

#### The effect of extract and fractions of *V. odorata* on cell viability

All different concentrations of the extract and fractions of *V. odorata* led to a minimal decrease in cell viability which was not significant (Figure 3).

#### The effect of extract and fractions of *V. odorata* on cellular tyrosinase activity

All tested concentrations of both methanol extract and  $\text{CH}_2\text{Cl}_2$  fraction, and three higher concentrations of BuOH fraction of *V. odorata* significantly decreased the cellular tyrosinase activity ( $P<0.05$ , Figure 4). Furthermore, three lower concentrations of both  $\text{H}_2\text{O}$  and EtOAc fractions, and two lower concentrations of *n*-hexane fraction of *V. odorata* led to a significant decrement ( $P<0.05$  to 0.001, Figure 4).

#### The effect of extract and fractions of *V. odorata* on melanin content

Three concentrations of methanol extract, BuOH and EtOAc fractions and two lower concentrations of both  $\text{H}_2\text{O}$  and  $\text{CH}_2\text{Cl}_2$  fractions could significantly decrease the melanin content in cells in comparison to negative control group ( $P<0.01$ , Figure 5).





Figure 3. The effect of extract and fractions of *Viola odorata* on cell viability. Data were presented as mean $\pm$ SEM




**Figure 4.** The effect of extract and fractions of *Viola odorata* on cellular tyrosinase activity. Data are presented as mean±SEM. \**P*<0.05, \*\**P*<0.01 and \*\*\**P*<0.001 compared to negative control. ++*P*<0.01: compared to KA

**The effect of methanol extract and different fractions of *V. odorata* on mushroom tyrosinase activity**

All tested concentrations of extract and fractions had non-significant lowering effect on mushroom tyrosinase activity (Figure 6).


**The effect of methanol extract and different fractions of *V. odorata* on ROS production**

Concentrations of 1-10 µg/ml of methanol extract, and BuOH and EtOAc fractions significantly decreased ROS production, while the concentrations of 1 and 5 µg/ml of both H<sub>2</sub>O and CH<sub>2</sub>Cl<sub>2</sub> fractions of *V. odorata* had significant decreased ROS in comparison to control untreated group (*P*<0.01 to 0.001, Figure 7).




**Figure 5.** The effect of extract and fractions of *Viola odorata* on melanin content. Data were presented as mean±SEM. \*\**P*<0.01 and \*\*\**P*<0.001 compared to negative control





**Figure 6.** The effect of extract and fractions of *Viola odorata* on mushroom tyrosinase activity. Data were presented as mean±SEM. \* $P < 0.05$ , \*\* $P < 0.01$  and \*\*\* $P < 0.001$  compared to negative control



**Figure 7.** The effect of extract and fractions of *Viola odorata* on *in vitro* reactive oxygen species (ROS) production. Data were presented as mean±SEM. \* $P < 0.05$ , \*\* $P < 0.01$  and \*\*\* $P < 0.001$  compared to negative control

## Discussion

According to available information, this study is the first original research on antimelanogenic activity of *V. odorata*. In this study the effect of different extracts and fractions of *V. odorata* including methanol extract and *n*-hexane,  $\text{CH}_2\text{Cl}_2$ , BuOH, EtOAc and  $\text{H}_2\text{O}$  fractions on proliferation, cellular and mushroom tyrosinase activity, melanin content and

ROS production were examined using B16F10 murine melanoma cell line.

Methanol extract and fractions of *V. odorata* showed no significant effect on cell viability as mentioned in Figure 3. Although *V. odorata* at concentrations 1-32  $\mu\text{g/ml}$  had minimal effect on viability of cells but it was not significant. Based on our results, concentrations equal and less than 32

µg/ml were chosen for more evaluations.

Hyperpigmentation disorders are considered as inflammatory problems and have different causes such as excessive activity of melanocytes and increased melanin aggregation that leads to dark areas called spot. In addition, many factors can lead to spots including pregnancy, oral contraceptive pills, aging, excessive use of anti-acne drugs, cosmetics, perfumes and some chemical materials, etc. (6).

Melanin is composed and secreted by melanocytes during a regular and complex process in human skin (26). Sunlight exposure and especially UV radiation stimulates melanocytes to synthesize melanin to protect skin (7). Skin pigmentation is determined by different physiologic processes of melanocytes creation, expression of the melanosomes residuals, melanosomes transferring to keratinocytes and melanin distribution in the skin (27-29). Successful treatment usually obtained by combining two or more mechanisms through synergism effect (30). Furthermore, it has been demonstrated that tyrosinase has an important role in melanin formation and transferring to adjacent cells (6).

Kojic acid (5-hydroxy-2-hydroxy methyl-4-pyron) is a natural compound that is obtained from *Aspergillus niger*, *Acetobacter* spp. and *Penicillium* spp., and is used at concentrations of 1 to 4% as skin whitening agent (31). In this study, kojic acid, as positive control, was used at concentration of 2 mM and it significantly decreased cellular and mushroom tyrosinase activity, melanin content and ROS production. It has been shown that kojic acid inhibits tyrosinase through chelating  $\text{Cu}^{2+}$  in the active site of the enzyme (32), antioxidant and free radical scavenging activity (7). In contrast of good efficacy, limited application of kojic acid is due to toxicity and the side effect of the substance in inducing allergic dermatitis (33).

Herbal products are so popular among people due to naturalness and the belief to be non-toxic substitutes, but studies have been shown the toxicity of some plant. In melanocytes, decrease in the skin pigments production and the toxic impact limited the use of plant for hyperpigmentation problem. Hydroquinone is one of the compounds which irritates skin, induces toxicity for melanocytes and causes contact dermatitis and skin depigmentation (34). Also, it has carcinogenic effects which have been reported on rodents (35). Hydroquinone derivatives, ether monobenzyl hydroquinone and mono methylether hydroquinone, could also inhibit tyrosinase and have toxicity similar to hydroquinone (36, 37).

Rate limiting and first step of melanin synthesis is oxidation of tyrosine to dopa-quinone through acting tyrosinase (38). Tyrosinase catalyze and oxides the monophenols and diphenols to quinines (6). Considering the results, methanol extract of *V.*

*odorata* at concentrations 1-20 µg/ml could significantly decrease tyrosinase activity in a concentration dependent manner. Among all extract and fractions of *V. odorata*, EtOAc extract that contains medium polarity components such as flavonoids was more effective than other extracts.

In the next step, the melanin content was measured to confirm the cellular tyrosinase results. All extract and fractions except the *n*-hexane extract, could significantly decrease melanin content. However, EtOAc fraction had stronger effects than other extract and fractions which supports the results obtained in cellular tyrosinase test.

Herbal antioxidants such as glutathione, phytic acid and ubi-quinone (39, 40) could decrease the intracellular oxidative stress and play a protective role against disorders associated with oxidative stress (7). Herbal antioxidants could decrease the oxidation of tyrosin to dopaquinone and melanin synthesis in the melanogenesis pathway (41). Aging process will be accelerate in all body parts including skin as a result of the gradual accumulation of ROS which causes skin damages in the skin exposed to sunlight. Melanin has an important role in absorbing free radicals which is produce in cytoplasm of the cells irritated by UV irradiation (42). Antioxidants have important role in decreasing free radicals and the melanin content. Considering the results, extract and fractions of *V. odorata* could reduce ROS induced by hydrogen peroxide and the EtOAc extract was more effective than other extract and fractions. Furthermore, it has been shown that concentrations between 25-800 µg/ml of *V. odorata* have antioxidant activity in radical scavenging by DPPH,  $\text{Fe}^{2+}$  chelating, nitric oxide and hydrogen peroxide chelating tests (17).

In addition, according to investigations and literature survey we have done, there is no similar study on the genus *Viola* which emphasizes more investigations on other species.

## Conclusion

In summary, it can be suggested that different extracts and fractions of *V. odorata* could decrease total melanin content, cellular tyrosinase activity and oxygen free radicals in B16F10 cells and could have promising therapeutic effects on skin inflammatory diseases such as hyperpigmentation. In addition, our evaluation presented that the ethyl-acetate fraction had best anti-melogenic activity compared to other fractions or extract. However, we suggest that more basic and clinical studies will be done on ethyl-acetate fraction to establish this effect.

## Acknowledgment

This study has been financially supported by the Research Council of Mashhad University of Medical Sciences. This paper is a part of Pharm D thesis of the

first author which has been registered as No. 922760 in Research Council of Mashhad University of Medical Sciences, Mashhad, Iran.

### Conflict of interest

Authors declare that there is no conflict of interest.

### References

- Murakami S and Miki Y. Human skin histology using high-resolution echography. *J Clin Ultrasound* 1989; 2: 77-82.
- Tran K T, Griffith L, Wells A. Extracellular matrix signaling through growth factor receptors during wound healing. *Wound Repair Regen* 2004; 3: 262-268.
- Wang Z, Qian Y, Li L, Pan L, Njunge LW, Dong L, et al. Evaluation of emulsion electrospun polycaprolactone/hyaluronan/epidermal growth factor nanofibrous scaffolds for wound healing. *J Biomater Appl* 2016; 6: 686-698.
- Tadokoro T, Kobayashi N, Zmudzka BZ, Ito S, Wakamatsu K, Yamaguchi Y, et al. UV-induced DNA damage and melanin content in human skin differing in racial/ethnic origin. *Faseb j* 2003; 9: 1177-1179.
- Ando H, Kondoh H, Ichihashi M, Hearing VJ. Approaches to identify inhibitors of melanin biosynthesis via the quality control of tyrosinase. *J Invest Dermatol* 2007; 4: 751-761.
- Chang TS. An updated review of tyrosinase inhibitors. *Int J Mol Sci* 2009; 6: 2440-75.
- Baek S H, Nam I J, Kwak H S, Kim K C, and Lee S H. Cellular anti-melanogenic effects of a *Euryale ferox* seed extract ethyl acetate fraction via the lysosomal degradation machinery. *Int J Mol Sci* 2015; 5: 9217-35.
- Wu ZS, Zhang M, Wang SJ. Effects of high-pressure argon and nitrogen treatments on respiration, browning and antioxidant potential of minimally processed pineapples during shelf life. *J Sci Food Agric* 2012; 11: 2250-9.
- Feyzabadi Z, Jafari F, Kamali SH, Ashayeri H, Badiie Aval S, Esfahani MM, Efficacy of *Viola odorata* in treatment of chronic insomnia. *Iran Red Crescent Med J* 2014; 12: e17511.
- Zarrabi M, Dalirfardouei R, Sepehrizade Z, and Kermanshahi R K. Comparison of the antimicrobial effects of semipurified cyclotides from Iranian *Viola odorata* against some of plant and human pathogenic bacteria. *J Appl Microbiol* 2013; 2: 367-75.
- Koochek M, Pipelzadeh M, and Mardani H. The effectiveness of *Viola odorata* in the prevention and treatment of formalin-induced lung damage in the rat. *J Herbs Spices Med Plants* 2003; 2: 95-103.
- Lindholm P, Goransson U, Johansson S, Claeson P, Gullbo J, Larsson R, et al. Cyclotides: a novel type of cytotoxic agents. *Mol Cancer Ther* 2002; 6: 365-9.
- Gerlach SL, Rathinakumar R, Chakravarty G, Goransson U, Wimley WC, Darwin S, et al. Anticancer and chemosensitizing abilities of cycloviolacin O2 from *Viola odorata* and psyle cyclotides from *Psychotria leptothyrsa*. *Biopolymers* 2010; 5: 617-25.
- Siddiqi HS, Mehmood MH, Rehman NU, Gilani AH. Studies on the antihypertensive and antidyslipidemic activities of *Viola odorata* leaves extract. *Lipids Health Dis* 2012; 6.
- Monadi A, Rezaie A. Evaluation of sedative and pre-anesthetic effects of *Viola odorata* Linn. extract compared with diazepam in rats. *Bull Environ Pharmacol Life Sci* 2013; 7: 125-131.
- Ganapaty S, Dash G K, Subburaju T, Suresh P. Diuretic, laxative and toxicity studies of *Cocculus hirsutus* aerial parts. *Fitoterapia* 2002; 1: 28-31.
- Ebrahimzadeh MA, Nabavi SM, Nabavi SF, Bahramian F, Bekhradnia AR. Antioxidant and free radical scavenging activity of *H. officinalis* L. var. *angustifolius*, *V. odorata*, *B. hyrcana* and *C. speciosum*. *Pak J Pharm Sci* 2010; 1: 29-34.
- Cestari TF, Dantas LP, and Boza JC. Acquired hyperpigmentations. *An Bras Dermatol* 2014; 1: 11-25.
- Sadraei H, Asghari G, and Emami S. Effect of *Rosa damascena* Mill. flower extract on rat ileum. *Res Pharm Sci* 2013; 4: 277-284.
- Seidel V. Initial and bulk extraction. In: Sarker SD, Latif Z, Gray AI, editors. *Natural Product Isolation*. 2nd ed. Humana Press 2006; 27-46.
- Shafaei-Bajestani N, Emami S A, Asili J, Tayarani-Najaran Z. Anti-apoptotic effect of taxodione on serum/glucose deprivation-induced PC12 cells death. *Cell Mol Neurobiol* 2014; 8: 1103-9.
- Qiao Z, Koizumi Y, Zhang M, Natsui M, Flores M J, Gao L, et al. Anti-melanogenesis effect of *Glechoma hederacea* L. extract on B16 murine melanoma cells. *Biosci Biotechnol Biochem* 2012; 10: 1877-83.
- Prakash L and Majeed M. Multifunctional skin tone lighteners from nature: An overview. *Eurocosmetics* 2009; 19-23.
- Aranda A, Sequedo L, Tolosa L, Quintas G, Burello E, Castell JV, and Gombau L. Dichloro-dihydro-fluorescein diacetate (DCFH-DA) assay: a quantitative method for oxidative stress assessment of nanoparticle-treated cells. *Toxicol in vitro* 2013; 2: 954-63.
- Huang HC, Hsieh WY, Niu YL, Chang TM. Inhibition of melanogenesis and antioxidant properties of *Magnolia grandiflora* L. flower extract. *BMC Complement Altern Med* 2012; article number 72.
- Hearing VJ. Determination of melanin synthetic pathways. *J Invest Dermatol* 2011; E1: E8-e11.
- Eller MS, Gilchrist BA. Tanning as part of the eukaryotic SOS response. *Pigment Cell Res* 2000; 94-7.
- Gilchrist BA, Park HY, Eller MS, Yaar M. Mechanisms of ultraviolet light-induced pigmentation. *Photochem Photobiol* 1996; 1: 1-10.
- Coelho SG, Zhou Y, Bushar HF, Miller SA, Zmudzka BZ, Hearing VJ, et al. Long-lasting pigmentation of human skin, a new look at an overlooked response to UV. *Pigment Cell Melanoma Res* 2009; 2: 238-41.
- Solano F, Briganti S, Picardo M, Ghanem G. Hypopigmenting agents: an updated review on biological, chemical and clinical aspects. *Pigment Cell Res* 2006; 6: 550-71.
- Kahn V. Effect of kojic acid on the oxidation of DL-DOPA, norepinephrine, and dopamine by mushroom tyrosinase. *Pigment Cell Res* 1995; 5: 234-40.
- Cabanes J, Chazarra S, Garcia-Carmona F. Kojic acid, a cosmetic skin whitening agent, is a slow-binding inhibitor of catecholase activity of tyrosinase. *J Pharm Pharmacol* 1994; 12: 982-5.
- Nakagawa M, Kawai K, Kawai K. Contact allergy to kojic acid in skin care products. *Contact Dermatitis* 1995; 1: 9-13.

34. Lawrence N, Bligard CA, Reed R, Perret WJ. Exogenous ochronosis in the United States. *J Am Acad Dermatol* 1988; 2: 1207-11.
35. Nordlund JJ, Grimes PE, and Ortonne JP. The safety of hydroquinone. *J Eur Acad Dermatol Venereol* 2006; 7: 781-7.
36. Kamakshi R. Fairness via formulations: a review of cosmetic skin-lightening ingredients. *J Cosmet Sci* 2012; 1: 43-54.
37. Naish-Byfield S, Cooksey C, Latter A, Johnson C, Riley P. *In vitro* assessment of the structure-activity relationship of tyrosinase-dependent cytotoxicity of a series of substituted phenols. *Melan res* 1991; 4: 273-288.
38. Halaban R, Patton RS, Cheng E, Svedine S, Trombetta ES, Wahl ML, et al. Abnormal acidification of melanoma cells induces tyrosinase retention in the early secretory pathway. *J Biol Chem* 2002; 17: 14821-8.
39. Pompella A, Visvikis A, Paolicchi A, De Tata V, Casini AF. The changing faces of glutathione, a cellular protagonist. *Biochem Pharmacol* 2003; 8: 1499-1503.
40. Draelos ZD. Novel topical therapies in cosmetic dermatology. *Curr Prob Dermatol* 2000; 5: 235-239.
41. Katsambas A D and Stratigos A J. Depigmenting and bleaching agents: coping with hyperpigmentation. *Clin Dermatol* 2001; 4: 483-488.
42. Mason HS. The chemistry of melanin; mechanism of the oxidation of dihydroxyphenylalanine by tyrosinase. *J Biol Chem* 1948; 1: 83-99.

Archive of SID