

Protective effects of phenolic acids on mercury-induced DNA damage in precision-cut kidney slices

Irma Edith Carranza-Torres^{1,2}, Ezequiel Viveros-Valdez³, Nancy Elena Guzmán-Delgado⁴, Sara García-Davis³, Javier Morán-Martínez¹, Nadia Denys Betancourt-Martínez¹, Isaías Balderas-Rentería⁵, Pilar Carranza-Rosales^{2*}

¹ Departamento de Biología Celular y Ultraestructura, Centro de Investigación Biomédica, Facultad de Medicina, Universidad Autónoma de Coahuila. Torreón, Coah. México

² Departamento de Biología Celular y Molecular, Centro de Investigación Biomédica del Noreste, Instituto Mexicano del Seguro Social, Monterrey, NL. México

³ Departamento de Química Analítica, Facultad de Ciencias Biológicas, Universidad Autónoma de Nuevo León, San Nicolás de los Garza, NL. México

⁴ División de Investigación, Unidad Médica de Alta Especialidad # 34, Instituto Mexicano del Seguro Social, Monterrey, NL. México

⁵ Laboratorio de Ingeniería Genética y Genómica, Facultad de Ciencias Químicas, Universidad Autónoma de Nuevo León, San Nicolás de los Garza, NL. México

ARTICLE INFO

Article type:

Original article

Article history:

Received: Feb 22, 2018

Accepted: Nov 12, 2018

Keywords:

Comet assay

Genotoxicity

Mercuric chloride

Phenolic compounds

Precision-cut tissue slices

ABSTRACT

Objective(s): Precision-cut tissue slices are considered an organotypic 3D model widely used in biomedical research. The comet assay is an important screening test for early genotoxicity risk assessment that is mainly applied on in vitro models. The aim of the present study was to provide a 3D organ system for determination of genotoxicity using a modified method of the comet assay since the stromal components from the original tissue make this technique complicated.

Materials and Methods: A modified comet assay technique was validated using precision-cut hamster kidney slices to analyze the antigenotoxic effect of the phenolic compounds caffeic acid, chlorogenic acid, and rosmarinic acid in tissue slices incubated with 15 μM HgCl_2 . Cytotoxicity of the phenolic compounds was studied in Vero cells, and by morphologic analysis in tissue slices co-incubated with HgCl_2 and phenolic compounds.

Results: A modification of the comet assay allows obtaining better and clear comet profiles for analysis. Non-cytotoxic concentrations of phenolic acids protected kidney tissue slices against mercury-induced DNA damage, and at the same time, were not nephrotoxic. The highest protection was provided by 3 $\mu\text{g}/\text{ml}$ caffeic acid, although 6 $\mu\text{g}/\text{ml}$ rosmarinic and 9 $\mu\text{g}/\text{ml}$ chlorogenic acids also exhibited protective effects.

Conclusion: This is the first time that a modification of the comet assay technique is reported as a tool to visualize the comets from kidney tissue slices in a clear and simple way. The phenolic compounds tested in this study provided protection against mercury-induced genotoxic damage in precision-cut kidney slices.

► Please cite this article as:

Carranza-Torres IE, Viveros-Valdez E, Guzmán-Delgado NE, García-Davis S, Morán-Martínez J, Betancourt-Martínez ND, Balderas-Rentería I, Carranza-Rosales P. Protective effects of phenolic acids on mercury-induced DNA damage in precision-cut kidney slices. Iran J Basic Med Sci 2019; 22:367-375. doi: 10.22038/ijbms.2019.30056.7242

Introduction

Precision-cut tissue slices represent a three-dimensional (3D) biological model closely resembling the organ from which they are prepared, where all cell types and tissue architecture are preserved, as well as metabolic activity and transport mechanisms (1-3). These characteristics are very treasured in the investigation of different biological activities. Several studies show that precision-cut kidney slices (PCKS) are a useful model for studying drug metabolism, nephrotoxicity, cryopreservation, and fibrosis (4-8) among other applications.

The comet assay is a quantitative technique through which visual evidence of DNA damage in individual eukaryotic cells can be measured. This method possesses a number of advantages when compared to other genotoxicity tests; one of them being that it's readily modifiable for adaptation to a variety of experimental requirements. Over time, many improvements and

variations of the method have been developed for a number of experimental purposes (9, 10); however, new applications for this method require standardization. A small number of DNA damage studies using the comet assay in cells from precision-cut tissue slices have been reported as tools for genotoxic compounds testing (1, 11-14). In these studies, the comet assay was used to study genotoxicity induced by several xenobiotics or nanomaterials in liver and lung slices (1, 12-14); and, to evaluate the antigenotoxic effects of flavonoids (11). Depending on the kind of tissue used, different modifications were made. For example, human or murine tissue slices were prepared for the assay by mincing the tissue (14), enzymatic digestion was applied to murine lung slices (12), and human and rat liver slices were directly gently squashed with a coverslip (1, 11, 13). These reports demonstrated the usefulness of 3D tissue culture models for genotoxicity studies given that the main advantage of the said models is that the results obtained are comparable to *in vivo* conditions.

*Corresponding author: Pilar Carranza-Rosales. 5Departamento de Biología Celular y Molecular, Centro de Investigación Biomédica del Noreste, Instituto Mexicano del Seguro Social, Monterrey, NL. México. Tel/Fax: +52-818190-4035; Email: pilarcarranza@cibinmty.net

So far, PCKS studies aimed at determining the degree of protection offered by bioactive compounds against genotoxic damage have not been published.

On the other hand, large population groups of people are currently exposed to low levels of xenobiotics such as mercury, which has been implicated in the rising prevalence of autism spectrum disorders (15, 16). Occupational or environmental exposure to mercury causes many other health complications (17). Mercuric chloride (HgCl_2) toxicity is related to oxidative stress (18, 19), it is well known that it is genotoxic at low doses (20), and may be a primary cause of precancerous injuries (21). For these reasons, there is an increasing interest in naturally occurring antioxidants capable of acting as chelating agents and free-radical scavengers against heavy metal poisoning (22). In relation to this, phenolic acids (PAs) are free-radical scavengers universally distributed in the kingdom Plantae; they are essential for plant growth, reproduction, and serve as a defense mechanism against pathogens (23). The antioxidant activity of phenolic compounds and their possible uses in processed foods and treatments against many different diseases have been reported (24, 25). In previous studies, we reported on the anti-radical and antiproliferative activities against cancer cell lines of the phenolic compounds caffeic acid (CA), rosmarinic acid (RA), and chlorogenic acid (CGA) (26, 27). These PAs are present in a variety of species used to make teas, as well as other edible, and medicinal plants, and exhibit diverse biological activities (28-30). In recent years, they have received particular attention due to their role in the prevention of several human diseases.

In this study, we made an adaptation to the comet assay technique in order to better analyze the protective effects of CA, RA, and CGA on the genotoxic damage induced by HgCl_2 using the PCKS model. Our results show that the modification made to the comet assay enabled us to analyze comets in tissue slices in an easy and clear manner and that PAs protected kidney tissue slices against mercury-induced DNA damage, and that they were not nephrotoxic.

Materials and Methods

Chemicals

Mercuric chloride (HgCl_2), phenolic compounds (CA, RA, and CGA), normal, and low-melting-point agaroses, 1, 1-diphenyl-2-picrylhydrazyl (DPPH), 4-[3-(4-iodophenyl)-2-(4-nitrophenyl)-2H-5-tetrazolium]-1,3-benzene disulfonate (WST-1), and other chemicals were purchased from Sigma-Aldrich Corp. (St. Louis, MO, USA). Cell culture medium and fetal bovine serum (FBS) were obtained from Gibco Invitrogen (Carlsbad, CA, USA).

1, 1-diphenyl-2-picrylhydrazyl (DPPH) radical-scavenging activity

Before experiments with tissue slices and comet assay, free radical scavenging activity of the PAs was confirmed. To do this, serial dilutions of PAs dissolved in methanol were mixed with DPPH solution (2 mg/l) in 96-well microplates and incubated for 30 min at room temperature. The change in absorbance at 517 nm was measured. Mean values were obtained from

triplicate experiments. The percentage of inhibition was calculated using the following equation:

% inhibition = $[(A0 - A1)/A0] \times 100$, where $A0$ is the absorbance of the control and $A1$ is the absorbance of the samples (26).

Radical-scavenging activities were expressed as 90% inhibitory concentrations (IC_{90}), and a smaller value indicated more effective radical scavenging. IC_{90} values were calculated from log-dose inhibition curves obtained using a nonlinear regression algorithm (Prism 4.0; GraphPad Software, Inc., La Jolla, CA, USA).

Cell line and culture

Vero cells (kidney epithelial cells; American Type Culture Collection: CCL 81) were used to determine the nontoxic doses of the phenolic compounds to be used in PCKS. Cells were grown in M199 medium supplemented with 5% (v/v) FBS, penicillin G (100 units/ml), and streptomycin (100 $\mu\text{g}/\text{mL}$). Cultures were maintained at 37 °C in a humidified 5% CO_2 atmosphere.

Cytotoxicity assay

Vero cells were plated in 96-well flat-bottomed plates at 5000 cells/well and allowed to adhere overnight. Subsequently, cells were treated with different concentrations of the compounds (0–50 $\mu\text{g}/\text{ml}$). After 24 hr, the WST-1 colorimetric assay was used to monitor cell growth (27). Cell growth percentage was used to obtain dose-response curves and determine the 10% inhibitory concentration (IC_{10}). Values were expressed as the percentage variation relative to the control. Cellular inhibition was measured as a function of inhibitory concentration, and a higher IC_{10} value indicated lower cytotoxicity.

Preparation of precision-cut kidney slices

PCKS were prepared from two month old male Syrian golden hamsters (*Mesocricetus auratus*). Hamsters were killed with an overdose of sodium pentobarbital (6 mg/100 g) and treated following institutional and international guidelines for the humanitarian care of animals used in experimental work. Once the hamsters were unconscious, the kidneys were quickly removed and placed into ice-cold Krebs–Henseleit buffer (KB buffer). The external membranes covering the kidneys were separated with a scalpel, and the organs were cored to yield cylinders of 5 mm diameter. The cores were then sliced in oxygenated KB buffer (4 °C, 95:5 $\text{O}_2:\text{CO}_2$) into ~150- μm -thick slices, using a Brendel–Vitron tissue slicer (Vitron Inc., Tucson, AZ, USA). After preparation, slices were kept in ice-cold KB solution until incubation. The whole procedure from organ excision to incubation typically took 2 hr. The maintenance and care of the experimental animals complied with Instituto Mexicano del Seguro Social and Ley General de Salud (México) regulations and international guidelines for the humane use of laboratory animals.

Incubation of PCKS

PCKS were cultured in 24-well culture plates (Corning Life Sciences, Acton, MA, USA) in Dulbecco's modified Eagle's medium/F12 serum-free medium (1.3 ml/well) supplemented with glucose (25 mM) and gentamicin

(50 µg/ml) at 37 °C in a 95% air/5% CO₂ atmosphere. The slices were incubated for 30 min in culture medium (untreated control) with individual PAs (9 µg/mL CGA, 3 µg/ml CA, 6 µg/ml RA), or 15 µM HgCl₂, and co-incubated with HgCl₂ plus each PA (CGA+HgCl₂, CA+HgCl₂, and RA+HgCl₂) at the same concentration mentioned above. During incubation, plates were stirred at 30 rpm to facilitate gas exchange and substrate diffusion in the slices and ensure continuous floating of the slices in the medium. All experiments were performed with slices from at least three different experiments, and duplicate slices were used for each experimental condition.

Preparation of slides for the comet assay

We made a modification to the comet technique in order to obtain separated nuclei from PCKS- cells as well as cells maintaining their structural integrity. To do this a mechanical homogenization was performed according to the method described by Hymer and Kuff (31), however, a filtration step was not applied to the homogenates and the use of detergents was avoided. The method used in the present work is described below. Fully frosted microscope slides were pre-coated with a layer of 0.75% normal agarose. After incubation with the compounds, the PCKS were homogenized using a Potter-Elvehjem tissue grinder (Fisher Scientific, Hampton, New Hampshire, USA) in saccharose buffer (0.25 M saccharose and 1 mM MgCl₂) (31). Once partially disintegrated, the material was centrifuged at 5000 rpm for 10 min, and pellets containing nuclei and cells were resuspended in 90 µL of low-melting-point agarose (0.75%) and placed on the normal agarose layer. Next, coverslips were placed on top of the samples, and each cellular pellet was squashed to release cells. The liberated nuclei and cells spread over the microscope slide but remained embedded in the agarose. Slides were kept at 4 °C for up to 5 min to allow the agarose to solidify, after which the coverslips were removed. Finally, each preparation was covered with a layer of low-melting-point agarose.

Comet assay

The subsequent steps of the comet assay were performed as described by Singh *et al.* (32). Previously prepared slides were incubated in cold lysis solution [2.5 M NaOH, 0.1 M ethylenediaminetetraacetic acid (EDTA), and 0.01 M Tris with 1% Triton X-100, pH 10] at 4 °C for at least 1 hr. Then, slides were placed in a horizontal electrophoresis box, and the DNA was allowed to unwind for 20 min in freshly prepared alkaline electrophoresis buffer (300 mM NaOH and 1 mM EDTA, pH 13). Electrophoresis was conducted at 4 °C for 20 min at 25 V and 300 mA. Slides were then neutralized with Tris buffer (0.4 M, pH 7.5) and stained with ethidium bromide. Slides were observed under a fluorescence microscope with a charge-coupled device camera and analyzed using an image analysis system (CASPer). For each treatment, 100 single cells were analyzed; three separate experiments were conducted for each series. DNA damage was expressed as the olive tail moment, which is defined as the product of the tail length and fraction of total DNA in the tail. The tail moment incorporates a measure of both the smallest

detectable size of migrating DNA (reflected in the comet tail length) and the number of relaxed/broken pieces (represented by the intensity of DNA in the tail). The following equation was used:

$$\text{Olive tail moment} = (\text{tail.mean} - \text{head.mean}) \times \text{tail\%DNA}/100.$$

It is generally accepted that higher olive moment values indicate more damage to DNA.

Morphologic analysis

To study the effects of PAs on renal tissue, PCKS were incubated for 12 hr in culture medium with PAs alone (9 µg/ml CGA, 3 µg/ml CA, and 6 µg/ml RA) or 15 µM HgCl₂, and co-incubated with mercury and PAs (9 µg/mL CGA+15 µM HgCl₂, 3 µg/mL CA+15 µM HgCl₂, and 6 µg/mL RA+15 µM HgCl₂). After incubation, slices were washed with phosphate-buffered saline solution at 37 °C and fixed with 10% neutral formalin for 24 hr at room temperature. Slices were embedded in paraffin using an automated tissue processor. Subsequently, 5-µm thick sections were prepared from paraffin blocks containing the samples using a microtome. These were then stained with hematoxylin and eosin and studied under a conventional light microscope. Histopathologic criteria were used to establish the morphologic characteristics of normal tissue and were compared with those of slices subjected to different treatments.

Statistical analysis

Statistical significance was determined using one-way analysis of variance followed by the Tukey's test, comparing all groups with the control group. The results were considered statistically significant at $P < 0.01$. Data were expressed as means \pm standard deviation.

Results




Antioxidant and cytotoxic activities

Caffeic, rosmarinic, and chlorogenic acids are phenolic compounds that can be found in some herbs and spices and were evaluated for their anti-radical activity by the DPPH method; the highest anti-radical activity was exhibited by CA (IC₅₀ = 3.12 µg/ml), followed by RA and CGA (IC₅₀ = 5.61 and 8.52 µg/ml, respectively). Regarding cytotoxicity in Vero cells (normal kidney cell line), the nontoxic doses (IC₁₀) of CA, RA, and CGA were 3, 6.16, and 7.53 µg/ml, respectively. CA exhibited the highest cytotoxic and anti-radical effects, revealing a direct relationship between anti-radical and cytotoxic properties (Table 1).

Comet assay in PCKS

When prepared the tissue slices were for the comet assay, the resulting images showed large masses of stromal tissue and few isolated cells (Figure 1A); due to this, we decided to establish a simpler and faster method to measure the DNA damage in PCKS-cells. In order to accomplish this goal, we mechanically disintegrated the tissue slices by gently homogenizing them in a tissue grinder and using saccharose buffer to prevent the released cells from lysing. Both single nuclei and cells were obtained by this method, we were able to visualize that the said nuclei retained their integrity and would be useful for the purpose of our studies (Figure 1B).

Table 1. Radical scavenging and cytotoxic activities of phenolic acids

Phenolic Acid	Chemical Structure	Radical Scavenger (EC ₉₀)	Cytotoxicity (IC ₁₀)
Chlorogenic acid (CGA)		8.52 ± 1.5	7.53 ± 1.7
Rosmarinic acid (RA)		5.61 ± 0.9	6.16 ± 1.2
Caffeic acid (CA)		3.12 ± 0.7	2.78 ± 0.4

Data show the amount of PAs needed to capture 90 percent of free radicals and inhibit only 10 percent of normal kidney cells (non-cytotoxic doses). CA possesses the best radical-scavenging activity but also is the most cytotoxic. Values ($\mu\text{g/ml}$) represent the mean and standard deviation of triplicate measurements

Protective effect of phenolic acids on HgCl₂-induced DNA damage

In this study, we used PCKS and the comet assay to analyze the protective effect of PAs against the genotoxic damage induced by sub-cytotoxic doses of HgCl₂. PAs efficiently decreased HgCl₂-induced damage; and, no statistically significant differences were found relative to the untreated control (Figure 2).

With regard to the morphology of the comets formed

under different incubation conditions, CA induced shapes with less DNA damage compared with untreated control slices, PCKS treated with 15 μM HgCl₂, and slices co-incubated with 3 $\mu\text{g/ml}$ CA+15 μM HgCl₂ (Figure 3). These results suggest that CA has a greater protective effect against mercury-induced damage in renal tissue.

Morphological analysis

To establish whether the phenolic compounds had toxic effects on renal tissue, PCKS were incubated for 12




Figure 1. Fluorescent microscope images of tissue slices for the comet assay. PCKS were incubated with 15 μM HgCl₂ and processed for comet assay with the conventional technique (a), where a mass of tissue contains many renal cells trapped on the stroma (arrows) which are not appropriated for visual scoring after the squashing method. (b) Cells released from a treated slice after completion of the procedure using the homogenization method with the saccharose buffer. Ethidium bromide stain. Total magnification A (100X), B (400X)




Figure 2. Olive tail moment of precision-cut kidney slices exposed to mercuric chloride and treated with phenolic acids. The olive tail moment was determined from each of the treatments. HgCl₂ (15 μM) was used as control of damage; negative control was DMEM/F12 medium. Kidney slices were co-incubated separately with 15 μM HgCl₂ and CA (3 μg/ml), RA (6 μg/ml) and CGA (9 μg/ml). Values represent the mean and standard deviation of 100 cells/ treatment. Normality of the data was evaluated with the Shapiro-Wilks test. The T Student's Test was applied, and significant differences were analyzed with the Tukey's test ($P \leq 0.01$). A significant difference exists between slices treated with HgCl₂ (a) and those co-incubated with the PAs (b, c, d). CA exhibited the biggest difference (c)

hr in the presence of the treatments as described in the Materials and Methods section. The results show that compared to the control slices, the phenolic compounds did not induce alterations in the histological structure of the kidney (Figure 4D). In contrast, HgCl₂ caused interstitial edema, proximal tubular necrosis, and apoptosis (Figure 4E). These results indicate that the three studied phenolic compounds are not nephrotoxic. Furthermore, in PCKS co-incubated with the phenolic compounds+HgCl₂ (Figure 4F-H), morphological damage was less evident, although slight edema was observed in the interstices of the kidney. CGA (9 μg/ml) and RA (6 μg/ml) induced hydropic regeneration, which is associated with an adaptive response. In the same way as results obtained in the comet assay, CA induced minor morphologic changes and has the most effective protection at the tissue level.

Discussion

The aim of the present study was to render a simple and reproducible 3D *ex vivo* PCKS model for the

determination of mercury-induced genotoxic damage using the comet assay, as well as to analyze the protective effect of PAs in the same model.

Comet assays directly applied to a complex three-dimensional system without preparation of single cells represent a challenge mainly due to the fact that cells are lying on top of one another, and to stromal components present in the original tissue. In order to adequately visualize the comets obtained from PCKS incubated with the different treatments, it was necessary to disaggregate cells from each tissue slice cells under isotonic conditions in a glass homogenizer in the presence of saccharose buffer to avoid cell lysis (31), this process resulted in purified intact nuclei and cells from renal tissue. Thus, for the first time, we report a modification of the original technique that allowed us to visualize the comets from PCKS in a clear, simple, and reproducible manner for the comet assay. This variation made by our laboratory permits avoiding problems with overlapping comets or false positives made by squashing a great number of cells in tissue slices between the




Figure 3. Comet assay from kidney slices incubated with mercuric chloride and phenolic acids. Representative cell images from PCKS incubated with 15 μM HgCl₂ showed the typical form of long-tail comets indicating DNA strand breaks in the cells (a). Untreated slices (b) showed round nuclei, with intact DNA, while in the kidney slices incubated with 3 μg/mL CA+15 μM HgCl₂ the damage was attenuated (c). Ethidium bromide stain. Total magnification (200X)




Figure 4. Histopathologic analysis of precision-cut kidney slices treated with mercuric chloride and phenolic acids. PCKS were incubated in the presence of (a) cell culture medium, (b) 9 µg/ml CGA, (c) 3 µg/ml CA, (d) 6 µg/ml RA, (e) 15 µM HgCl₂, (f) CGA+HgCl₂, (g) CA+HgCl₂, and (h) RA+HgCl₂. Untreated slices showed morphologic characteristics of normal tissue, as well as preserved tubular structure (a). Slices incubated in the presence of phenolic acids alone show no signs of damage or morphological alterations (b, c, d). In contrast, mercury induced interstitial edema, tubular necrosis and the presence of apoptotic nuclei (e), dotted lines. In slices co-incubated with HgCl₂ and compounds, the damage induced by mercury was ameliorated; this was evident because of the conserved histologic structure, showing only mild interstitial edema. With CGA and RA hydropic degeneration was observed, whereas, with CA, less degenerative changes were observed (f, g, h). H&E stain. Total magnification 400X

coverslip and glass slide.

On the other hand, since its development by Ostling and Johanson (33), the comet assay is widely used to test the genotoxicity of chemical compounds both *in vitro* and *in vivo*. One advantage of the technique is that cells from various tissues of a wide variety of organisms can be studied (10). However, questions have arisen about the results obtained from cell cultures; in many cases, cells in culture cannot adequately metabolize some toxins because they lack important components of metabolic pathways such as cytochrome P450 (34, 35). This problem is addressed by alternative 3D systems, in which all cell types reside in their original tissue-matrix configuration and metabolic activity is preserved. Consequently, research laboratories have begun to use precision-cut slices to study the genotoxic effects of xenobiotics (12-14) and the protective effects of natural products (11), among other applications. This experimental approach has important advantages over *in vitro* and *in vivo* models. This model closely resembles the *in vivo* tissue function, which is highly desirable for the evaluation of new or existing compounds. In addition, precision-cut tissue slices offer the advantage of greater reproducibility in the experiments carried out with the same organ, it also allows to work simultaneously with different organs from the same animal, being able to obtain a large number of slices that can be cultivated, decreasing the number of experimental animals (36, 37). Few reports describe the use of the comet assay to measure DNA damage in *ex vivo* models such as murine and human liver and lung slices; however, reports show that this assay is a useful tool to study genotoxicity induced by xenobiotics such as tert-Butyl hydroperoxide, benzo (α) pyrene, formalin, 2-Nitrofluorene, and nanomaterials (1, 12-14); and to evaluate antigenotoxic effects of flavonoids (11).

In relation to the protective effect on the radical scavenging activity, the three PAs share the presence of the catechol group (3', 4'-dihydroxy group) in their structure (Table 1), and it is well known that at low concentrations, some compounds that have catechol groups provide protection to primary cultures of mice cortical neurons against methylmercury induced damage (38). Our results suggest a positive relationship between the radical scavenging effects and cytotoxicity (Table 1). In fact, it has been reported that the antioxidant or pro-oxidant activity of the phenolics depends on their concentration (39). Also, the cytotoxicity of some polyhydroxy benzenes studied by QSAR models indicates parallelism between polyphenol cytotoxicity and the rates of their single-electron oxidation, and points to the leading role of formation of the reactive oxygen their cytotoxicity (40). The catechol group has also the ability to donate electrons, which is related to its antioxidant capacity *in vitro* and *in vivo* (41), as well as for the formation of metal complexes (42). Transitional metals as Hg have a major role in the generation of oxygen free radicals in living organisms, in fact, the toxicity of HgCl₂ is attributed to the high affinity of Hg (II) to the thiol groups, which depletes cellular proteins such as GSH (43), and therefore increases the production of free radicals that can lead to lipid peroxidation, protein modification, and DNA damage. Compounds acting as chelators like the PAs used in this work, can inactivate metal ions and potentially inhibit the harmful processes related to mercury poisoning (44), which in turn can explain the results observed in the present study on the protective effect by scavenging free radicals. Phenolic compounds may inhibit and/or attenuate the damage generated by oxidative stress via several mechanisms, including the antioxidant and free-radical-scavenging activities and their ability to inhibit, activate, or protect

specific enzymes in the body (45). In a similar way, to prevent mercury-induced damage, natural products like plant extracts, essential oils, and edible species have been studied (46, 47); for example, Abarikwu *et al.* (46) demonstrated that oil extracted from *Moringa oleifera* diminishes the negative effects of mercury-induced oxidative stress and decreases malondialdehyde levels and superoxide dismutase and catalase activities in rats.

Mercury-induced oxidative and renal stress response is studied using different procedures (48) and, although CA, RA, and CGA have been evaluated by several authors using different test systems (49), in the present work we report for the first time the use of an *ex vivo* 3D kidney model, where we find results similar to those found in murine models. For example, researchers (50), observed that pretreatment with methanolic extracts from leaves of *Capparis spinosa*, which is rich in phenolic compounds, restores the healthy histology of renal tissue relative to the damage evident in Albino rats intoxicated with cisplatin. Our results are in accordance with reports that have demonstrated the protective effects of CA against the damage induced by paracetamol, ultraviolet light, hydrogen peroxide, lambda-cyhalothrin, ochratoxin A, and pilocarpine (51-56). It has also been reported that low doses of CA protect against endothelial damage caused by oxidative stress (57). Reportedly, CA protects against damage associated with ochratoxin A, tumor promoter 12-O-tetradecanoylphorbol-13-acetate (TPA), carbon tetrachloride, paraquat, hypochlorite, and hydrogen peroxide (55, 58-62). Similarly, RA protects against the damage caused by pilocarpine, aflatoxin B1, ochratoxin A, hydrogen peroxide, and doxorubicin (56, 63-65). All the reported findings were obtained analyzing murine models and/or 2D cell culture. The genoprotective effects of phenolic acid against HgCl₂-induced damage seem to be related to their anti-radical activity. This was also proposed by Rao and Sharma (66), who found that vitamin E protects against reproductive damage caused by HgCl₂ in mice. Rao *et al.* (67) also found that vitamin C protects against genotoxic damage induced by HgCl₂ in the peripheral blood, which they attributed to the antioxidant activity of the vitamin.

Conclusion

We have shown that application of the modified comet assay to 3D *ex vivo* models such as PCKS is a useful and complementary tool for determination of genotoxic damage induced by xenobiotic agents, and to evaluate antigenotoxic compounds. This study also shows that CA, RA, and CGA protect PCKS against mercury-induced genotoxic damage and that these compounds are not nephrotoxic. CA showed the best anti-radical activity, and also protected at the tissue level.

Acknowledgement

The authors thank Consejo Nacional de Ciencia y Tecnología (CONACyT) for postdoctoral fellowship to Irma Edith Carranza-Torres (CVU 209488). The results presented in this paper were from a student thesis that was partially supported by a grant from PAICYT/UANL: SA153-15. The authors also thank QBP Consuelo Coronado-Martínez for the excellent technical assistance.

Conflict of interest

All authors declared no conflict of interest in present manuscript.

References

- Plazar J, Hreljac I, Pirić P, Filipić M, Groothuis GM. Detection of xenobiotic-induced DNA damage by the comet assay applied to human and rat precision-cut liver slices. *Toxicol In Vitro* 2007; 21:1134-1142.
- Graaf IA, Groothuis GM, Olinga P. Precision-cut tissue slices as a tool to predict metabolism of novel drugs. *Expert Opin Drug Metab Toxicol* 2007;3: 879-898.
- Starokozhko V, Greupink R, van de Broek P, Soliman N, Ghimire S, de Graff IAM, *et al.* Rat precision-cut liver slices predict drug-induced cholestatic injury. *Arch Toxicol* 2007; 91:3403-3413.
- De Kanter R, De Jager MH, Draaisma AL, Jurva JU, Olinga P, Meijer DK, *et al.* Drug-metabolizing activity of human and rat liver, lung, kidney and intestine slices. *Xenobiotica* 2002; 32:349-362.
- Catania JM, Parrish AR, Kirkpatrick DS, Chitkara M, Bowden GT, Henderson CJ, *et al.* Precision-cut tissue slices from transgenic mice as an *in vitro* toxicology system. *Toxicol In Vitro* 2003; 17:201-205.
- Baverel G, Knouzy B, Gauthier C, El Hage M, Ferrier B, Martin G, *et al.* Use of precision-cut renal cortical slices in nephrotoxicity studies. *Xenobiotica* 2013; 43:54-62.
- de Graaf IA, Draaisma AL, Schoeman O, Fahy GM, Groothuis GM, Koster HJ. Cryopreservation of rat precision-cut liver and kidney slices by rapid freezing and vitrification. *Cryobiology* 2007; 54:1-12.
- Poosti F, Pham BT, Oosterhuis D, Poelstra K, van Goor H, Olinga P, *et al.* Precision-cut kidney slices (PCKS) to study development of renal fibrosis and efficacy of drug targeting *ex vivo*. *Dis Model Mech* 2015; 8:1227-1236.
- Liao W, McNutt MA, Zhu WG. The comet assay: a sensitive method for detecting DNA damage in individual cells. *Methods* 2009; 48:46-53.
- Langie SA, Azqueta A, Collins AR. The comet assay: past, present, and future. *Front Genet* 2015; 6:266.
- Plazar J, Filipić M, Groothuis GM. Antigenotoxic effect of Xanthohumol in rat liver slices. *Toxicol In Vitro* 2008; 22:318-327.
- Switalla S, Knebel J, Ritter D, Dasenbrock C, Krug N, Braun A, *et al.* Determination of genotoxicity by the Comet assay applied to murine precision-cut lung slices. *Toxicol In Vitro* 2013; 27:798-803.
- Jetten MJ, Claessen SM, Dejong CH, Lahoz A, Castell JV, van Delft JH, Kleinjans JC. Interindividual variation in response to xenobiotic exposure established in precision-cut human liver slices. *Toxicology* 2014; 323:61-69.
- Maser E, Schulz M, Sauer UG, Wiemann M, Ma-Hock L, Wohlleben W, Hartwig A, Landsiedel R. *In vitro* and *in vivo* genotoxicity investigations of differently sized amorphous SiO₂ nanomaterials. *Mutat Res Genet Toxicol Environ Mutagen* 2015; 794:57-74.
- Choi J, Chang JY, Hong J, Shin S, Park JS, Oh S. Low-level toxic metal exposure in healthy weaning-age infants: association with growth, dietary intake, and iron deficiency. *Int J Environ Res Publ Health* 2017; 14:388.
- Li H, Li H, Li Y, Liu Y, Zhao Z. Blood mercury, arsenic, cadmium, and lead in children with autism spectrum disorder. *Biol Trace Elem Res* 2018; 181:31-37.
- Wise JTF, Wang L, Zhang Z, Shi X. The 9th conference on metal toxicity and carcinogenesis: the conference overview. *Toxicol Appl Pharmacol* 2017; 331:1-5.
- Woods JS, Calas CA, Aicher LD. Stimulation of porphyrinogen oxidation by mercuric ion. II. Promotion of oxidation from the interaction of mercuric ion, glutathione, and mitochondria-

- generated hydrogen peroxide. *Mol Pharmacol* 1990; 38:261-266.
19. Carranza-Rosales P, Said-Fernandez S, Sepulveda-Saavedra J, Cruz-Vega DE, Gandolfi AJ. Morphologic and functional alterations induced by low doses of mercuric chloride in the kidney OK cell line: ultrastructural evidence for an apoptotic mechanism of damage. *Toxicology* 2005; 210:111-121.
 20. Schurz F, Sabater-Vilar M, Fink-Gremmels J. Mutagenicity of mercury chloride and mechanisms of cellular defense: the role of metal-binding proteins. *Mutagenesis* 2000; 15:525-530.
 21. Piccoli C, D'Aprile A, Scrima R, Ambrosi L, Zefferino R, Capitano N. Subcytotoxic mercury chloride inhibits gap junction intercellular communication by a redox- and phosphorylation-mediated mechanism. *Free Rad Bio Med* 2012; 52:916-927.
 22. Ramos Elizagaray SI, Soria EA. Arsenic immunotoxicity and immunomodulation by phytochemicals: potential relations to develop chemopreventive approaches. *Recent Pat Inflamm Allergy Drug Discov* 2014; 8:92-103.
 23. Karakaya S. Bioavailability of phenolic compounds. *Crit Rev Food Sci Nutr* 2004; 44:453-464.
 24. Caleja C, Ribeiro A, Barreiro MF, Ferreira IC. Phenolic compounds as nutraceuticals or functional food ingredients. *Curr Pharm Des* 2017; 23:2787-2806.
 25. Peyrol J, Riva C, Amiot MJ. Hydroxytyrosol in the prevention of the metabolic syndrome and related disorders. *Nutrients* 2017; 9:306.
 26. Viveros-Valdez E, Rivas-Morales C, Carranza-Rosales P, Mendoza S, Schmeda-Hirschmann G. Free radical scavengers from the Mexican herbal tea "poleo" (*Hedeoma drummondii*). *Z Naturforsch C* 2008; 63:341-346.
 27. Viveros-Valdez E, Rivas-Morales C, Oranday-Cárdenas A, Castro-Garza J, Carranza-Rosales P. Antiproliferative effect from the mexican poleo (*Hedeoma drummondii*). *J Med Food* 2010; 13:740-742.
 28. Petersen M, Simmonds MS. Rosmarinic acid. *Phytochemistry* 2003; 62:121-125.
 29. Jiang RW, Lau KM, Hon PM, Mak TC, Woo KS, Fung KP. Chemistry and biological activities of caffeic acid derivatives from *Salvia miltiorrhiza*. *Curr Med Chem* 2005; 12:237-246.
 30. Belkaid A, Currie JC, Desgagnés J, Annabi B. The chemopreventive properties of chlorogenic acid reveal a potential new role for the microsomal glucose-6-phosphate translocase in brain tumor progression. *Canc Cell Int* 2006; 6:12.
 31. Hymer WC, Kuff EL. Isolation of nuclei from mammalian tissues through the use of Triton X-100. *J Histochem Cytochem* 1964; 12:359-363.
 32. Singh NP, McCoy MT, Tice RR, Schneider EL. A simple technique for quantitation of low levels of DNA damage in individual cells. *Exp Cell Res* 1998; 175:184-191.
 33. Ostling O, Johanson KJ. Microelectrophoretic study of radiation-induced DNA damages in individual mammalian cells. *Biochem Biophys Res Commun* 1984; 123:291-298.
 34. Aninat C, Piton A, Glaise D, Le Charpentier T, Langouet S, Morel F, et al. Expression of cytochromes P450, conjugating enzymes and nuclear receptors in human hepatoma HepaRG cells. *Drug Metab Dispos* 2006; 34:75-83.
 35. Nelson LJ, Navarro M, Treskes P, Samuel K, Tura-Ceide O, Morley SD, et al. Acetaminophen cytotoxicity is ameliorated in a human liver organotypic co-culture model. *Sci Rep* 2015; 5:17455.
 36. Olinga P, Schuppan D. Precision-cut liver slices: a tool to model the liver ex vivo. *J Hepatol* 2013; 58:1252-1253.
 37. de Graaf IA, Olinga P, de Jager MH, Merema MT, de Kanter R., van de Kerkhof EG, et al. Preparation and incubation of precision-cut liver and intestinal slices for application in drug metabolism and toxicity studies. *Nat Protoc* 2010; 5:1540-1551.
 38. Olgún N, Müller ML, Rodríguez-Farré E, Suñol C. Neurotransmitter amines and antioxidant agents in neuronal protection against methylmercury-induced cytotoxicity in primary cultures of mice cortical neurons. *Neurotoxicology* 2018; pii:S0161-813X(18)30309-7.
 39. Simić A, Manojlović D, Segan D, Todorović M. Electrochemical behavior and antioxidant and prooxidant activity of natural phenolics. *Molecules* 2007; 12:2327-2340.
 40. Nemeikaite-Ceniene A, Imbrasaite A, Sergediene E, Cenas N. Quantitative structure-activity relationships in prooxidant cytotoxicity of polyphenols: role of potential of phenoxyl radical/phenol redox couple. *Arch Biochem Biophys* 2005; 441:182-190.
 41. De La Cruz JP, Ruiz-Moreno MI, Guerrero A, López-Villodres JA, Reyes JJ, Espartero JL, et al. Role of the catechol group in the antioxidant and neuroprotective effects of virgin olive oil components in rat brain. *J Nutr Biochem* 2015; 26:549-555.
 42. Andjelkovic M, Van Camp J, De Meulenaer B, Depaemelaere G, Socaciu C, Verloo M, et al. Iron-chelation properties of phenolic acids bearing catechol and galloyl groups. *Food Chem* 2006; 98:23-31.
 43. Augusti PR, Conterato GM, Somacal S, Sobieski R, Spohr PR, Torres JV, et al. Effect of astaxanthin on kidney function impairment and oxidative stress induced by mercuric chloride in rats. *Food Chem Toxicol* 2008; 46:212-219.
 44. Calabrese EJ, Iavicoli I, Calabrese V, Cory-Slechta DA, Giordano J. Elemental mercury neurotoxicity and clinical recovery of function: A review of findings, and implications for occupational health. *Environ Res* 2018; 163:134-148.
 45. Ferrari CK. Functional foods, herbs and nutraceuticals: towards biochemical mechanisms of healthy aging. *Biogerontology* 2004; 5:275-289.
 46. Abarikwu SO, Benjamin S, Ebah SG, Obilor G, Agbam G. Protective effect of Moringa oleifera oil against HgCl₂-induced hepato- and nephro-toxicity in rats. *J Basic Clin Physiol Pharmacol* 2017; 28:337-345.
 47. Joshi D, Mittal DK, Shukla S, Srivastav SK, Dixit VA. *Curcuma longa* Linn. extract and curcumin protect CYP 2E1 enzymatic activity against mercuric chloride-induced hepatotoxicity and oxidative stress: A protective approach. *Exp Toxicol Pathol* 2017; 69:373-382.
 48. Halliwell B, Whiteman M. Measuring reactive species and oxidative damage *in vivo* and in cell culture: how should you do it and what do the results mean? *Br J Pharmacol* 2004; 142:231-255.
 49. Sevgi K, Tepe B, Sarikurkcü C. Antioxidant and DNA damage protection potentials of selected phenolic acids. *Food Chem Toxicol* 2015; 77:33.
 50. Tlili N, Feriani A, Saadoui E, Nasri N, Khaldi A. *Capparis spinosa* leaves extract: Source of bioantioxidants with nephroprotective and hepatoprotective effects. *Biomed Pharmacother* 2017; 87:171-179.
 51. Janbaz KH, Saeed SA, Gilani AH. Studies on the protective effects of caffeic acid and quercetin on chemical-induced hepatotoxicity in rodents. *Phytomedicine* 2004; 11:424-430.
 52. Neradil J, Veselsk R, Slanina J. UVC-protective effect of caffeic acid on normal and transformed human skin cells *in vitro*. *Folia Biol. (Praha)* 2003; 49:197-202.
 53. Kang KA, Lee KH, Zhang R, Piao M, Chae S, Kim KN. Jeon, et al. Caffeic acid protects hydrogen peroxide induced cell damage in WI-38 human lung fibroblast cells. *Biol Pharm Bull* 2006; 29:1820-1824.
 54. Abdallah FB, Fetoui H, Fakhfakh F, Keskes L. Caffeic acid and quercetin protect erythrocytes against the oxidative stress and the genotoxic effects of lambda-cyhalothrin *in vitro*. *Biol Pharm Bull* 2011; 31:92-100.
 55. Cariddi LN, Sabini MC, Escobar FM, Montironi I, Mañas F, Iglesias D, et al. Polyphenols as possible bioprotectors against cytotoxicity and DNA damage induced by ochratoxin A. *Environ Toxicol Pharmacol* 2015; 39:1008-1018.

56. Coelho VR, Vieira CG, de Souza LP, da Silva LL, Pflüger P, Regner GG, *et al.* Behavioral and genotoxic evaluation of rosmarinic and caffeic acid in acute seizure models induced by pentylentetrazole and pilocarpine in mice. *Naunyn Schmiedeberg Arch Pharmacol* 2016; 389:1195-1203.
57. Migliori M, Cantaluppi V, Mannari C, Bertelli AAE, Medica D, Quercia AD, *et al.* Caffeic acid, a phenol found in white wine, modulates endothelial nitric oxide production and protects from oxidative stress-associated endothelial cell injury. *PLOS One* 2015; 10:21.
58. Huang MT, Smart RC, Wong CQ, Conney AH. Inhibitory effect of curcumin, chlorogenic acid, caffeic acid, and ferulic acid on tumor promotion in mouse skin by 12-O-tetradecanoylphorbol-13-acetate. *Cancer Res* 1998; 48:5941-5946.
59. Kapil A, Koul IB, Suri OP. Antihepatotoxic effects of chlorogenic acid from *Anthocephalus cadamba*. *Phytother Res* 1995; 9:189-193.
60. Tsuchiya T, Suzuki O, Igarashi K. Protective effects of chlorogenic acid on paraquat-induced oxidative stress in rats. *Biosci Biotechnol Biochem* 1996; 60:765-768.
61. Gugliucci A, Bastos DH. Chlorogenic acid protects paraoxonase 1 activity in high density lipoprotein from inactivation caused by physiological concentrations of hypochlorite. *Fitoterapia* 2009; 80:138-142.
62. Pavlica S, Gebhardt R. Protective effects of ellagic and chlorogenic acids against oxidative stress in PC12 cells. *Free Radic Res* 2005; 39:1377-1390.
63. Renzulli C, Galvano F, Pierdomenico L, Speroni E, Guerra MC. Effects of rosmarinic acid against aflatoxin B1 and ochratoxin-A-induced cell damage in a human hepatoma cell line (Hep G2). *J Appl Toxicol* 2004; 24:289-296.
64. Lee HJ, Cho HS, Park E, Kim S, Lee SY, Kim CS, *et al.* Rosmarinic acid protects human dopaminergic neuronal cells against hydrogen peroxide-induced apoptosis. *Toxicology* 2008; 250:109-115.
65. Furtado MA, de Almeida LC, Furtado RA, Cunha WR, Tavares DC. Antimutagenicity of rosmarinic acid in Swiss mice evaluated by the micronucleus assay. *Mutat Res* 2008; 657:150-154.
66. Rao MV, Sharma PS. Protective effect of vitamin E against mercuric chloride reproductive toxicity in male mice. *Reprod Toxicol* 2010; 15:705-712.
67. Rao MV, Chinoy NJ, Suthar MB, Rajvanshi MI. Role of ascorbic acid on mercuric chloride-induced genotoxicity in human blood cultures. *Toxicol In Vitro* 2001; 15:649-654.