

ORIGINAL ARTICLE

Iran J Allergy Asthma Immunol
June 2015; 14(3):292-297.

Plasma Adiponectin Concentrations and Adiponectin Gene Polymorphisms Are Associated with Bronchial Asthma in the Chinese Li Population

Yipeng Ding¹, Hua Yang², Haiwu He³, Huifang Shi³, Ping He¹, Shouchun Yan⁴, and Tianbo Jin^{2,5}

¹ Emergency Department of Hainan Provincial People's Hospital, Haikou, 570102, China

² School of Life Sciences, Northwest University, Xi'an 710069, China

³ Respiratory Department of Hainan Nongken Jeneral Hospital, Haikou, 570311 China

⁴ Emergency Department of First People's Hospital of Xi'an, Xi'an, 710002, China

⁵ National Engineering Research Center for Miniaturized Detection Systems, Xi'an 710069, China

Received: 25 August 2014; Received in revised form: 31 October 2014; Accepted: 22 November 2014

ABSTRACT

The purpose of this study was to determine the clinical significance of changes in the plasma adiponectin concentration in patients with bronchial asthma and to test the association between the single nucleotide polymorphisms (SNPs) rs2241766 and rs1501299 in the *ADIPOQ* gene and bronchial asthma in the Chinese Li population.

We selected 120 cases and 120 controls, and plasma adiponectin, interleukin (IL)-6, and tumor necrosis factor- α (TNF- α) levels were measured by enzyme-linked immunosorbent assay (ELISA). In addition, we genotyped two tag single nucleotide polymorphisms (tSNPs) and evaluated their association with bronchial asthma using the χ^2 test and genetic model analysis.

Compared to controls, patients with acute exacerbation of bronchial asthma showed significantly lower adiponectin and significantly higher IL-6 and TNF- α levels ($p < 0.01$). A positive association was found between the rs1501299 SNP and acute exacerbation (OR = 1.62; 95% CI = 1.08-2.43; $p = 0.019$).

The inverse correlation between the plasma adiponectin concentration and asthma exacerbation indicates that adiponectin may play a protective role in the pathogenesis of asthma. Meanwhile, our findings suggest that *ADIPOQ* polymorphisms influence the risk of developing bronchial asthma in Chinese Li population.

Keywords: Adiponectin; Asthma; Case-control studies

INTRODUCTION

Asthma is a serious public health problem in many

regions, including developing countries, where its incidence and mortality are increasing.¹ Over the last few decades, the prevalence of asthma has increased in many places around the world.² Asthma is a complex diseases involving complex interactions between multiple genes and environmental factors.^{3,4}

Adiponectin is a protein secreted from adipose

Corresponding Authors: Tianbo Jin, MD, PhD;
Mailbox: 386, #229 North Taibai Road, Xi'an 710069, Shaanxi,
China. Tel/Fax: (+86 29) 8830 3800, E-mail: tianbojin1973@gmail.com

tissue in abundant amounts into the blood circulation. In mice, adiponectin has been shown to attenuate allergen-induced airway inflammation and hyper-responsiveness via induction of anti-inflammatory factors.⁵⁻⁷ Other studies have suggested that asthma is associated with intermediate biological and physiological phenotypes, including plasma adiponectin, interleukin (IL)-6, and tumor necrosis factor- α (TNF- α). Investigation shows, Chinese Li population asthma prevalence was 3.38%, significantly higher than the 1.50% prevalence of asthma in China.^{8,9} To investigate the potential role of *ADIPOQ* in the pathogenesis of asthma in the Chinese Li population, we determined the levels of plasma adiponectin, IL-6, and TNF- α in 120 bronchial asthma patients and 120 healthy controls of the Li nationality from the Hainan Province of China and determined whether there is an association between exacerbation of symptoms and the *ADIPOQ* single nucleotide polymorphisms (SNPs) rs2241766 and rs1501299.

MATERIALS AND METHODS

Study Participants

We recruited 120 Chinese Li (58 females, 62 males; median age = 53.5 ± 6.9 years) diagnosed with bronchial asthma according to the guidelines developed by the Chinese Society of Respiratory Diseases Asthma Study Group in 1997. We used two different classifications for current asthma: 1) acute exacerbation was defined as episodes of recurrent wheezing, shortness of breath, chest tightness, or cough concurrent with a positive bronchodilator test; and 2) remission was defined as a diminishment of asthmatic symptoms and signs with or without treatment, restoration of lung function to the level before the onset of asthma attack, and retention of normal lung function for more than 2 weeks. The inclusion criteria for patient enrollment included no use of corticosteroids within the previous 1 week and no use of anti-asthma medications, such as a bronchodilator, within the 24 hours before blood sample collection.^{10,11} A total of 36 current asthma patients were classified as having acute asthma exacerbation and the remaining 84 patients were classified as being in remission. For control, 120 randomly selected healthy individuals (60 females, 60 males; median age 50.2 ± 5.8 years) were taken as a control group, without previous or present medical history of respiratory disorders or symptoms (persistent

cough, sputum production, or dyspnea), seen by their practitioner for regular checkup, were recruited consecutively as controls. All participants were from Li populations of Hainan province.

Blood Sample Collection and Testing

The Human Research Committee of the People's Hospital of Hainan Province for Approval of Research Involving Human Subjects approved the use of human samples in this study. We also obtained signed informed consent from each study participant. Fasting blood samples were obtained from the elbow vein from all study participants. Blood samples were centrifuged at 2,500 r/min for 5 min at 4°C, and the resulting plasma was stored at -20°C. The plasma concentrations of total adiponectin, IL-6, and TNF- α were measured by enzyme-linked immunosorbent assay (ELISA) (adiponectin: Huian Biotech, Shenzheng, China; IL-6 and TNF- α : Sengxiong Biotech, Shanghai, China).

DNA Preparation and Genotyping

DNA was extracted following the phenol-chloroform protocol. The rs2241766 and rs1501299 variants of *ADIPOQ* were genotyped by polymerase chain reaction (PCR) amplification and restriction fragment length polymorphism (RFLP) analysis. The primers used for PCR amplification of *ADIPOQ* were designed using PrimerPremier 5.0 software, and the primers were synthesized and sequenced by Boya Biotech (Shanghai, China). The sequences of the PCR primers and restriction enzymes used were as follows: CTGAGATGGACGGAGTCCTTT (forward) and CCAAATCACTTCAGGTTGCTT (reverse), with *Sma*I (45 T>G) and *Mva*I 269I (276 G>T). We selected two tag SNPs (tSNPs) in *ADIPOQ*. All products of DNA cleavage were analyzed by electrophoresis on a 2% agarose gel and visualized using GIS Gel Imaging System software.

Statistical Analysis

Microsoft Excel and SPSS 16.0 statistical package (SPSS, Chicago, IL) were used to perform statistical analyses. All *p*-values in this study were two-sided, and a *p*-value ≤ 0.05 was considered to indicate a significant difference. With the normality of data distribution for adiponectin, IL-6, and TNF- α . The frequency of each SNP in control subjects was validated by testing for departure from the Hardy-Weinberg equilibrium

(HWE) using an exact test. Differences in the tSNP genotype distribution between cases and controls were evaluated using the χ^2 test.¹² Unconditional logistic regression analysis was used to calculate the age-adjusted odds ratio (OR) and 95% confidence interval (CI).¹³ We evaluated the genetic association between genotype and bronchial asthma risk under different genetic models (co-dominant, dominant, recessive, over-dominant, and log-additive) using SNPstats, a website-based software available at <http://bioinfo.iconcologia.net>.¹⁴

RESULTS

Basic characteristics (e.g., gender, age, and pathology) of the cases and controls are listed in Table 1. No statistically significant differences were detected with respect to age, sex, and living environment

between the asthmatic and control groups, verifying the comparability of these two groups.

As shown in Table 2, adiponectin, IL-6, and TNF- α levels between cases and controls were compared by non-parametric Kruskal-Wallis test since adiponectin, IL-6, and TNF- α were not normally distributed in the outcome groups. Compared to the control group, patients with acute exacerbation of bronchial asthma showed significantly lower adiponectin and significantly higher IL-6 and TNF- α levels ($p < 0.01$). The level of adiponectin in patients in remission was significantly higher than that in patients with acute-stage asthma ($p < 0.01$) but was still lower than that in the controls ($p < 0.05$). The levels of both IL-6 and TNF- α in patients in remission were significantly lower than in patients with acute exacerbation ($p < 0.01$), but there were no differences in IL-6 and TNF- α levels between patients in remission and controls.

Table 1. Basic characteristics of cases and controls

Variables		Cases (n = 120)		Controls (n = 120)	
		no.	%	no.	%
Sex	Male	62	51.7	60	50
	Female	58	48.3	60	50
Age (years)	Median age	53.5 \pm 6.9		50.2 \pm 5.8	
Histologic type	acute asthma exacerbation	36	30		
	remission stage	84	70		

Table 2. Comparison of plasma concentrations of adiponectin, IL-6, and TNF- α between healthy controls, patients with acute exacerbation of asthma, and patients in asthma remission

Group	n	Adiponectin (ng/L)	IL-6 (ng/L)	TNF- α (ng/L)
Control	120	24.6 \pm 8.9	26.3 \pm 11.5	29.5 \pm 12.6
Exacerbation	36	11.7 \pm 6.2 ^a	56.4 \pm 19.3 ^a	62.8 \pm 18.1 ^a
Remission	84	19.3 \pm 7.5 ^{b,c}	30.7 \pm 11.4 ^c	35.6 \pm 14.6 ^c

Concentration values are expressed as mean \pm SD.

^aCompared with Control group, $p < 0.01$; ^bCompared with Control group, $p < 0.05$;

^cCompared with Exacerbation group, $p < 0.01$

Table 3. Association statistics for SNPs in ADIPOQ

SNPID	Base change	MAF (Case)	MAF (Control)	HWE p	OR	95% CI	p -value
rs2241766	T>G	0.30	0.30	0.08	1.02	0.69-1.51	0.921
rs1501299	G>T	0.33	0.23	0.44	1.62	1.08-2.43	0.019*

MAF: minor allele frequency; CI: confidence interval; OR: odds ratio.

* $P < 0.05$ indicates statistical significance.

Table 4. Association between *ADIPOQ* tSNPs and the risk of bronchial asthma, based on logistic tests

Model	Genotype	Control (n, %)	Case (n, %)	OR (95% CI)	p-value	AIC	BIC
Co-dominant	G/G	73 (60.8%)	61 (50.8%)	1.00	0.058	333	343.4
	G/T	39 (32.5%)	40 (33.3%)	1.23 (0.70-2.14)			
	T/T	8 (6.7%)	19 (15.8%)	2.84 (1.16-6.94)			
Dominant	G/G	73 (60.8%)	61 (50.8%)	1.00	0.12	334.3	341.2
	G/T-T/T	47 (39.2%)	59 (49.2%)	1.50 (0.90-2.51)			
Recessive	G/G-G/T	112 (93.3%)	101 (84.2%)	1.00	0.023*	331.5	338.5
	T/T	8 (6.7%)	19 (15.8%)	2.63 (1.10-6.28)			
Over-dominant	G/G-T/T	81 (67.5%)	80 (66.7%)	1.00	0.89	336.7	343.7
	G/T	39 (32.5%)	40 (33.3%)	1.04 (0.61-1.78)			
Log-additive	-	-	-	1.51 (1.04-2.20)	0.03*	332	339

MAF: minor allele frequency; OR: odd ratio; CI: confidence interval.

* $P < 0.05$ indicates statistical significance.

As shown in Table 3, two SNPs were genotyped in patients and controls. Both tSNPs were in HWE in the controls. We found that the rs1501299 tSNP in *ADIPOQ* was correlated with an increased risk of bronchial asthma at the 5% level (OR = 1.62; 95% CI, 1.08-2.43; $p = 0.019$).

We assumed that the minor allele of each tSNP was a risk factor compared with the wild-type allele. The results of age- and gender-adjusted unconditional logistic regression analyses of different genetic models are presented in Table 4. In the recessive model of rs1501299 in *ADIPOQ*, genotype "TT" was associated with increased bronchial asthma risk (OR = 2.63; 95% CI, 1.10-6.28; $p = 0.023$), whereas genotype "T" was associated with an increased risk in the log-additive model (OR = 1.51; 95% CI, 1.04-2.20; $p = 0.03$).

DISCUSSION

In our current study, patients with acute exacerbation of bronchial asthma showed significantly lower adiponectin and significantly higher IL-6 and TNF- α levels ($p < 0.01$). A positive association was found between the rs1501299 SNP in *ADIPOQ* and acute exacerbation (OR = 1.62; 95% CI = 1.08-2.43; $p = 0.019$). The inverse correlation between the plasma adiponectin concentration and asthma exacerbation indicates that adiponectin may play a protective role in the pathogenesis of asthma. Meanwhile, our findings suggest that *ADIPOQ* polymorphisms influence the risk of developing bronchial asthma in Hainan Chinese Li population.

The adiponectin gene spanned 17 kb on chromosome 3q27, consisting of three exons and two introns.¹⁵ *ADIPOQ* encodes a protein consisting of 244 amino acids, a signal peptide, a collagen-like domain at its N-terminus, and a globular domain at its C-terminus.

The adiponectin protein is exclusively secreted from adipose tissue, mainly the white adipose tissue. Adiponectin modulates a number of metabolic processes, including glucose regulation and fatty acid metabolism, through its anti-diabetic, anti-inflammatory, anti-atherogenic, and anti-proliferative activities.^{16,17} Impaired adiponectin multimerization due to mutation is usually the causative factor for asthma. Asthma is a complex syndrome, broadly defined as inflammation of the airways associated with airway hyper-responsiveness (AHR) and mucus hypersecretion.¹⁸ Studies have demonstrated that adipose tissues express multiple adipocytokines that may influence airway inflammation and airway remodeling.^{19,20} In a murine model of acute asthma, treatment with adiponectin was shown to attenuate airway inflammation and AHR.⁶ Eosinophils, mast cells, T lymphocytes, neutrophils, airway epithelial cells, and various other types of cells participate in the pathogenesis of asthma. These cells release a number of cellular products, including histamine, arachidonic acid metabolites, and inflammatory cytokines such as IL-6, IL-8, and TNF- α , which induce development of airway obstruction, AHR, and inflammation. More specifically, it has been shown that adiponectin suppresses the production of TNF- α and IL-6 in

lipopolysaccharide-activated human and porcine macrophages.^{21,22} Adiponectin also suppresses production of the pro-inflammatory cytokine interferon-gamma (IFN- γ) by human macrophages.²³ These studies indicate that adiponectin may be an inhibitory regulator of the immune system and thus may suppress the inflammatory responses associated with asthma.²⁴

We determined whether there is an association between two SNPs in *ADIPOQ* (rs2241766 and rs1501299) and the risk of bronchial asthma in people of the Li nationality residing in Hainan Province, China. Our findings suggest a positive association between the rs1501299 SNP in *ADIPOQ* and acute exacerbations of asthma. Although the functions of most *ADIPOQ* SNPs remain unclear, Genome-wide association studies have identified several potentially functional *ADIPOQ* variants.²⁵⁻²⁷ It has been shown that two common SNPs, rs2241766 and rs1501299, may affect susceptibility to a variety of disorders, including obesity,²⁸⁻³⁰ type 2 diabetes mellitus,³¹⁻³³ cardiovascular disease,³⁴ colon cancer,³⁵ and chronic obstructive pulmonary disease.^{36,37}

In conclusion, we examined the hypotheses that polymorphisms in *ADIPOQ* may affect susceptibility to bronchial asthma and that low APN levels may increase asthmatic inflammation in the Chinese Li population. To the best of our knowledge, this is the first study that has investigated the association between the *ADIPOQ* SNPs rs2241766 and rs1501299 and bronchial asthma risk in persons of Li nationality residing in the Hainan Province of China. Our results are different from many previous studies finding. We confirmed *ADIPOQ* as an important asthma susceptibility gene in our study. Moreover, we illustrated that the gene plays a key role in asthma. However, our results need to be verified through repeat of our experiments in other populations, and the mechanisms underlying the role of adiponectin in asthma also require further investigation.

ACKNOWLEDGEMENTS

We would like to thank the clinicians and other hospital staff who contributed to the blood sample and data collection for this study. We thank all of the individuals involved for their participation in this study. This study was supported by Hainan Natural Science Fund (2002-203-30222).

REFERENCES

1. Bousquet J, Bousquet PJ, Godard P, Daures JP. The public health implications of asthma. *Bull World Health Organ* 2005; 83(7):548-54.
2. Asher MI, Montefort S, Bjorksten B, Lai CK, Strachan DP, Weiland SK, et al. Worldwide time trends in the prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and eczema in childhood: ISAAC Phases One and Three repeat multicountry cross-sectional surveys. *Lancet* 2006; 368(9537):733-43.
3. Ober C. Perspectives on the past decade of asthma genetics. *J Allergy Clin Immunol* 2005; 116(2):274-8.
4. Thomsen SF, van der Sluis S, Kyvik KO, Skytthe A, Backer V. Estimates of asthma heritability in a large twin sample. *Clin Exp Allergy* 2010; 40(7):1054-61.
5. Arita Y, Kihara S, Ouchi N, Takahashi M, Maeda K, Miyagawa J, et al. Paradoxical decrease of an adipose-specific protein, adiponectin, in obesity. *Biochem Biophys Res Commun* 1999; 257(1):79-83.
6. Shore SA, Terry RD, Flynt L, Xu A, Hug C. Adiponectin attenuates allergen-induced airway inflammation and hyperresponsiveness in mice. *J Allergy Clin Immunol* 2006; 118(2):389-95.
7. Shore SA. Obesity, airway hyperresponsiveness, and inflammation. *J Appl Physiol* (1985) 2010; 108(3):735-43.
8. Ding YP, Yao HX, Tang XL, He HW, Shi HF, Lin L, Li M, Chen S, Chen J and Wang HJ. [Prevalence and risk factors of bronchial asthma among Li nationality in Hainan province]. *Zhonghua Yu Fang Yi Xue Za Zhi* 2011; 45(2):155-7.
9. Zhong NS. [The prevention, treatment and future study of asthma in China]. *Zhonghua Jie He He Hu Xi Za Zhi* 2005; 28(12):809-11.
10. Gao H, Xue Q, Lin Y, Wang L, Zhu G, Zhao, et al. Role of beta-adrenoceptor at different stages of bronchial asthma. *Chin Med J (Engl)* 2001; 114(12):1317-9.
11. Guo YS, Xu YP, Xu YH. [Expression of T-bet, GATA-3 and FOXP3 mRNA in asthmatic peripheral blood mononuclear cells stimulated by Dermatophagoides farinae antigen]. *Zhonghua Jie He He Hu Xi Za Zhi* 2007; 30(2):112-5.
12. Adamec C. [Example of the Use of the Nonparametric Test. Test X2 for Comparison of 2 Independent Examples]. *Cesk Zdrav* 1964; 12:613-9.
13. Bland JM, Altman DG. Statistics notes. The odds ratio. *BMJ* 2000; 320(7247):1468.

14. Sole X, Guino E, Valls J, Iniesta R, Moreno V. SNPStats: a web tool for the analysis of association studies. *Bioinformatics* 2006; 22(15):1928-9.
15. Takahashi M, Arita Y, Yamagata K, Matsukawa Y, Okutomi K, Horie M, et al. Genomic structure and mutations in adipose-specific gene, adiponectin. *Int J Obes Relat Metab Disord* 2000; 24(7):861-8.
16. Viengchareun S, Zennaro MC, Pascual-Le Tallec L, Lombes M. Brown adipocytes are novel sites of expression and regulation of adiponectin and resistin. *FEBS Lett* 2002; 532(3):345-50.
17. Hotta K, Funahashi T, Bodkin NL, Ortmeyer HK, Arita Y, Hansen BC, et al. Circulating concentrations of the adipocyte protein adiponectin are decreased in parallel with reduced insulin sensitivity during the progression to type 2 diabetes in rhesus monkeys. *Diabetes* 2001; 50(5):1126-33.
18. Anderson WJ, Watson L. Asthma and the hygiene hypothesis. *N Engl J Med* 2001; 344(21):1643-4.
19. Shore SA, Fredberg JJ. Obesity, smooth muscle, and airway hyperresponsiveness. *J Allergy Clin Immunol* 2005; 115(5):925-7.
20. Aaron SD, Fergusson D, Dent R, Chen Y, Vandemheen KL, Dales RE. Effect of weight reduction on respiratory function and airway reactivity in obese women. *Chest* 2004; 125(6):2046-52.
21. Wulster-Radcliffe MC, Ajuwon KM, Wang J, Christian JA, Spurlock ME. Adiponectin differentially regulates cytokines in porcine macrophages. *Biochem Biophys Res Commun* 2004; 316(3):924-9.
22. Yokota T, Oritani K, Takahashi I, Ishikawa J, Matsuyama A, Ouchi N, et al. Adiponectin, a new member of the family of soluble defense collagens, negatively regulates the growth of myelomonocytic progenitors and the functions of macrophages. *Blood* 2000; 96(5):1723-32.
23. Wolf AM, Wolf D, Rumpold H, Enrich B, Tilg H. Adiponectin induces the anti-inflammatory cytokines IL-10 and IL-1RA in human leukocytes. *Biochem Biophys Res Commun* 2004; 323(2):630-5.
24. Baek HS, Kim YD, Shin JH, Kim JH, Oh JW, Lee HB. Serum leptin and adiponectin levels correlate with exercise-induced bronchoconstriction in children with asthma. *Ann Allergy Asthma Immunol* 2011; 107(1):14-21.
25. Heid IM, Henneman P, Hicks A, Coassin S, Winkler T, Aulchenko YS, et al. Clear detection of ADIPOQ locus as the major gene for plasma adiponectin: results of genome-wide association analyses including 4659 European individuals. *Atherosclerosis* 2010; 208(2):412-20.
26. Jee SH, Sull JW, Lee JE, Shin C, Park J, Kimm H, et al. Adiponectin concentrations: a genome-wide association study. *Am J Hum Genet* 2010; 87(4):545-52.
27. Ling H, Waterworth DM, Stirnadel HA, Pollin TI, Barter PJ, Kesaniemi YA, et al. Genome-wide linkage and association analyses to identify genes influencing adiponectin levels: the GEMS Study. *Obesity (Silver Spring)* 2009; 17(4):737-44.
28. Park JW, Park J, Jee SH. ADIPOQ Gene Variants Associated with Susceptibility to Obesity and Low Serum Adiponectin Levels in Healthy Koreans. *Epidemiol Health* 2011; 33:e2011003.
29. Guzman-Ornelas MO, Chavarria-Avila E, Munoz-Valle JF, Armas-Ramos LE, Castro-Albarran J, Aguilar Aldrete ME, et al. Association of ADIPOQ +45T>G polymorphism with body fat mass and blood levels of soluble adiponectin and inflammation markers in a Mexican-Mestizo population. *Diabetes Metab Syndr Obes* 2012; 5:369-78.
30. Yuksel H, Sogut A, Yilmaz O, Onur E, Dinc G. Role of adipokines and hormones of obesity in childhood asthma. *Allergy Asthma Immunol Res* 2012; 4(2):98-103.
31. Li Y, Li X, Shi L, Yang M, Yang Y, Tao W, et al. Association of adiponectin SNP+45 and SNP+276 with type 2 diabetes in Han Chinese populations: a meta-analysis of 26 case-control studies. *PLoS One* 2011; 6(5):e19686.
32. Melistas L, Mantzoros CS, Kontogianni M, Antonopoulou S, Ordovas JM, Yiannakouris N. Association of the +45T>G and +276G>T polymorphisms in the adiponectin gene with insulin resistance in nondiabetic Greek women. *Eur J Endocrinol* 2009; 161(6):845-52.
33. Boer-Martins L, Figueiredo VN, Demacq C, Martins LC, Consolin-Colombo F, Figueiredo MJ, et al. Relationship of autonomic imbalance and circadian disruption with obesity and type 2 diabetes in resistant hypertensive patients. *Cardiovasc Diabetol* 2011; 10:24.
34. Menzaghi C, Trischitta V, Doria A. Genetic influences of adiponectin on insulin resistance, type 2 diabetes, and cardiovascular disease. *Diabetes* 2007; 56(5):1198-209.
35. Al-Harithy RN, Al-Zahrani MH. The adiponectin gene, ADIPOQ, and genetic susceptibility to colon cancer. *Oncol Lett* 2012; 3(1):176-80.
36. Yuan Y, Jiang H, Kuang J, Hou X, Feng Y, Su Z. Genetic variations in ADIPOQ gene are associated with chronic obstructive pulmonary disease. *PLoS One* 2012; 7(11):e50848.
37. Fasshauer M, Paschke R, Stumvoll M. Adiponectin, obesity, and cardiovascular disease. *Biochimie* 2004; 86(11):779-84.