Iran J Allergy Asthma Immunol October 2015; 14(5):543-544.

Prevalence of Specific IgE to Wheat Flour Allergens in Romanian Pediatric Population

Gabriel Samasca^{1,2}, Genel Sur³, Mihaela Iancu⁴, Iulia Lupan⁵, and Diana Deleanu^{1,6}

¹ Department of Immunology and Allergology, Iuliu Hatieganu University of Medicine and Pharmacy, Cluj-Napoca, Romania

² Emergency Hospital for Children, Cluj-Napoca, Romania

³ Department of Pediatrics II, Iuliu Hatieganu University of Medicine and Pharmacy, Cluj-Napoca, Romania

⁴ Department of Medical Informatics and Biostatistics, Iuliu Hatieganu University of Medicine

and Pharmacy, Cluj-Napoca, Romania

⁵ Department of Molecular Biology and Biotechnology, Babes-Bolyai University, Cluj-Napoca, Romania ⁶ Department of Internal Medicine, Regional Institute of Gastroenterology and Hepatology, Prof. Dr. Octavian Fodor Urgent Care Clinic, Romania

Received: 29 January 2015; Received in revised form: 26 March 2015; Accepted: 25 April 2015

INTRODUCTION

This is a short report on the incidence of wheatspecific IgE allergens in the pediatric population of Romania.

MATERIALS AND METHODS

The present study was conducted from 2012 to 2013 on 1097 children to determine the frequency of wheat-specific IgE antibodies as determined using EUROIMMUN immunoblot tests. The calculation of the prevalence of wheat-specific sIgE antibodies did not require study inclusion criteria for patients. This allowed performance of testing on more children to calculate the prevalence of these antibodies in the pediatric population as a whole.

The gender distribution was 59.7% males and 40.3% females. The test used to detect wheat-specific IgE antibodies was an *in vitro* semi-quantitative

Keywords: Allergens; Epidemiology; Food allergy; IgE

Corresponding Author: Gabriel Samasca, PhD;

method for the detection of inhaled allergens or food allergens in serum or plasma. The measuring range and implicit results are represented as enzyme-allergosorbent test scores of 0 to 6 according to manufacturer recommendations. A score of 0 signifies no spectifc antibody detected (concentration < 0.35 kU/l); 1 signifies very low detection of antibodies and sensitization, often without clinical symptoms (concentration 0.35 - 0.7 kU/l);.2 signifies low antibody titer and sensitization, often with noticeable clinical symptoms (concentration 0.7- 3.5 kU/l); 3 signifies significant antibody titer, clinical symptoms usually present (concentration 3.5-17.5 kU/l); 4 signifies high antibody titer, with clinical symptoms (concentration 17.5 - 50 kU/l); 5 signifies very high antibody titer (concentration 50 - 100 kU/l); 6 signifies extremely high antibody titer (concentration> 100 kU/l).

The prevalence of sensitization was calculated to be f4 according to class 1 positivity, where positivity shows no clinical symptoms to which the body responds. The prevalence of allergy was calculated as f4 for classes 2, 3, 4, 5, and 6, where positivity implies the presence of clinical symptoms to which the body responds.

"Sensitization" is the body producing IgE that develops specifically for that particular allergen, but without experiencinge a reaction.^{1,2} The immune

Copyright© Autumn 2015, Iran J Allergy Asthma Immunol. All rights reserved.

Published by Tehran University of Medical Sciences (http://ijaai.tums.ac.ir)

Department of Immunology, Iuliu Hațieganu University of Medicine and Pharmacy, Cluj-Napoca, Romania. Tel: (+40 74) 0252 795; E-mail: Gabriel.Samasca@umfcluj.ro

reponse of an allergy begins with sensitization.^{3,4}

The present study focused on the following allergens: f4 wheat flour, f49 apple, f35 potato, f31 carrot, f17 hazelnut, f13 peanut, f14 soy, f9 rice, e204 bovine serum albumin, f78 casein, f77 beta-lactoglobulin, f76 alpha-lactalbumin, f3 fish, f2 cow's milk, f75 egg yolk, f1 egg white, m6 Alternaria alternata, m3 Aspergillus fumigatus, m2 Cladosporium herbarum, d2 Dermatophagoides farinae, d1 Dermatophagoides Pter, and gx mixture of herbs.

RESULTS

No statistically significant association between patient gender and the presence of f4 was observed $(\chi^2=0.27, df = 1, n = 1097, p = 0.61)$. The sIgE antibody detected as f4 was not present in 1012 patients. Very low sIgE antibody titers for f4 (frequently no clinical symptoms where sensitization is present) were detected in 65 patients. Low sIgE antibody titers for f4 (sensitization, frequently with clinical symptoms in the upper range of class) were detected in 13 patients. Significant sIgE antibody titers for f4 (clinical symptoms usually present) were detected in 5 patients. Very high sIgE antibody titers for f4 were detected in 2 patients. The prevalence of sIgE as sensitization to f4 according to class 1 positivity was 5.92% and the prevalence of sIgE to allergy to f4 according to classes 2, 3, 4, 5, and 6 positivity was 1.82%.

No statistically significant differences between class frequencies for f4 and patient gender was observed (Fisher's exact test, test statistic = 1.62, df = 4, n= 1097, P= 0.85). There was a weak to moderate positive association between f4 and the following parameters f49, f35, f31, f17, f14, f9, e204, f78, f77, f76, f3, f2, f75, f1, m6, m2, d2 and gx (correlation coefficient p<0.30, p<0.001). No association was observed

between f4 and m3 (p=0.13) or d1 (Spearman correlation coefficient: p = 0.06.

CONCLUSION

Patients showing sensitization to wheat flour should be considered at risk for allergy to wheat flour. Male gender was not a predisposing factor for wheat flour allergy. The following possible triggers for wheat flour sensitization and wheat flour allergy were noted for allergens that have not been reported thus far: mites, *Dermatophagoides farinae*, molds, *Alternaria alternata*, and *Cladosporium herbarum*.

ACKNOWLEDGEMENTS

This paper was published under the framework of the European Social Fund, Human Resources Development Operational Programme 2007-2013, project no. POSDRU/159/1.5/S/138776.

REFERENCES

. IgE's Role in Allergic Asthma https://www.aafa.org/display.cfm?id=8&sub=16&cont=54.

- Kekki OM, Scheynius A, Poikonen S, Koskinen A, Kautiainen H, Turjanmaa K. Sensitization to Malassezia in children with atopic dermatitis combined with food allergy. Pediatr Allergy Immunol 2013; 24(3):244-9.
- 3. IgE in Clinical Allergy and Allergy Diagnosis http://www.worldallergy.org/professional/allergic_diseas es_center/ige.
- Mansouri M, Pourpak Z, Mozafari H, Abdollah Gorji F, Shokouhi Shoormasti R. Follow-up of the wheat allergy in children; consequences and outgrowing the allergy. Iran J Allergy Asthma Immunol 2012; 11(2):157-63.

Iran J Allergy Asthma Immunol, Autumn 2015/544 Published by Tehran University of Medical Sciences (http://ijaai.tums.ac.ir)