

Novel ionic liquid 1,3-disulfonic acid imidazolium hydrogen sulfate {[Dsim]HSO₄} efficiently catalyzed *N*-boc protection of amines

M.A. Zolfigol^{a,*}, V. Khakyzadeh^a, A.R. Moosavi-Zare^a, G. Chehardoli^b, F. Derakhshan-Panah^a, A. Zare^c, O. Khaledian^a

^a Faculty of Chemistry, Bu-Ali Sina University, Hamedan, 6517838683, Iran

^b School of Pharmacy, Hamedan University of Medical Sciences, Zip Code 65178, Hamedan, Iran

^c Department of Chemistry, Payame Noor University, Tehran, P.O. Box: 19395-4697, Iran

Received 20 January 2012; revised 18 March 2012; accepted 22 May 2012

KEYWORDS

Ionic liquid; 1,3-Disulfonic acid imidazolium hydrogen sulfate {[Dsim]HSO₄}; *N*-boc protection; Amine. **Abstract** A simple and highly efficient method for the protection of amines from their corresponding *N*-boc derivatives is reported. It is interesting that only 0.1 mol % of novel ionic liquid 1,3-disulfonic acid imidazolium hydrogen sulfate {[Dsim]HSO₄} can efficiently catalyze the reaction. Clean reaction, very short reaction times, high yields, easy preparation and high TOF (turn over frequency) values of the catalyst in comparison with reported catalysts are some advantages of this work.

© 2012 Sharif University of Technology. Production and hosting by Elsevier B.V. All rights reserved.

1. Introduction

lonic Liquids (ILs) have received considerable interest as eco-friendly solvents, catalysts and reagents in the context of green synthesis, because of their unique properties, such as low volatility, non-flammability, high thermal stability, negligible vapor pressure and ability to dissolve a wide range of materials. Among them, ILs exhibiting a Brønsted acid character possess useful characteristics of solid acids and mineral liquid acids. Such ILs have been designed to replace traditional mineral liquid acids, like sulfuric acid and hydrochloric acid, in a range of chemical transformations [1,2]. In continuation of our ongoing program on the synthesis and applications of acidic ILs in organic synthesis, we have recently introduced a new category of ionic liquids, namely Sulfonic Acid Functionalized Imidazolium Salts (SAFIS) [3–6]. This class of ILs has been used as an efficient catalyst and reagent in organic transformations,

* Corresponding author. Fax: +98 811 8272404.

E-mail addresses: zolfi@basu.ac.ir, mzolfigol@yahoo.com (M.A. Zolfigol). Peer review under responsibility of Sharif University of Technology.

ELSEVIER Production and hosting by Elsevier

such as synthesis of bis(indolyl)methans [3], preparation of N-sulfonyl imines [4], nitration of phenols [5], synthesis of 1-amidoalkyl-2-naphthols [6], preparation of benzimidazoles [7], and chemoselective trimethylsilyl protection of hydroxyl groups [8]. In this work, we synthesize another analogue of SAFIS, namely, 1,3-disulfonic acid imidazolium hydrogen sulfate {[Dsim]HSO4} (as an ionic liquid), *via* the reaction of 1,3-disulfonic acid imidazolium chloride {[Dsim]Cl}, with H_2SO_4 (99.99%), which exhibits many interesting properties (Scheme 1).

Functional group protection and deprotection methods are important in the synthesis of the target molecule. The protection of amines is one of the most fundamental and useful transformations in organic synthesis, especially in peptide synthesis [9,10].

Several reagents and methods have been used for N-*tert*butoxycarbonylation of amines [11–13]. These methodologies suffer from a variety of drawbacks, such as long reaction times, low yields, high toxicity and the use of expensive reagents. Commercially available *di-tert*-butyl dicarbonate (*diboc*) is an efficient reagent for clean and rapid introduction of the bocprotecting group [14–17].

Herein, we report the selective *tert*-butoxycarbonylation of various amines, using *diboc* catalyzed by 1,3-disulfonic acid imidazolium hydrogen sulfate {[Dsim]HSO₄} as a novel

1026-3098 © 2012 Sharif University of Technology. Production and hosting by Elsevier B.V. All rights reserved. doi:10.1016/j.scient.2012.05.003



Scheme 1: The preparation of 1,3-disulfonic acid imidazolium hydrogen sulphate.

Scheme 2: Selective *tert*-butoxycarbonylation of various amines using *diboc* catalyzed by [Dsim]HSO₄.



Figure 1: TG/DTG diagrams of the catalyst.

ionic liquid, at room temperature with high yields and very short reaction times. Interestingly, this method has none of the above-mentioned drawbacks for the N-boc protection of amines at all (Scheme 2).

2. Results and discussion

The structure of 1,3-disulfonic acid imidazolium hydrogen sulfate was identified by IR, ¹H and ¹³C NMR and mass spectra.

The IR spectrum of the catalyst shows a broad peak at $3100-3400 \text{ cm}^{-1}$ related to the OH in the SO₃H group. Moreover, the two peaks observed at 1085 cm^{-1} and 1285 cm^{-1} correspond to vibrational modes of the N–SO₂ bond.

The ¹H NMR spectrum of [Dsim]HSO₄ show the unmistaken acidic hydrogen (SO₃H) peaks at 13.55 and 11.95 ppm. To prove that this peak is really related to the hydrogen of SO₃H in the compound, we also run ¹H NMR of [Dsim]Cl and H₂SO₄ in DMSO $- d_6$. In these spectra, the peaks of the two acidic hydrogen groups of [Dsim]HSO₄, [Dsim]Cl and H₂SO₄ were observed at 13.55 and 11.95, 13.34, and 12.54 ppm, respectively. This observation showed that [Dsim]HSO₄ was exactly synthesized. Other details regarding the NMR signals are provided in the experimental section.

The mass spectrum of the compound gave the correct molecular ion peak at 326. The thermogram analysis of the catalyst was studied at a range of temperatures between 25 to 600 °C. The thermogram (TG) of the catalyst shows two weight losses (Figure 1). The first strong loss weight was observed after 350 °C and the second strong weight loss appears after 540 °C. Therefore, the molecular decomposition occurred for this catalyst after 350 °C.

To optimize the reaction conditions, as a model, the reaction of 4-methoxyaniline with *diboc* (Scheme 3) was tested in

Table 1: Effect of different	amounts of the	catalyst on	the reaction of	4-
methoxyaniline with diboc.				

Catalysts	Mol% of	catalyst	Temp.(°C)	Time(min)	Yield ^a (%)
-	-		r.t.	60	20
H_2SO_4	0.1		r.t.	60	23
H_2SO_4	1		rat.	60	31
[Dsim]Cl	0.1		r.t.	15	88
[Dsim]Cl	1		r.t.	10	90
[Dsim]HSO ₄	0.1		r.t.	5	98
[Dsim]HSO ₄	1		r.t.	5	98
^a Isolated yield.					
^a Isolated yield.					



Scheme 3: The reaction of 4-methoxyaniline with diboc catalyzed by [Dsim]HSO₄.

Table 2: The effect of various solvents on the reaction of 4-methoxyaniline
with <i>diboc</i> catalyzed by [Dsim]HSO ₄ .

Entry	Solvent	Time (min)	Yield ^a (%)		
1	EtOH	5	98		
2	CH ₃ CN	15	88		
3	CHCl ₃	18	90		
4	CH_2Cl_2	15	90		
5	n-hexane	20	87		
6	Acetone	12	89		
3 7 1 4 1 1 1					

^a Isolated yield.

the presence of different amounts of $[Dsim]HSO_4$ at room temperature in ethanol as solvent. The results are summarized in Table 1. To prove that $[Dsim]HSO_4$ was completely synthesised, and [Dsim]Cl or H_2SO_4 converted to $[Dsim]HSO_4$, the model reaction was also performed in different amounts of [Dsim]Cl or H_2SO_4 . As it can be seen in Table 1, the best results were obtained when the reaction was carried out using 0.1 mol% of $[Dsim]HSO_4$ at room temperature.

In another study, we studied the reaction of 4-methoxyaniline (1 mmol) to its corresponding N-boc protected derivative, using [Dsim]HSO₄ as catalyst (0.1 mol %) and *diboc* (1 mmol) in a variety of solvents, at room temperature (Table 2). The results showed that absolute ethanol is the best solvent in terms of time and product yield (Table 2).

To explore the generality of the catalyst, a wide range of amines underwent N-boc protection using this procedure to provide the corresponding *tert*-butoxycarbonylated products in good to excellent yields. We also applied our reaction conditions to the N-boc protection of some aliphatic amines. The results were summarized in Table 3.

In a plausible mechanism, which is shown in Scheme 4, we suggest that the carbonyl group of *diboc* is activated by

Entry	Amine	Product	Time	Yield ^a (%)	Ref.
	HO	Boc			
1	NH NH	С N OH	3 min	93	[13]
	\bigcap				
2	HN	N⁄ Boc	1 min	99	[13]
	NH ₂	H N. Boc			
3			15 min	90	[15]
	OH				
4	H ₂ N	HN V Boc	4 min	99	[12]
	NH₂	Boc NH			
_					[40]
5	011	Boc	33 min	85	[13]
	H ₂ N	HN			
6	~~~0×		5 min	98	[13]
		HN			
7	$H_2N \sim$	Boc	30 min	95	[13]
0	H ₂ N	Boc	30 min	95	[14]
0	NH ₂		20 11111	60	[14]
9		N Boc	2 h	75	[13]
10	H ₂ N	HN' V NH Boc Boc	12 min	70	[13]
	NH ₂	CI			
11	CI	N Boc H	10 min	70	[14]
	NH ₂	Br			
12	Br	N Boc H	15 min	90	[14]
	NHa	Boc			
13	HaN. A Br	H N A	12 h	50	[13]
14		Boc	35 min	55	[13]
	-	•			[]



Scheme 4: The purposed mechanism for the protection of amines with *diboc* using [Dsim]HSO₄ as catalyst.

Catalyst/condition	Catalyst amount (mol%)	Time (min)	Yield ^a (%)	TON ^b	TOF ^c (min ⁻¹)	Ref.
[Dsim]HSO ₄ /ethanol/r.t.	0.1	15	90	900	60	_d
Thiourea/toluene/60 °C	10	40	95	9.5	0.23	[11]
NH ₂ SO ₃ H/solvent-free/r.t.	5	10	96	19.2	1.92	[12]
NH ₂ SO ₃ H/solvent-free/r.t.	5	5	99	19.8	3.96	[12]
[H ₂ -cryptand 222](Br ₃) ₂	0.1	300	80	800	2.66	[13]
$Cu(BF_4)_2 \cdot xH_2O/solvent-free/35 °C$	1	5	98	98	19.6	[14]
[(HMIm)BF ₄]/Solvent-free/35 °C	1	15	100	100	6.66	[15]
La(NO ₃) ₃ · 6H ₂ O/solvent-free/35 °C	5	2	100	20	10	[16]
Thiourea/toluene/60 °C	10	40	95	9.5	0.23	[17]

Table 4: Comparison of the results of the reaction of aniline with *diboc* catalyzed by [Dsim]HSO₄ with those obtained by the reported catalysts.

^a Isolated yield.

^b Turn-over number.

^c Turn-over frequency.

^d Our work.



Figure 2: The reaction of 4-methoxyaniline with *diboc* in the presence of reused $[Dsim]HSO_4$ at room temperature.

hydrogen bond formation with the acidic hydrogens of [Dsim]HSO₄. Afterward, the nitrogen of the amine attacks the carbonyl group. By opening and closing the Π bond of the carbonyl group, the amine is protected, and one molecule of CO₂ and *tert*-butyl alcohol is also prepared.

In another study, recyclability of the catalyst was tested upon the reaction of the N-boc protection of 4-methoxyaniline. The reaction mixture was extracted by ethyl acetate and separated from the catalyst. Afterward, the reused catalyst was employed for another reaction. We observed that the catalytic activity of the catalyst was restored within the limits of experimental error for five successive runs (Figure 2).

Furthermore, we compared the Turn Over Number (TON) and Frequency (TOF) values of our catalyst with the reported catalysts on the reaction of aniline with *diboc* (Table 4). The TON and TOF values showed that [Dsim]HSO₄ was more effective than the reported catalysts.

3. Conclusion

In summary, we have prepared, characterized and applied a novel ionic liquid, namely; 1,3-disulfonic acid imidazolium hydrogen sulfate {[Dsim]HSO₄}, as a reusable catalyst for the chemoselective N-boc protection of amines. The reaction was carried out at room temperature with high yields and very short reaction times.

4. Experimental

All Chemicals were purchased from Merck, Sigma-Aldrich[®] or Fluka Chemical Company. The products were identified by

comparison of their ¹H NMR, TLC and melting points with those in authentic samples. The progress of the reactions was monitored by TLC using silica gel SIL G/UV 254 plates. The ¹H NMR (300 or 90 MHz) was run on a Bruker Avance DPX FT-NMR spectrometer (δ in ppm). Melting points were recorded on a Stuart Scientific Apparatus SMP3 (UK) in open capillary tubes.

4.1. Procedure for the preparation of ionic liquid [Dsim]HSO₄ (Scheme 1)

Chlorosulfonic acid (1.1885 g, 10.2 mmol) was added dropwise to a round-bottomed flask (100 mL) containing imidazole (0.340 g, 5 mmol) in dry CH₂Cl₂ (50 mL), over a period of 20 min at room temperature. After the addition was completed, the reaction mixture was stirred for 12 h under pressure of nitrogen (to remove the produced HCl), stood for 5 min, and the CH₂Cl₂ was decanted. The residue was washed with dry CH_2Cl_2 (3 × 50 mL) and dried under vacuum to give [Dsim]Cl as a viscous pale yellow oil in 95% yield, 1.257 g [6]. Then, sulfuric acid 99.99% (0.49 g, 5 mmol) was added dropwise to [Dsim]Cl (1.3233 g, 5 mmol) over a period of 5 min at room temperature under a continuous flow of nitrogen to remove the HCl gas that is produced. The resulting mixture was stirred for 12 h at 60 °C and then heated at 90 °C under powerful vacuum for 20 min to give [Dsim]HSO₄ as a viscous green oil in 99% vield.

1,3-disulfonic acid imidazolium hydrogen sulfate{[Dsim]HSO₄}:

Brown oil; IR (Nujol) 624, 1031, 1053, 1085, 1285, 1324, 3100–3400 cm⁻¹; ¹H NMR(300 MHz, DMSO – d_6), δ (ppm) 7.22(*s*, 2H), 8.44(*s*, 1H), 11.95(*s*, 1H), 13.55(*s*, 2H); ¹³C NMR (75 MHz, DMSO – d_6) : δ (ppm)119.5, 134.0; MS : m/z = 326(M^+), 229(M^+ – HSO₄), 148(M^+ – HSO₄SO₃H), 97(HSO₄), 67(M^+ – HSO₄SO₃HSO₃H).

4.2. General procedure for the N-boc protection of amines

A solution of *diboc* (1 mmol) in ethanol (2 mL) was added to a round-bottomed flask containing [Dsim]HSO₄ (0.1 mol%), and the resulting solution was stirred at room temperature for 10 s. Then, amine (1 mmol) was added, and the reaction mixture was stirred at room temperature for the appropriate time (see Table 3). Afterwards, the solution of the product and the unreacted starting materials were separated from [Dsim]HSO₄ by decanting (the catalyst could be used for another reaction). After drying and evaporation of the solvent, the solid residue (crude product) was purified by recrystallization from *n*-hexane or ethyl acetate (5 mL) or by column chromatography on silica gel eluted with ethyl acetate/*n*-hexane (1/3).

4.3. Selected spectral data of the products

Tert-butyl 4-(2-*hydroxyethyl*)*piperazine*-1-*carboxylate* (1): (lit. [13]); IR (Nujol): 3450, 2976, 1700 cm⁻¹; ¹H NMR (90 MHz, CDCl₃), δ 1.09(*s*, 9H), 2.11(*m*, 6H), 3.08(*m*, 4H), 3.30(*t*, 2H), 3.68(*br*, 1H).

Tert-butyl 4-*hydroxyphenylcarbamate* (4): (lit. [13]); IR (KBr): 3360, 2969, 1705, 1535 cm⁻¹; ¹H NMR (90 MHz, CDCl₃), δ 1.48(*s*, 9H), 5.35(*s*, 1H), 6.36(*s*, 1H), 6.60–6.82(*m*, 2H), 7.02 –7.35(*m*, 2H).

Tert-butyl 4-*methoxyphenylcarbamate* (6): (lit. [15]); IR (KBr): 3420, 2970, 1693, 1535 cm⁻¹. ¹H NMR(90 MHz, CDCl₃), δ 1.47(*s*, 9H), 3.73(*s*, 3H), 6.42(*s*, 1H), 6.65–6.96(*m*, 2H), 7.15–7.32(*m*, 2H).

Tert-butyl m-tolylcarbamate (8): (lit. [14]); IR (KBr): 3327, 2960, 1698, 1533 cm⁻¹; ¹H NMR(90 MHz, CDCl₃), δ 1.52(*s*, 9H), 2.30(*s*, 3H), 6.62–7.36(*m*, 5H).

Tert-butyl4,4'-methylenebis (4,1-phenylene) dicarbamate (10): (lit. [13]); IR (KBr): 3325, 2960, 1683, 1539 cm⁻¹; ¹H NMR(90 MHz, CDCl₃) : δ 1.49(*s*, 18H), 3.85(*s*, 2H), 6.42(*s*, 2H), 7.02–7.35(*m*, 8H).

Tert-butyl 4-*chlorophenylcarbamate* (11): (lit. [14]); IR (KBr): 3304, 2961, 1700, 1593 cm⁻¹; ¹H NMR(90 MHz, CDCl₃), δ 1.49(*s*, 9H), 6.49(*s*, 1H), 7.19–7.35(*m*, 4H).

Tert-butyl 4-*bromophenylcarbamate* (12): (lit. [14]); IR (KBr): 3360, 2960, 1700, 1591 cm⁻¹; ¹H NMR(90 MHz, CDCl₃), δ 1.48(*s*, 9H), 6.63(*s*, 1H), 7.26–7.43(*m*, 4H).

Tert-butyl naphthalen-1-ylcarbamate (16): (lit. [13]); IR (KBr): 3250, 2972, 1690, 1545 cm⁻¹; ¹H NMR(90 MHz, CDCl₃), δ 1.55(*s*, 9H), 6.86(*s*, 1H), 7.35–7.60(*m*, 4H), 7.75–7.96(*m*, 3H).

Tert-butyl 1,2-*phenylenedicarbamate* (17): (lit. [13]); IR (KBr): 3375, 2976, 1685, 1531 cm⁻¹; ¹H NMR(90 MHz, CDCl₃), δ 1.48(s, 18H), 6.71(s, 2H), 7.04–7.49(m, 4H).

Acknowledgment

The authors gratefully acknowledge the support of this work by the Research Affairs Office of Bu-Ali Sina University (Grant number 32-1716; entitled development of chemical methods, reagent and molecules).

References

- Peppel, T., Roth, C., Fumino, K., Paschek, D., Köckerling, M. and Ludwig, R. "The influence of hydrogen bonding on the physical properties of ionic liquids", *Angew. Chem. Int. Ed.*, 29, pp. 6661–6665 (2011).
- [2] Zare, A., Hasaninejad, A., Salimi Beni, A., Moosavi-Zare, A.R., Merajoddin, M., Kamali, E., Akbari-Seddigh, M. and Parsaee, Z. "Ionic liquid 1-butyl-3-methylimidazolium bromide ([Bmim]Br) as a green and neutral reaction media for the catalyst-free synthesis of 1-amidoalkyl-2-naphthols", *Sci. Iran., Trans. C*, 18, pp. 433–438 (2011).
- [3] Zolfigol, M.A., Khazaei, A., Moosavi-Zare, A.R. and Zare, A. "Ionic liquid 3-methyl-1-sulfonic acid imidazolium chloride as a novel and highly efficient catalyst for the very rapid synthesis of bis(indolyl)methanes under solvent-free conditions", Org. Prep. Proced. Int., 42, pp. 95–102 (2010).
- [4] Zolfigol, M.A., Khazaei, A., Moosavi-Zare, A.R. and Zare, A. "3-methyl-1-sulfonic acid imidazolium chloride as a new, efficient and recyclable catalyst and solvent for the preparation of N-sulfonyl imines at room temperature", J. Iran. Chem. Soc., 7, pp. 646–651 (2010).
- [5] Khazaei, A., Zolfigol, M.A., Moosavi-Zare, A.R. and Zare, A. "An efficient method for the nitration of phenols with NaNO₂ in the presence of 3-methyl-1-sulfonic acid imidazolium chloride", *Sci. Iran., Trans. C*, 17, pp. 31–36 (2010).

- [6] Zolfigol, M.A., Khazaei, A., Moosavi-Zare, A.R., Zare, A. and Khakyzadeh, V. "Rapid synthesis of 1-amidoalkyl-2-naphthols over sulfonic acid functionalized imidazolium salts", *Appl. Catal.*, A, 400, pp. 70–81 (2011).
- [7] Khazaei, A., Zolfigol, M.A., Moosavi-Zare, A.R., Zare, A., Ghaemi, E., Khakyzadeh, V., Asgari, Z. and Hasaninejad, A. "Sulfonic acid functionalized imidazolium salts/FeCl₃ as novel and highly efficient catalytic systems for the synthesis of benzimidazoles at room temperature", *Sci. Iran., Trans. C*, 18, pp. 1365–1371 (2012).
- [8] Khaligh, N.G. "Preparation, characterization and use of 3-methyl-1sulfonic acid imidazolium hydrogen sulfate as an eco-benign, efficient and reusable ionic liquid catalyst for the chemoselective trimethylsilyl protection of hydroxyl groups", J. Mol. Catal. A: Chem., 349, pp. 63–70 (2011).
- [9] Greene, T.W. and Wuts, P.G.M., Protective Groups in Organic Synthesis, 3rd Edn., John Wiley & Sons, New York (1999).
- [10] Jia, X., Huang, Q., Li, J., Li, Sh. and Yang, Q. "Environmentally benign N-boc protection under solvent- and catalyst-free conditions", *Synlett*, pp. 0806–0808. 03.207 (2007).
- [11] Khaksar, S., Heydari, A., Tajbakhsh, M. and Vahdat, S.M. "Hydrogen bond catalyzed chemoselective N-tert-butoxycarbonylation of amines", *Tetrahedron Lett.*, 49, pp. 3527–3529 (2008).
- [12] Upadhyaya, D.J., Barge, A., Stefania, R. and Cravotto, G. "Efficient, solventless N-boc protection of amines carried out at room temperature using sulfamic acid as recyclable catalyst", *Tetrahedron Lett.*, 48, pp. 8318–8322 (2007).
- [13] Chehardoli, Gh., Zolfigol, M.A., Khakyzadeh, V., Golbedaghi, R.A., Hall, N. and Blackman, A.G., "Synthesis and crystal structure determination of [H₂ cryptand222](Br₃)₂: a unique tribromide catalyst for the catalytic chemoselective N-boc protection of amines", *J. Chin. Chem. Soc.*, 58, pp. 538-543 (2011).
- [14] Chankeshwara, S.V. and Chakraborti, A.K. "Copper(II) tetrafluoroborate as a novel and highly efficient catalyst for N-tert-butoxycarbonylation of amines under solvent-free conditions at room temperature", *Tetrahedron Lett.*, 47, pp. 1087–1091 (2006).
- [15] Sunitha, S., Kanjilal, S., Reddy, P.S. and Prasad, R.B.N. "An efficient and chemoselective Brønsted acidic ionic liquid-catalyzed N-boc protection of amines", *Tetrahedron Lett.*, 49, pp. 2527–2532 (2008).
- [16] Suryakiran, N., Prabhakar, P., Reddy, T.S., Rajesh, K. and Venkateswarlu, Y. "Facile N-tert-butoxycarbonylation of amines using La(NO₃)₃ · 6H₂O as a mild and efficient catalyst under solvent-free conditions", *Tetrahedron Lett.*, 47, pp. 8039–8042 (2006).
- [17] Wang, G., Li, C., Li, J. and Jia, X. "Catalyst-free water-mediated N-boc deprotection", *Tetrahedron Lett.*, 50, pp. 1438–1440 (2009).

Mohammad Ali Zolfigol was born in 1966 in Iran. He obtained his B.S. degree from Arak University, Iran, his M.S. from Isfahan University of Technology, Iran, and his Ph.D. from Shiraz University, Iran. He became a faculty member of Bu-Ali Sina University in 1997 and Professor in 2005. In 2003, he was selected as distinguished researcher by the Ministry of Science, Research and Technology of Iran. He was also awarded at the Khwarizmi International Festival and at COMSTEC, in 2008. His research interests include: the discovery and development of new synthetic methods by the synthesis and application of new solid-supported reagents, especially silica-based resins.

Vahid Khakyzadeh was born in 1984 in Iran. He received his B.S. degree in Pure Chemistry, in 2008, and his M.S. degree in Organic Chemistry, from Bu-Ali Sina University, Hamedan, Iran, in 2010, where he is currently working on his Ph.D. degree in the same subject.

He was elected as outstanding M.S. and Ph.D. student in Bu-Ali Sina University in 2009 and 2012. His research interests include: applications of ionic liquids, solvent-free conditions in organic synthesis, and synthesis and application of nano magnetite catalysts in organic reactions.

Ahmad Reza Moosavi-Zare was born in Iran, in 1974. He received his B.S. degree in Chemistry from Yazd University, Iran, in 1998, and his M.S. degree in Organic Chemistry from Mohagheghe Ardebili University (Iran) in 2006. He is currently working on his Ph.D. degree in Organic Chemistry, in the Organic Chemistry Department at Bu-Ali Sina University, Hamedan, Iran. He was elected as outstanding Ph.D. student in Bu-Ali Sina University in 2012. His research interests include: applications of ionic liquids, solvent-free conditions and microwave irradiation in organic synthesis, and green chemistry.

Gholamabbas Chehardoli was born in Iran, in 1973. He received his B.S. degree in Chemistry from Bu-Ali Sina University, Iran, in 1998, and M.S. and Ph.D. degrees in Organic Chemistry from this same university in 2000 and 2006, respectively. He is currently teaching Organic Chemistry in the School of Pharmacy at Hamedan University of Medical Sciences, I.R. Iran. His research fields include: methodology in organic chemistry.

1590

Fatemeh Derakhshan-Panah was born in Iran, in 1987. She received her B.S. and M.S. degrees in Chemistry from Bu-Ali Sina University, Iran, in 2009 and 2011, respectively. She was introduced as outstanding M.S. student in Bu-Ali Sina University in 2010. Her research fields include: methodology in organic chemistry.

Abdolkarim Zare was born in Iran, in 1977. He received his B.S. degree in Chemistry from Yazd University, Iran, in 1998, and his M.S. and Ph.D. degrees in Organic Chemistry from Shiraz University, Iran, in 2002, and 2006, respectively. He is currently faculty member at Payame Noor University (PNU). His research fields include: applications of solvent-free conditions, ionic liquids and microwave irradiation in organic synthesis, and study of methodology in organic chemistry.

Omid Khaledian was born in Iran in 1987. He received his B.S. degree in Pure-Chemistry from Tabriz University, Iran, in 2010, and is currently working on his M.S. degree in Organic Chemistry in the Organic Chemistry Department at Bu-Ali Sina University, Hamedan, Iran. His research interests include: applications of ionic liquids, solvent-free conditions, and green chemistry.