

Moment Inequalities and Applications for Negative Dependence Random Variables

Archive of SID

F.Rivaz*, M.Amini** and A.Bozorgnia***

* Department of Statistics, Tarbiat Modares University

** Department of Mathematics, Sistan and Baluchestan University

*** Department of Statistics, Ferdowsi University

Abstract

In this paper, we extend some moment inequalities for partial sums of negative dependence (ND) random variables. Based on these inequalities, some useful inequalities are obtained.

www.SID.ir

Key Words: Negative Dependence, Moment Inequalities.

1.Introduction and preliminaries Then

Archive of SID

Let $\{X_n, n \geq 1\}$ be a sequence of random variables defined on the probability space (Ω, \mathcal{F}, p) . The moment inequalities and their applications for independent random variables have been studied by Petrov (1995). Shao (2000) and Su (1997) studied moment inequalities for NA random variables. In this paper, we extend these inequalities for ND random variables. To prove our main results we need the following definition, lemmas and theorems.

Definition 1. The random variables X_1, \dots, X_n are said to be ND if we have

$$P\left[\bigcap_{j=1}^n (X_j \leq x_j)\right] \leq \prod_{j=1}^n P[X_j \leq x_j], \quad (1)$$

and

$$P\left[\bigcap_{j=1}^n (X_j > x_j)\right] \leq \prod_{j=1}^n P[X_j > x_j], \quad (2)$$

for all $x_1, \dots, x_n \in R$. An infinite sequence $\{X_n, n \geq 1\}$ is said to be ND if every finite subset X_1, \dots, X_n is ND. The conditions (1) and (2) are equivalent for $n = 2$, but these do not agree for $n \geq 3$ (see [2])

Lemma 1. ([1]) Let X_1, \dots, X_n be ND random variables and f_1, \dots, f_n be a sequence of Borel functions which all are monotone increasing (or all are monotone decreasing), then $f_1(X_1), \dots, f_n(X_n)$ are ND random variables.

Lemmas 2. ([1]) Let X_1, \dots, X_n be ND nonnegative random variables.

$$E\left[\prod_{j=1}^n X_j\right] \leq \prod_{j=1}^n E[X_j].$$

Corollary 1. Let X_1, X_2, \dots, X_n be ND random variables, then for every real t we have

$$Ee^{tS_n} \leq \prod_{i=1}^n Ee^{tX_i}.$$

Lemmas 3. ([3]) If X be an arbitrary random variable and $p > 0$. Then

$$E|X|^p = p \int_0^\infty x^{p-1} P[|X| \geq x] dx.$$

Lemmas 4. ([3]) Let $\{X_n, n \geq 1\}$ be a sequence of random variables with $E|X|^p \leq \infty$ for some $p \geq 1$ and $S_n = \sum_{i=1}^n X_i$. Then

$$E|S_n|^p \leq n^{p-1} M_{p,n},$$

where $M_{p,n} = \sum_{j=1}^n E|X_j|^p$.

Theorem 1. ([3]) Let X be a random variable and $\beta_r = E|X|^r$ for any $0 < r < s$. Then

$$\beta_r^{\frac{1}{r}} \leq \gamma^{\frac{1}{r}-\frac{1}{s}} \cdot \beta_s^{\frac{1}{s}},$$

where $\gamma = P[X \neq 0]$.

Lemmas 5. ([3]) If Y is a random variable with d.f. $\frac{1}{n} \sum_{k=1}^n P[X_k \leq x]$. Then for any $r > 0$,

i)

$$E|Y|^r = \frac{1}{n} \sum_{k=1}^n E|X_k|^r$$

ii)

$$P[Y \neq 0] = \frac{1}{n} \sum_{k=1}^n P[X_k \neq 0].$$

Corollary 2. Applying Theorem 1 to Y with $n=2$ and $p \geq 2$, we conclude that

$$B_n^{\frac{1}{2}} \leq \left[\sum_{k=1}^n P[X_k \neq 0] \right]^{\frac{1}{2} - \frac{1}{p}} M_{p,n}^{\frac{1}{p}},$$

where $B_n = \sum_{j=1}^n EX_j^2$.

2. The Main Results

In this section, first we extend an inequality for ND r.v.s that Petrov [3] proved for independent r.v.'s and Su (1997) proved for NA r.v.'s. Then by using this inequality, we obtain some moment inequalities for ND random variables.

Theorem 2. Let X_1, X_2, \dots, X_n be ND random variables with $E(X_i) = 0$, $EX_i^p < \infty$, $i = 1, 2, \dots, n$ and $p \geq 2$. Then for any $t > p/2$ and $x > 0$,

$$P[|S_n| \geq x] \leq \sum_{j=1}^n P[|X_j| \geq \frac{x}{t}] + 2e^t \left[1 + \frac{x^2}{tB_n} \right]^{-t} \quad (3)$$

Proof. Since proof is similar to Lemma 2.3 (Ref.[3]) and Theorem 1 (Ref.[5]), thus we omit details. We define $Y_j = \min(X_j, y)$, $y > 0$ and $T_n = \sum_{j=1}^n Y_j$. It is clear that for every real x that $x > y$,

$$[S_n \geq x] \subseteq [T_n \geq x] \cup [T_n \neq S_n].$$

Hence for every $h > 0$ and by Markov's inequality

$$\begin{aligned} P[S_n \geq x] &\leq P[T_n \geq x] + P[T_n \neq S_n] \\ &\leq e^{-hx} Ee^{hT_n} \\ &\quad + \sum_{j=1}^n P[X_j \geq y]. \quad (4) \end{aligned}$$

Let $F_j(x) = P[X_j \leq x]$ for all real x and $j = 1, 2, \dots, n$. Since $Y_j \leq X_j$ and $EY_j \leq EX_j = 0$. Thus for every $h > 0$,

$$\begin{aligned} Ee^{hY_j} &\leq 1 + \int_0^y (e^{hx} - 1 - hx) dF_j(x) \\ &\quad + (e^{hy} - 1 - hy) P[X_j \geq y]. \quad (5) \end{aligned}$$

Since the function $g(x) = \frac{e^{hx} - 1 - hx}{x^2}$, for every $h > 0$ is non-decreasing, hence by (5) obtain

$$\begin{aligned} Ee^{hY_j} &\leq 1 + \frac{e^{hy} - 1 - hy}{y^2} EX_j^2 \\ &\leq \exp\left[\frac{e^{hy} - 1 - hy}{y^2} EX_j^2\right]. \end{aligned}$$

By Lemma 1, the random variables Y_1, Y_2, \dots, Y_n are ND because the random variables X_1, X_2, \dots, X_n are ND. Therefore,

$$\begin{aligned} e^{-hx} Ee^{hT_n} &\leq e^{-hx} \prod_{j=1}^n Ee^{hY_j} \\ &\leq \exp\left[-hx + \frac{e^{hy} - 1 - hy}{y^2} B_n\right]. \quad (6) \end{aligned}$$

We put here $h = \frac{1}{y} \ln(\frac{xy}{B_n} + 1)$, then

$$\begin{aligned} e^{-hx} Ee^{hT_n} &\leq \exp\left[-\frac{x}{y} \ln\left(\frac{xy}{B_n} + 1\right) \right. \\ &\quad \left. + \frac{x}{y} - \frac{B_n}{y^2} \ln\left(\frac{xy}{B_n} + 1\right) \right] \\ &\leq \exp\left[\frac{x}{y} - \frac{x}{y} \ln\left(\frac{xy}{B_n} + 1\right) \right]. \end{aligned}$$

Note that an analogous estimate holds for $P[-S_n \geq x]$, with replacement of $Z_j = \min(-X_j, y)$ and $T'_n = \sum_{j=1}^n Z_j$ by $Y_j = \min(X_j, y)$ and $T_n = \sum_{j=1}^n Y_j$. Since by Lemma 1 Z_1, Z_2, \dots, Z_n are ND, hence for every $x > 0$ and $h > 0$,

$$P[|S_n| \geq x] \leq P[S_n \geq x]$$

$$\begin{aligned}
& + P[-S_n \geq x] \leq P[T_n \geq x] \\
& + P[T'_n \geq x] + P[T'_n \neq -S_n] \\
& \leq \sum_{j=1}^n P[|X_j| \geq y] + e^{-hx} (Ee^{hT_n} \\
& + Ee^{hT'_n}) \\
& \leq \sum_{j=1}^n P[|X_j| \geq y] \\
& + 2 \exp\left[\frac{x}{y} - \frac{x}{y} \ln\left(\frac{xy}{B_n} + 1\right)\right],
\end{aligned}$$

Let $t = \frac{x}{y} > p/2$, we have

$$\begin{aligned}
P[|S_n| \geq x] & \leq \sum_{j=1}^n P[|X_j| \\
& \geq \frac{x}{t}] + 2e^t \left[1 + \frac{x^2}{tB_n}\right]^{-t}
\end{aligned}$$

Hence complete the proof.

Corollary 3. Under the assumptions of Theorem 2 we have

$$\begin{aligned}
\text{i)} \quad E|S_n|^p & \leq A_p(M_{p,n} + B_n^{p/2}), \quad (7)
\end{aligned}$$

$$\begin{aligned}
\text{ii)} \quad E|S_n|^p & \leq 2A_p n^{p/2-1} M_{p,n}. \quad (8)
\end{aligned}$$

Where $A_p > 0$ depends only on p .

Proof.

i) By Lemma 2 and (3) for every $x > 0$ and $p \geq 2$, we have

$$\begin{aligned}
E|S_n|^p & = p \int_0^\infty x^{p-1} P[|S_n| \geq x] dx \\
& \leq p \sum_{j=1}^n \int_0^\infty x^{p-1} P[|X_j|
\end{aligned}$$

$$\begin{aligned}
& \geq x/t] dx \\
& + 2pe^t \int_0^\infty x^{p-1} \left[1 + \frac{x^2}{tB_n}\right]^{-t} dx \\
& = t^p M_{p,n} \\
& + pe^t t^{p/2} \beta(p/2, t - p/2) B_n^{p/2} \\
& \leq A_p (M_{p,n} + B_n^{p/2}),
\end{aligned}$$

where,

$$\begin{aligned}
A_p & = \max(p^p, p^{1+p/2} e^p \beta(p/2, p/2)) \\
& \text{for } t = p.
\end{aligned}$$

ii) By Lemma 4 and Jensen's inequality for every $p \geq 2$, we have

$$\begin{aligned}
E|S_n|^p & \leq A_p (M_{p,n} + E(\sum_{j=1}^n X_j^2)^{p/2}) \\
& \leq A_p (1 + n^{p/2-1}) M_{p,n} \\
& \leq 2A_p n^{p/2-1} M_{p,n}.
\end{aligned}$$

Corollary 4. Under the assumptions of Theorem 2,

$$\begin{aligned}
E|S_n|^p & \leq A_p [1 \\
& + (\sum_{j=1}^n P[X_j \neq 0])^{p/2-1}] M_{p,n}. \quad (9)
\end{aligned}$$

Proof. By Corollaries 2 and 3 we have

$$\begin{aligned}
E|S_n|^p & \leq A_p (M_{p,n} + B_n^{p/2}) \\
& \leq A_p [1 + (\sum_{j=1}^n P[X_j \neq 0])^{p/2-1}] M_{p,n}
\end{aligned}$$

This complete the proof.

Remark 2. If the sum $\sum_{j=1}^n P[X_j \neq 0]$ grows slower than n , particular, if $\sum_{j=1}^n P[X_j \neq 0] = O(n^\alpha)$, $0 < \alpha < 1$, then (6) gives a better estimate than (5) does.

www.SID.ir

Theorem 3. Let X_1, X_2, \dots, X_n are

ND random variables and $EX_k = 0$ for $k = 1, 2, \dots, n$. Then for every $p > 1$,

Archive of SID

$$E|S_n|^p \leq A_p(M_{p,n} + D_n^p), \quad (10)$$

Where $D_n = \sum_{j=1}^n E|X_j|$ and A_p is a positive constant depending only on p .

Proof Since $g(x) = \frac{e^{hx} - hx - 1}{x}$ for any $h > 0$ is a nondecreasing function; hence, by applying it in proof of Theorem 2 and $EX_j \leq E|X_j|$ for $h = \frac{1}{y} \ln(1 + \frac{y}{D_n}) > 0$, we obtain

$$P[|S_n| \geq x] \leq \sum_{j=1}^n P[|X_j| \geq \frac{x}{t}] + 2e[1 + \frac{x}{tD_n}]^{-t}, \quad (11)$$

where $t = \frac{x}{y} > p > 1$, and $x > 0$. Now by using (11) we have

$$\begin{aligned} E|S_n|^p &\leq t^p M_{p,n} \\ &+ 2pe \int_0^\infty x^{p-1} (1 + \frac{x}{tD_n})^{-t} dx \\ &= t^p M_{p,n} \\ &+ 2pt^p D_n^p e \int_0^\infty u^{p-1} (1 + u)^{-t} du \\ &= t^p M_{p,n} + 2pt^p D_n^p e^t \beta(p, t - p), \end{aligned}$$

by putting $t = 2p$ and $A_p = \max((2p)^p, (2p)^{p+1} e^{2p} \beta(p, p))$, hence complete the proof.

Theorem 4. Let X_1, X_2, \dots, X_n are ND random variables with $EX_k = 0$ for $k = 1, 2, \dots, n$. Then for every $p > 1$,

$$E|S_n|^p \leq A_p [1 + (\sum_{j=1}^n P[X_j \neq 0])^{p-1}] M_{p,n}.$$

Proof By Theorem 1 for $r = 1$ and $s = p > 1$, we have

$$D_n \leq [\sum_{j=1}^n P[X_j \neq 0]]^{1-1/p} M_{p,n}^{1/p},$$

now Theorem 3 for every $p > 1$, complete the proof.

Acknowledgments

The authors are very grateful to the referees for the suggestions which allowed us to improve this paper.

5. References

- [1] Bozorgnia, A., Patterson, R.F and Taylor, R.L. (1996) Limit theorems for dependent random variables: World Congress Nonlinear Analysts' 92. P.1639-1650.
- [2] Ebrahimi, N and Ghosh, M. (1981) Multivariate negative dependence. Comm. Statist., A10 (4), 307-337.
- [3] Petrov, V.V. (1995). Limit theorems of probability theory sequences of independent random variables. Clarendon Press. Oxford.
- [4] Shao, Q.M. (2000). A comparison theorem on moment inequalities between negatively associated and independent random variables. Journal of Theor. Probab., 13, 343-353.
- [5] Su, C., Zhao, L.C. and Wang, Y.B. (1997). Moment inequalities and weak convergence for NA sequences. (ser A) Sci. in China, 40, 172-182.