

DIFFERENTIAL SENSITIVITY OF NH_4^+ AND NO_2^- OXIDATION TO CdCl_2 AND AlCl_3 IN AN AQUEOUS SYSTEM

L. Rowhani and A. Moradshahi*

Department of Biology, Faculty of Sciences, Shiraz University, Shiraz, Islamic Republic of Iran

Abstract

The effects of different concentrations of CdCl_2 and AlCl_3 on NH_4^+ and NO_2^- oxidation in an aqueous system were investigated. AlCl_3 had no effect on NH_4^+ removal and its oxidation to NO_2^- , whereas NO_2^- disappearance was greatly enhanced by AlCl_3 . In contrast, CdCl_2 delayed removal of NH_4^+ and its subsequent oxidation to NO_2^- and also decreased the rate of NO_2^- disappearance. When compared to the controls, NO_3^- production from NH_4^+ oxidation decreased severely by CdCl_2 whereas slight increase in NO_3^- production was observed in the presence of AlCl_3 . The results presented provide evidence for differential sensitivity of NH_4^+ and NO_2^- oxidation to AlCl_3 and CdCl_2 in an aquatic environment.

Keywords: Sensitivity; CdCl_2 ; AlCl_3

Introduction

Nitrification, the biological oxidation of NH_4^+ to NO_3^- is carried out by nitrifying bacteria; *Nitrosomonas* which oxidizes NH_4^+ to NO_2^- and *Nitrobacter* which oxidizes NO_2^- to NO_3^- . Addition of toxic substances to soil and water bodies may exert a significant impact on the performance of nitrifying bacteria and hence on the nitrogen balance in the environment [2,7,14]. Lee *et al.* [7] reported inhibitory effects of Cu^{2+} and Ni^{2+} on *Nitrosomonas* sp. and *Nitrobacter* sp. in continuous flow stirred tank reactors. Sato *et al.* [11-13] found that the growth inhibition of *Nitrosomonas europaea* in batch system was highly correlated to the amine compounds of copper and nickel. Significant reduction of nitrifying activity was detected when simulated acid rain was added to copper-amended soil [2]. Addition of lead salts influenced the nitrification processes in soil

[6]. The nitrate formation decreased with increase in lead salts concentrations. Singha *et al.* [14] studied the effect of Zn^{2+} , Cd^{2+} and Ni^{2+} alone and in combination on the N transformation in a loamy-sand soil. Addition of metals in combination caused a severe reduction in nitrification and ammonification. Comparison of the results of bacteria-plate-counts in the presence of heavy metals showed that all tested heavy metals had more toxic effects on isolated soil microorganisms in agar media than in soil [1].

Removal of ammonia from water bodies is an important factor in the survival of aquatic biota. Uptake of ammonia by phytoplanktons and biological nitrification are among the major mechanisms for the removal of ammonia [3,5]. The presence of AlCl_3 and CdCl_2 in agricultural and industrial waste waters is an ever increasing hazard to various biological activities. To our knowledge, there is no report of impacts of AlCl_3

* E-mail: moradshahi@biology.susc.ac.ir

or CdCl_2 on nitrification in an aquatic system. Experiments on the effects of heavy metals on nitrification are mainly carried out in different soil types. In the present study, we report the effects of the AlCl_3 and CdCl_2 on the fate of NH_4^+ and NO_2^- in an aqueous system.

Materials and Methods

Water samples containing different populations of bacteria including *Nitrosomonas* sp. and *Nitrobacter* sp. were collected in 25-L containers from a pond 25 km west of Shiraz, immediately transferred to the laboratory and kept aerated at room temperature. Either AlCl_3 or CdCl_2 was added to 1-L flasks containing 900 ml of water sample to produce the concentration range of 0, 15, 60, 120 and 300 μM AlCl_3 and 0, 5, 25, 50 and 100 μM CdCl_2 . Then, either ammonium sulfate or sodium nitrite was added to each flask to give a final concentration of 2600 $\mu\text{g/L}$ for $\text{NH}_4^+\text{-N}$ and 1200 $\mu\text{g/L}$ for $\text{NO}_2^-\text{-N}$ at the beginning of experiments. The pH of each flask was adjusted to 7.8 with 0.1 M NaOH and the flasks were placed on orbital shaker (model SGM-300-010U) at 130 rpm and kept at room temperature ($\sim 25^\circ\text{C}$) for the entire length of the experiments. There were three replicates for each experiment.

Samples were taken every other day from each flask and the amount of NH_4^+ , NO_2^- and NO_3^- were determined by the colorimetric procedure described by Stainton *et al.* [15]. Ammonium concentration was measured at 640 nm and NO_2^- concentration was determined at 543 nm using Shimatzo double - beam spectrophotometer (model UV-160 A). Nitrate was analyzed after being reduced to NO_2^- by passing through a cadmium column and measuring NO_2^- as described.

Results

Effects of AlCl_3 and CdCl_2 on NH_4^+ Removal

As shown in Figure 1, the NH_4^+ content of water samples was depleted 11 days after the start of the experiment. The presence of 15, 60, 120 and 300 μM AlCl_3 in the media had no effect on the NH_4^+ depletion. In contrast, NH_4^+ depletion was affected by CdCl_2 (Fig. 2). In the absence of CdCl_2 , NH_4^+ disappeared after 11 days from the start of the experiment. In the presence of 5, 25 and 50 μM CdCl_2 , the removal of NH_4^+ began after a lag period and completed on day 15, 21, and 25, respectively. The lag period was longer with increased CdCl_2 concentrations, but the rate of NH_4^+ removal (slopes of the lines) were nearly identical up to 50 μM

CdCl_2 . At 100 μM CdCl_2 , only 54% decrease in NH_4^+ was observed after 27 days from the start of the experiment. Since the values for three replications in each treatment were close to each other, to avoid crowding, only the means are presented and the bars for standard errors are not included in the figures.

Effects of the AlCl_3 and CdCl_2 on NO_2^- Removal

As shown in Figure 3, AlCl_3 stimulated the removal of NO_2^- . In the absence of AlCl_3 , a 75% decrease in NO_2^- content of water samples occurred after 21 days from the start of the experiment. The rate of NO_2^- removal increased with increasing AlCl_3 concentration. At 300 μM AlCl_3 , complete depletion of NO_2^- occurred on day 17. In contrast, NO_2^- removal decreased with increasing CdCl_2 concentration. In control water samples, NO_2^- depletion was completed after 23 days, where as at 100 μM CdCl_2 , NO_2^- content remained relatively unchanged (Fig. 4).

Effects of AlCl_3 and CdCl_2 on NO_2^- Production from NH_4^+ and Its Subsequent Disappearance

In the presence of different concentrations of AlCl_3 , the production of NO_2^- increased up to 11 to 13 days from the start of experiment (Fig. 5). These results and the trend of NH_4^+ removal shown in Figure 1, suggest that NH_4^+ oxidation is the principal source of observed NO_2^- production. Nitrite produced by NH_4^+ oxidation subsequently disappeared from the media (Fig. 5). The pattern of NO_2^- disappearance as affected by different concentrations of AlCl_3 is similar to the pattern of NO_2^- removal shown in Figure 3. This suggests the endogenously produced NO_2^- and exogenous NO_2^- are probably removed by the same mechanism.

In the presence of different concentrations of CdCl_2 (Fig. 6), the increase in NO_2^- closely followed the decrease in NH_4^+ shown in Figure 2. The production of NO_2^- from NH_4^+ oxidation was delayed by CdCl_2 , but the amounts of NO_2^- produced in the presence of 5, 25 and 50 μM CdCl_2 were essentially the same as in control (Fig. 6). CdCl_2 slowed the rate of NO_2^- disappearance which is similar to the pattern of exogenous NO_2^- removal shown in Figure 4.

Effect of AlCl_3 and CdCl_2 on NO_3^- Production

The amount of nitrate as the end product of nitrification was not significantly affected by AlCl_3 . At 0, 15, 60, 120 and 300 μM AlCl_3 , the amounts of $\text{NO}_3^-\text{-N}$ produced from NH_4^+ oxidation by day 31 were 1975, 2114, 2114, 2118 and 2114 $\mu\text{g/L}$, respectively. In




Figure 1. NH_4^+ removal as affected by different concentrations (μM) of AlCl_3 .




Figure 2. NH_4^+ removal as affected by different concentrations (μM) of CdCl_2 .




Figure 3. Effects of different concentrations (μM) of AlCl_3 on NO_2^- removal.




Figure 4. Effects of different concentrations (μM) of CdCl_2 on NO_2^- removal.




Figure 5. Production of NO_2^- from NH_4^+ and its subsequent disappearance as affected by different concentrations (μM) of AlCl_3 . At the start of experiment, $2600 \mu\text{g/L}$ NH_4^+-N , as ammonium sulfate, was added to each flask.




Figure 6. Production of NO_2^- from NH_4^+ and its subsequent disappearance as affected by different concentrations (μM) of CdCl_2 . At the start of experiment, $2600 \mu\text{g/L}$ NH_4^+-N , as ammonium sulfate, was added to each flask.

contrast, NO_3^- production from NH_4^+ oxidation was severely reduced in the presence of CdCl_2 . By day 57, NO_3^- -N produced at 0, 5, and 25 μM CdCl_2 were 1975, 1805 and 756 $\mu\text{g/L}$, respectively. At 50 and 100 μM CdCl_2 , NO_3^- production was completely inhibited. Although these results are in agreement with those shown in Figures 5 and 6, the amounts of NO_3^- recovered were not stoichiometrically equal to the amount of NH_4^+ added at the beginning of the experiments. Nitrate assimilation by microorganisms present in culture solutions or lost by denitrification may account for the differences.

Discussion

In the present work, biological nitrification might have been responsible for the removal of NH_4^+ and NO_2^- . If so, it is suggested that AlCl_3 possibly enhanced NO_2^- oxidation by increasing the activity of *Nitrobacter* sp. but had no effect on the oxidation of NH_4^+ by *Nitrosomonas* sp. In contrast, CdCl_2 possibly reduced the activities of both groups of bacteria. Differential sensitivity of nitrifying bacteria to light, Cu^{2+} and Ni^{2+} has been reported by several investigators [3,7]. Olson [10] and Ward *et al.* [16] have reported the differential photoinhibition of the two steps of the nitrification process and later Diab and Shilo [3] showed that *Nitrobacter* sp. was more sensitive to light than *Nitrosomonas* sp. which could possibly cause accumulation of nitrite in water bodies especially in fish ponds. Lee *et al.* [7] showed that *Nitrosomonas* sp. was more sensitive to Cu^{2+} and Ni^{2+} than *Nitrobacter* sp. Inhibition of nitrification by cadmium has been shown in several experiments [3,4,14]. Laboratory addition of 10, 100 and 500 mg Cd^{2+}/kg dry soil as CdCl_2 , lowered the ability of soil to nitrify 100 μg added NH_4^+ -N/g dry soil [4]. Nitrification in a loamy-sand soil amended with sewage sludge decreased significantly in the presence of Zn^{2+} , Cd^{2+} and Ni^{2+} [14]. Influence of Al^{3+} on nitrification has also been studied by several investigators [8,9]. In alfalfa and straw-amended soil, aluminum sulfate retarded nitrification [8]. Mandal and Parsons [9] showed that addition of AlCl_3 and 0.1 M HCl to peat, inhibited nitrification but had stimulatory effect on ammonification. They suggested the presence of an active acid adapted strain of nitrifiers in the peat.

Further investigations on the effects of Al^{3+} and heavy metals on the activity and viability of isolated *Nitrosomonas* and *Nitrobacter* are in progress for better understanding of the nitrogen balance in aquatic environment.

Acknowledgements

Thanks are due to the Research Council of Shiraz University for financial support of this research.

References

1. Beck T.H. Investigations into the toxic effect on microflora of the heavy metals common in refinery waste products. *Arch. Agron. Soil Sci.*, **144**: 613-627 (1981).
2. Chao W.L. Effects of acid rain on the nitrifying activity of copper amended soil. *Lett. Appl. Microbiol.*, **21**: 23-24 (1995).
3. Diab S. and Shilo M. Effect of light on the activity and survival of *Nitrosomonas* sp. and *Nitrobacter* sp. isolated from fishponds. *Is. J. Aquaculture*, **40**: 50-56 (1988).
4. Dusek L. The effect of cadmium on the activity of nitrifying populations in two different grassland soils. *Plant Soil*, **177**: 43-53 (1995).
5. Hargreaves J.A. Nitrogen biogeochemistry of aquaculture ponds. *Aquaculture*, **166**: 3-4 (1998).
6. Jacobsen S. and Selmer-Olsen A.R. Influence of heavy metal pollution on nitrogen mineralization in soils. 1. Effects of lead salts. Aas (Norway). Norges Landbrukshoegskole. *Lop* (1985).
7. Lee Y.W., Ong S.K., and Sato C. Effects of heavy metals on nitrifying bacteria. *Wat. Sci.Tech.*, **36**: 69-74 (1997).
8. Maftoun M. and Banihashemi Z. Effects of added sulfur, aluminium sulfate and ferrous sulfate on CO_2 evolution, microbial population and nitrification in alfalfa- and straw-amended soil. *Agrochimica*, **25**: 318-326 (1981).
9. Mandal R. and Parsons J.W. The effects of aluminium chloride and hydrochloric acid on the carbon and nitrogen mineralization in peat. *International J. Trop. Agric.*, **13**: 1-4 (1998).
10. Olson R.J. Differential photoinhibition of marine nitrifying bacteria, A possible mechanism for the formation of the primary nitrite maximum. *J. Mar. Res.*, **39**: 227-238 (1981).
11. Sato C., Leung S.W., and Schnoor J.L. *Chemistry for Environmental Engineering*. 3rd ed. McGraw-Hill (1988).
12. Sato C., Schnoor J.L., and McDonald D.B. Characterization of effects of copper, cadmium and nickel on the growth of *Nitrosomonas europaea*. *Envir. Toxicol. Chem.*, **5**: 403-416 (1986).
13. Sato C., Schnoor J.L., and McDonald D.B. Effects of copper and nickel on the growth of *Nitrosomonas europaea*. *Toxicity Assessment, An int. Quart.*, **1**: 357-376 (1986).
14. Singha D.D., Anoop S., and Gupta A.P. Nitrogen transformation in sewage-sludge amended soil as influenced by addition of zinc, cadmium and nickel. *Indian J. Agric. Sci.*, **68**: 96-100 (1998).
15. Stainton M.P., Capel M.G., and Armstrong F.A.I. The chemical analysis of fresh water. *Env. Can. Spec. Publ.*, **5**: 119 (1974).
16. Ward B.B., Olson R.J., and Perry M.J. Microbial nitrification rates in the primary nitrite maximum of southern California. *Deep Sea Res.*, **29**: 247-255 (1982).