

## Topological Lumpiness and Topological Extreme Amenability

A.H. Riazi\*

Faculty of Mathematics and computer science, Amirkabir University of  
Technology, Tehran, Islamic Republic of Iran

### Abstract

In this paper we give some characterizations of topological extreme amenability. Also we answer a question raised by Ling [5]. In particular we prove that if  $T$  is a Borel subset of a locally compact semigroup  $S$  such that  $M(S)^*$  has a multiplicative topological left invariant mean then  $T$  is topological left lumpy if and only if there is a multiplicative topological left invariant mean  $M$  on  $M(S)^*$  such that  $M(\chi_T)=1$ , where  $\chi_T$  is the characteristic functional of  $T$ . Consequently if  $T$  is a topological left lumpy locally compact Borel subsemigroup of a locally compact semigroup  $S$ , then  $T$  is extremely topological left amenable if and only if  $S$  is.

**Keywords:** Mean; Topological extreme amenability; Left lumpy

### 1. Introduction

Let  $S$  be a locally compact (Hausdorff) semigroup. Let  $C_0(S)$  be the subalgebra of  $CB(S)$  consisting of functions which vanish at infinity. Let  $M(S)^*$  be the Banach space of all bounded regular Borel (signed) measures on  $S$  with total variation norm.

Let  $M_0(S) = \{ \mu \in M(S) : \mu \geq 0 \text{ and } \|\mu\| = 1 \}$  be the set of all probability measures in  $M(S)$ . It is known that  $M(S) = C_0(S)^*$  via the correspondence  $\mu \rightarrow \bar{\mu}$  where  $\bar{\mu}(f) = \int f d\mu$  for any  $f$  in  $C_0(S)$  [4, § 14]. Consider the continuous dual  $M(S)^*$  of  $M(S)$ . Denote by  $1$  the element  $1$  in  $M(S)^*$  such that  $1(\mu) = \mu(S)$  for any  $\mu$  in  $M(S)$ .

Also if  $T$  is a Borel subset of  $S$  we define the Borel characteristic functional  $\chi_T$  of  $T$  in  $M(S)^*$  by

$\chi_T(\mu) = \mu(T)$ ,  $\mu \in M(S)$ . An element  $M$  in  $M(S)^{**}$  is called a mean on  $M(S)$  if  $M(1)=1$  and  $M(F) \geq 0$ , whenever  $F \geq 0$ . An equivalent definition for a mean is that

$$\inf \{ F(\mu) : \mu \in M_0(S) \} \leq M(F) \leq \sup \{ F(\mu) : \mu \in M_0(S) \}$$

for any  $F$  in  $M(S)^*$ . We also note that  $M \in M(S)^{**}$  is a mean if and only if  $\|M\|=M(1)=1$ . Each probability measure  $\mu$  in  $M_0(S)$  is a mean on  $M(S)^{**}$  if we put  $\mu(F)=F(\mu)$ , for any  $F$  in  $M(S)^*$ . An application of Hahn-Banach separation theorem shows that  $M_0(S)$  is weak\* dense in the set of all means on  $M(S)^*$ .

Under pointwise operations and supremum norm  $C_0(S)$  becomes a Banach algebra. Arens product can thus be defined in  $C_0(S)^{**}$ . In particular, we have the

\*E-mail: riazi@aut.ac.ir

following defining formulas for any  $f, g$  in  $C_0(S)$ ,  $m$  in  $C_0(S)^*$  and  $\theta, \varphi$  in  $C_0(S)^{**}$ .

$$(m \odot f)(g) = m(fg)$$

$$(\varphi \odot m)(f) = \varphi(m \odot f)$$

$$(\theta \odot \varphi)(m) = \theta(\varphi \odot m)$$

This product induces a multiplication in  $M(S)^*$  via the identification  $M(S) = C_0(S)^*$ . For  $F, G$  in  $M(S)^*$  we denote the multiplication of  $F$  and  $G$  by  $F \times G$ . In [5] it is shown that  $F \times G$  is defined via the following three steps:

(i) For any  $\mu \in M(S)$  and  $f \in C_0(S)$ ,  $\mu_f \in M(S)$  is defined by

$$\int g d\mu_f = \int gf d\mu \text{ for all } g \in C_0(S)$$

(ii) For any  $\mu \in M(S)$  and  $G \in M(S)^*$ ,  $G \times \mu \in M(S)$  is defined by

$$\int f d(G \times \mu) = G(\mu_f) \text{ for all } f \in C_0(S)$$

(iii) For any  $F, G \in M(S)^*$ ,  $F \times G \in M(S)^*$  is defined by

$$(F \times G)(\mu) = F(G \times \mu) \text{ for all } \mu \in M(S).$$

Then  $M(S)^*$  becomes a commutative Banach algebra with identity [5, theorem 1.2.3].

For each  $\mu$  in  $M(S)$  define an operator  $l_\mu : M(S)^* \rightarrow M(S)^*$  by  $l_\mu F(\nu) = F(\mu * \nu)$ ,  $\nu \in M(S)$ , we denote  $l_\mu F$  by  $\mu \odot F$ . A mean  $M$  on  $M(S)^*$  is called topological left invariant (TLIM) if  $M(\mu \odot F) = M(F)$  for all  $F \in M(S)^*$  and for all  $\mu \in M(S)$ . A topological left invariant mean  $M$  on  $M(S)^*$  is called a multiplicative topological left invariant mean (MTLIM) if

$$M(F \times G) = M(F)M(G) \text{ for all } F, G \in M(S)^*.$$

If there is a MTLIM on  $M(S)^*$  we say that  $S$  is extremely topological left amenable (ETLA). For results concerning ETLA semigroups see [5] and [6].

## 2. Main Results

Note that for elements  $M, N$  in  $M(S)^{**}$  their Arens product is denoted by  $M \odot N$  and is defined by

$$(M \odot N)(F) = M(N_L(F)) \text{ for all } F \text{ in } M(S)^*$$

where  $N_L : M(S)^* \rightarrow M(S)^*$  is defined by  $N_L(\mu) = N(\mu \odot F)$ ,  $\mu \in M(S)$ . See [1] and [2].

First we prove two Lemmas.

**Lemma 2.1.** Suppose  $M$  and  $N$  are functionals in  $M(S)^{**}$ .

(i) If  $M$  and  $N$  are means on  $M(S)^*$  then  $M \odot N$  is also a mean on  $M(S)^*$ .

(ii) For each  $\mu \in M(S)$  and each  $F \in M(S)^*$  we have

$$M_L(\mu \odot F) = \mu \odot M_L(F)$$

(iii) If  $M$  is a topological left invariant mean, then  $M \odot N$  is also topological left invariant.

**Proof.** (i) It is easy to see that for each  $\mu \in M(S)$  and  $1 \in M(S)^*$  we have  $\mu \odot 1 = 1(\mu)$ , hence

$$(M \odot N)(1) = M(N_L(1)) = M(1) = 1$$

Also  $\|M \odot N\| \leq \|M\| \|N\|$ , hence  $M \odot N$  is a mean on  $M(S)^*$ .

(ii) For each  $\nu \in M(S)$

$$\begin{aligned} M_L(\mu \odot F)(\nu) &= M(\nu \odot (\mu \odot F)) \\ &= M((\mu * \nu) \odot F) \\ &= M_L(F)(\mu * \nu) \\ &= (\mu \odot M_L(F))(\nu) \end{aligned}$$

Thus  $M_L(\mu \odot F) = \mu \odot M_L(F)$ .

(iii) Suppose  $M$  is topological left invariant, then for each  $\mu \in M(S)$  and  $F \in M(S)^*$  we have

$$\begin{aligned} (M \odot N)(\mu \odot F) &= M(N_L(\mu \odot F)) \\ &= M(\mu \odot N_L(F)) \\ &= M(N_L(F)) \\ &= (M \odot N)(F) \end{aligned}$$

where we have used (ii) in the second equality. So  $M \odot N$  is topological left invariant, whenever  $M$  is.

**Lemma 2.2.** For each  $s \in S$ ,  $F \in M(S)^*$  and  $M \in M(S)^{**}$  we have

- (i)  $(\varepsilon_s)_L(F) = F \odot \varepsilon_s$
- (ii)  $(M \odot \varepsilon_s)(F) = M(F \odot \varepsilon_s)$
- (iii)  $(\varepsilon_s)_L(F \times G) = (F \times G) \odot \varepsilon_s$

$$= (F \odot \varepsilon_s) \times (G \odot \varepsilon_s)$$

(iv) If  $M$  is multiplicative, then  $M \odot \varepsilon_s$  is so.

**Proof.** (i)

$$\begin{aligned} (\varepsilon_s)_L(F)(\mu) &= \varepsilon_s(\mu \odot F) = (\mu \odot F)(\varepsilon_s) \\ &= F(\mu * \varepsilon_s) = (F \odot \varepsilon_s)(\mu) \end{aligned}$$

hence  $(\varepsilon_s)_L(F) = F \odot \varepsilon_s$ .

$$(ii) (M \odot \varepsilon_s)(F) = M((\varepsilon_s)_L(F)) = M(F \odot \varepsilon_s)$$

where we have used (i) in the second equality.

(iii) the first equality follows from (i) and the second one follows from [5, p.27]

(iv) Suppose  $M \in M(S)^{**}$  is multiplicative. Then:

$$\begin{aligned} (M \odot \varepsilon_s)(F \times G) &= M((\varepsilon_s)_L(F \times G)) \\ &= M((F \odot \varepsilon_s) \times (G \odot \varepsilon_s)) \\ &= M(F \odot \varepsilon_s) M(G \odot \varepsilon_s) \\ &= ((M \odot \varepsilon_s)(F))((M \odot \varepsilon_s)(G)) \end{aligned}$$

where we have used (iii) in the second equality and (ii) in the last equality.

The following theorem is an extension of [5, theorem 3.2.1]. But first we need a definition.

**Definition 2.3.** Let  $S$  be a locally compact semigroup and  $T$  a Borel subset of  $S$ .  $T$  is said to be topological left lumpy in  $S$  if it satisfies the following condition.

(TLL) For each  $\delta > 0$  and  $\mu \in M_o(S)$  with compact support, there exists  $a \in S$  such that  $\mu * \varepsilon_a(T) > 1 - \delta$ .

It is known that (TLL) is equivalent to each of the following conditions:

(TLL)<sub>1</sub> For any  $\delta > 0$  and  $\nu \in M_o(S)$  with compact support, there exists  $\mu \in M_o(S)$  with compact support such that

$$\mu(T) > 1 - \delta \quad \text{and} \quad (\nu * \mu)(T) > 1 - \delta$$

(TLL)<sub>2</sub> For any  $\delta > 0$  and  $\nu \in M_o(S)$  with compact support, there exists  $\mu \in M_o(S)$  with compact support such that

$$\mu(T) > 1 - \delta \quad \text{and} \quad (\nu * \mu)(T) > 1 - \delta$$

See [7, pp. 571-574 and addendum on p.585] for more details. See also [3].

**Theorem 2.4.** Suppose  $T$  is a Borel subset of a locally compact semigroup  $S$ . Suppose  $M(S)^*$  has a MTLIM then the following statements are equivalent:

(i)  $T$  is topological left lumpy.

(ii) There is a MTLIM on  $M(S)^*$  such that  $M(\chi_T) = 1$ .

**Proof.** (i)  $\Rightarrow$  (ii). Let  $F = \{\mu_1, \dots, \mu_k\}$  be a finite subset of  $M_o^c(S)$  (The elements in  $M_o(S)$  with compact support). For each  $\varepsilon > 0$  there is  $s = s_{(F, \varepsilon)} \in S$  such that  $\frac{\mu_1 + \dots + \mu_k}{k} * \varepsilon_s(T) > 1 - \frac{\varepsilon}{2}$  (by TLL), in particular  $\mu_i * \varepsilon(T) > 1 - \varepsilon$ ,  $1 \leq i \leq k$ .

Let  $F$  be the collection of all finite (nonempty) subsets of  $M_o^c(S)$ . Put  $\Delta = F \times (0, \infty)$  and order  $\Delta$  as following:

$$(F_1, \alpha_1) \geq (F_2, \alpha_2) \Leftrightarrow F_2 \subseteq F_1 \quad \text{and} \quad \alpha_1 < \alpha_2$$

By above discussion there is a net  $\{s_\alpha\}$  of elements of  $S$  with  $\gamma = (F, \alpha) \in \Delta$ . Since the set of means on  $M(S)^*$  is weak\* compact the net  $\{\varepsilon_{s_\alpha}\}$  has a subnet  $\{\varepsilon_{s_\beta}\}$  which converges weak\* to a mean  $N$  on  $M(S)^*$  and also for each  $\mu \in M_o^c(S)$  we have

$$\begin{aligned} N(\mu \odot \chi_T) &= \lim_{\beta} (\mu \odot \chi_T)(\varepsilon_{s_\beta}) \\ &= \lim_{\beta} (\mu * \varepsilon_{s_\beta})(T) = 1 \end{aligned} \tag{1}$$

Now suppose  $M$  is MTLIM on  $M(S)^*$ . Since the Arens product is weak\* continuous in the second variable and using Lemma 2.2 (iv) we conclude that  $M \odot N$  is multiplicative. Also since  $M$  and  $N$  are means and  $M$  is topological left invariant, by using Lemma 2.1 we conclude that  $M \odot N$  is a MTLIM on  $M(S)^*$ . Now since  $M_o^c(S)$  is weak\* dense in the set of means on  $M(S)^*$ , by using (1) we obtain  $(M \odot N)(\chi_T) = 1$ .

(ii)  $\Rightarrow$  (i) Suppose  $M$  is a MTLIM on  $M(S)^*$  such that  $M(\chi_T) = 1$ . If  $\{\mu_\alpha\}$  is a net in  $M_o^c(S)$  which converges to  $M$  in weak\* topology, then for each  $\nu \in M_o^c(S)$  we have

$$\omega^* - \lim_{\alpha} (\nu * \mu_\alpha - \mu_\alpha) = \nu \odot M - M = 0$$

Since  $\lim_{\alpha} \mu_{\alpha}(T) = M(\chi_T) = 1$  and for each  $\nu \in M_0^c(S)$

$$(\nu * \mu_{\alpha})(\chi_T) = \chi_T(\nu * \mu_{\alpha}) = (\nu * \mu_{\alpha})(T)$$

We conclude that for each  $\nu \in M_0^c(S)$ ,  $\lim_{\alpha} (\nu * \mu_{\alpha})(T) = 1$ .

So for each  $\nu \in M_0^c(S)$  and each  $\delta > 0$  there is  $\mu = \mu_{\alpha} \in M_0^c(S)$  such that  $(\nu * \mu)(T) > 1 - \delta$ . Therefore by (TLL)<sub>2</sub>,  $T$  is topological left lumpy.

Let  $S$  be a locally compact semigroup and  $T$  a locally compact Borel subsemigroup of  $S$ . We recall some of the constructions in [8] and [9].

Let  $B(S)$  be the  $\sigma$ -algebra of Borel subsets of  $S$ .

(1) Let  $\mu \in M(S)$ , then  $\mu_T$  is the restriction of  $\mu$  to  $B(T)$  and  $\mu_T \in M(T)$ .

(2) Let  $F \in M(T)^*$ , then  $F' \in M(S)^*$  is well-defined by  $F'(\mu) = F(\mu_T)$  for any  $\mu \in M(S)$ .

(3) Let  $M \in M(S)^{**}$ , then  $M_0 \in M(T)^{**}$  is well-defined by  $M_0(F) = M(F')$

For any  $F \in M(T)^*$ .

**Lemma 2.5.** (a)  $F \times \mu_T = (F' \times \mu)_T$  for  $F \in M(T)^*$  and  $\mu \in M(S)$ .

(b)  $(F \times G)' = F' \times G'$  for any  $F, G \in M(T)^*$ .

**Proof.** (a) For any  $A \in B(T)$  we denote  $\xi_A$  for characteristic function of  $A$  in  $T$  and  $\chi_T$  for characteristic function of  $A$  is  $S$ .

$$\begin{aligned} (F \times \mu_T)(A) &= \int \xi_A d(F \times \mu_T) = F((\mu_T)_{\xi_A}) \\ &= F((\mu_{\chi_A})_T) = F'(\mu_{\chi_A}) \\ &= \int \chi_A d(F' \times \mu) \\ &= (F' \times \mu)(A) = (F' \times \mu)_T(A) \end{aligned}$$

(b) For any  $\mu \in M(S)$  by (a) we have

$$\begin{aligned} (F' \times G')(\mu) &= F'(G' \times \mu) = F((G' \times \mu)_T) \\ &= F(G \times \mu_T) \\ &= (F \times G)(\mu_T) = (F \times G)'(\mu) \end{aligned}$$

We now state the main result of this paper which answers a question raised by J.M. Ling, See [5, then P. 51].

**Theorem 2.6.** Let  $T$  be a topological left lumpy locally compact Borel subsemigroup of a locally compact semigroup  $S$ . Then  $T$  is ETLA if and only if  $S$  is ETLA.

**Proof.** Suppose  $T$  is ETLA, then by [5, Theorem 3.2.3]  $S$  is ETLA.

Conversely suppose  $S$  is ETLA, by theorem 2.4 there is a MTLIM on  $M(S)^*$  such that  $M(\chi_T) = 1$ . Then  $M_0(F) = M(F')$  defines a TLIM on  $M(T)^*$ , we show that  $M_0$  is multiplicative

$$\begin{aligned} M_0(F \times G) &= M((F \times G)') = M(F' \times G') \\ &= M(F')M(G') \\ &= M_0(F)M_0(G) \end{aligned}$$

**Corollary 2.7.** Let  $T$  be a left ideal of a locally compact semigroup  $S$ . Then  $M(T)^*$  has a MTLIM if and only if  $M(S)^*$  has a MTLIM.

**Proof.** It suffices to show that every left ideal is topological left lumpy. Let  $t \in T$ . If  $K \subseteq S$  is compact then  $Kt \subseteq ST \subseteq T$ . Consider the Dirac measure  $\varepsilon$  at  $t$ . For any  $\mu \in M_0(S)$  with  $\mu(K) = 1$ , we have  $\mu * \varepsilon_t(T) = \int \chi_T(xt) d\mu(x) = \int_K \chi_T(xt) d\mu(x) = \mu(K) = 1$ , hence  $T$  is topological left lumpy.

## References

1. Arens R.F. The adjoint of a bilinear operator. *Proc. Amer. Math. Soc.*, **2**: 839-848 (1951).
2. Civin P. and Yood B. The second conjugate space of a Banach algebra as an algebra. *Pacific. J. Math.*, **3**: 847-870 (1961).
3. Day M.M. Lumpy subsets in left amenable locally compact semigroups. *Ibid.*, **62**: 87-92 (1976).
4. Hewitt E. and Ross K.A. *Abstract Harmonic Analysis I*. Springer-Verlag, Berlin (1963).
5. Ling J.M. Amenable and extremely amenable locally compact semi groups. *Ph.D. Thesis*, University of Calgary (1990).
6. Riazi A. Extremely amenable locally compact semigroups. *Atti. Sem. Mat. Fis. Univ. Modena XLV*, 441-447 (1997).
7. Wong J.C.S. On topological analogues of left thick subsets in semi groups. *Pacific. J. Math.*, **83**: 571-585 (1979).
8. Wong J.C.S. A characterisation of locally compact amenable subsemigroups. *Canada. Math. Bull.*, **23**: 305-312 (1979).
9. Wong J.C.S. Absolutely continuous measures on locally compact semigroups. *Ibid.*, **18**: 127-131 (1975).