Refolding of Lysozyme Upon Interaction with β-Cyclodextrin

G. Rezaei Behbehani,^{1,*} A. Taherkhani,^{2,3} L. Barzegar, A.A. Saboury,⁴ and A. Divsalar^{4,5}

¹Chemistry Department, Faculty of Science, Islamic Azad University, Takestan Branch, Takestan, Islamic Republic of Iran ²Member of Young Researchers Club, Islamic Azad University, Takestan Branch, Takestan, Islamic Republic of Iran

³Department of Physics, Islamic Azad University, Takestan Branch, Takestan, Islamic Republic of Iran ⁴Institute of Biochemistry and Biophysics, University of Tehran, Tehran, Islamic Republic of Iran ⁵Department of Biological Sciences, Tarbiat Moallem University, Tehran, Islamic Republic of Iran

Received: 9 April 2011 / Revised: 1 June 2011 / Accepted: 12 June 2011

Abstract

Effects of β -cyclodextrin, β CD, on refolding of lysozyme was investigated at pH 12 employing isothermal titration calorimetry (ITC) at 300K in 30mM Tris buffer solution. β CD was employed as an anti-aggregation agent and the heats obtained for lysozyme+ β CD interactions are reported and analyzed in terms of the extended solvation model. It was indicated that there are two sets of identical and non-cooperative sites for β CD. Enthalpic force in the first binding sites is more important than entropic one, indicating that electrostatic interaction plays an important role in the interaction of lysozyme with β CD. The interaction in the second binding sites is stronger and both enthalpy and entropy driven but hydrophobic interaction has more important than electrostatic force. These results suggest that the effects of β CD on lysozyme refolding are attributed to its ability to suppress aggregation of the protein.

Keywords: Lysozyme; Isothermal titration calorimetry; β-Cyclodextrin; Binding parameters

Introduction

Cyclodextrins (CDs) have been reported to suppress aggregate formation during the refolding of a wide range of proteins. Their potency is often ascribed to their affinity for aromatic amino acids, whose surface exposure would otherwise lead to protein association. However, no detailed structural studies are available. CDs, consisting of six, seven, or eight D-glucopyranose units, which are referred to as α -, β -, and γ cyclodextrins, respectively. CDs inhibited the chemically induced aggregation and its inhibition was generally in the order of γ -CDs < α -CDs < β -CDs. Hydrophilic CDs reduced the thermally induced unfolding of lysozyme, suggesting that CDs destabilize native lysozyme or stabilize the unfolded state of lysozyme [1-4]. Electrophoresis data indicate that CDs, which promoted lysozyme refolding, arrested

^{*} Corresponding author, Tel.: +98(281)3780040, Fax: +98(281)3780040, E-mail: grb402003@yahoo.com

aggregation at the stage of smaller soluble aggregates [3]. The presence of both anionic and cationic substituents on the same CD molecule was found to partially restore its renaturation ability. The reports show that Beta-cyclodextrin improved thermal stability and biological activity of lysozyme [4-10]. Previously published results suggest that the effects of CDs on protein refolding are attributed to their ability to suppress aggregation of proteins. CDs may show properties similar to chaotropic agents, which may help explain their anti-aggregation and protein refolding ability. Biological assay exhibited that lysozyme and lyophilized protein/cyclodextrin better maintained protein biological activity compared to lyophilized lysozyme in absence of cyclodextrin [1-14]. CDs have hydrophobic cavities that prevent direct interactions on the hydrophobic surfaces of proteins and in this way suppress protein aggregation [12-14].

The tendency of lysozyme to aggregate is most distinct at pH 12. Thus, exposure to alkaline pH of 12 serves as a convenient approach to initiate the aggregation of lysozyme. In these conditions we can follow the anti-aggregation effect of β CD clearly.

Materials and Methods

Hen egg-white lysozyme was obtained from Sigma. The isothermal titration calorimetric experiments were carried out on a VP-ITC ultra sensitive titration calorimeter. The microcalorimeter consists of a reference cell and a sample cell of 1.8mL in volume, with both cells insulated by an adiabatic shield. All solutions were thoroughly degassed before use by stirring under vacuum. The sample cell was loaded with lysozyme solution (1.26 mM) and the reference cell contained buffer solution. The solution in the cell was stirred at 307 rpm by the syringe (equipped with micro propeller) filled with β CD solution (30mM) to ensure rapid mixing. Injections were started after baseline stability had been achieved. The titration of lysozyme with β CD were carried out on a VP-ITC ultra sensitive titration calorimeter (MicroCal, LLC, Northampton, MA). The measurements were performed at a constant temperature of 27.0±0.02 °C and the temperature was controlled using a Poly-Science water bath. The titration of lysozyme with β CD solution involved 30 injections, the first injection was 5 μ L and the remaining ones were 10 μ L. To correct the thermal effects due to β CD dilution, control experiments were done in which identical aliquots were injected into the buffer solution with the exception of lysozyme. In the ITC experiments, the heat changes associated with processes occurring at a constant temperature are measured.

Results and Discussion

The determined heats for lysozyme+ β CD interaction were shown graphically in Figure 1.

We have shown previously [15-27] that the heats of the macromolecules+ligands interactions in the aqueous solvent systems can be reproduced by the following equation:

$$q = q_{\max} x'_{B} - \delta^{\theta}_{A} (x'_{A}L_{A} + x'_{B}L_{B})$$

-($\delta^{\theta}_{B} - \delta^{\theta}_{A}$)($x'_{A}L_{A} + x'_{B}L_{B}$) x'_{B} (1)

The parameters δ_A^{θ} and δ_B^{θ} reflect to the net effect of β CD on the lysozyme stability in the low and high dextrin concentrations respectively. The positive values for δ_A^{θ} or δ_B^{θ} indicate that β CD stabilizes the lysozyme structure and vice versa. x'_B can be expressed as follows:

$$x'_{\rm B} = \frac{px_{\rm B}}{x_{\rm A} + px_{\rm B}} \tag{2}$$

p > 1 or p < 1 indicate positive or negative cooperativity of macromolecule for binding with ligand respectively; p=1 indicates that the binding is noncooperative. $x'_{\rm B}$ is the fraction of bound β CD and $x'_{\rm A} = 1 - x'_{\rm B}$ is the fraction of unbound β CD. We can express $x_{\rm B}$ as follows:

$$x_{\rm B} = \frac{\left[CD\right]}{\left[CD\right]_{\rm max}} \qquad x_{\rm A} = 1 - x_{\rm B} \tag{3}$$

 $[\beta CD]$ is the concentration of βCD after every injection and [CD] is the maximum consternation of βCD upon saturation of all lysozyme molecule. L_A and L_B can be calculated from heats of dilution of CD in water, q_{dilut} , as follows:

$$L_{\rm A} = q_{\rm dilut} + x_{\rm B} \left(\frac{\partial q_{\rm dilut}}{\partial x_{\rm B}} \right), L_{\rm B} = q_{\rm dilut} - x_{\rm A} \left(\frac{\partial q_{\rm dilut}}{\partial x_{\rm B}} \right)$$
(4)

The heats of lysozyme+ β CD interactions were fitted to Eq. 1 over the entire β CD concentrations. In the fitting procedure, the only adjustable parameter (*p*) was changed until the best agreement between the experimental and calculated data was approached.

There are two distinct sets of binding sites on lysozyme, which are clear in Figure 1. The dissociation equilibrium constant (K_d) and the number of binding sites "g" can be determined by the following equation [15-27]:

$$\frac{\Delta q}{q_{\text{max}}}M_0 = (\frac{\Delta q}{q})L_0 \frac{1}{g} - \frac{K_d}{g}$$
(5)

Where $\Delta q = q_{\text{max}} - q$ and q represents the heat value at a certain β CD (L_0) and lysozyme (M_0) concentrations and q_{max} represents the heat value upon saturation of all lysozyme molecule. Therefore, the plot of $(\frac{\Delta q}{q_{\text{max}}})M_0$ vs. $(\frac{\Delta q}{q})L_0$ should be a linear plot with slope of "1/g" and

the vertical-intercept of $\frac{K_d}{g}$, which "g" and K_d can be obtained (Table 1). If q and q_{max} are calculated per mole of lysozyme, then the standard molar enthalpy of binding for each binding site, ΔH^0 , will be $\Delta H^0 = \frac{q_{max}}{g}$. The change in the standard Gibbs free energy, ΔG^0 , and change in standard entropy of binding, ΔS^0 , could be calculated by using association equilibrium constant, $K_a = 1/K_d$, and ΔH^0 value in equations 6 and 7, respectively.

$$\Delta G^{0} = -RT \ln K_{a} \tag{6}$$

$$\Delta S^{0} = \frac{\Delta H^{0} - \Delta G^{0}}{T}$$
⁽⁷⁾

The binding parameters in Table 1 suggest that the effects of BCD on lysozyme refolding are attributed to its ability to suppress aggregation of lysozyme. BCD reduced the unfolding of lysozyme as evidenced by large values of association equilibrium constants $(K_a=49967.89 \text{ and } 130462 \text{ M}^{-1} \text{ at the first and second set}$ of binding sites respectively), suggesting that β CD stabilize native or unfolded state of lysozyme. The binding process for inhibition of lysozyme aggregation at the first set of binding sites was both enthalpy and entropy driven (Table 1), but electrostatic interaction plays an important role in the binding processes. The interaction in the second set of binding sites is stronger and both enthalpy and entropy driven but hydrophobic interaction is more important than electrostatic force for the inhibition of lysozyme aggregation (Table 1). βCD has a stronger affinity for lysozyme at the second set of binding sites, as evidenced by larger association equilibrium constant. A negative δ_{A}^{θ} value ($\delta_{A}^{\theta} = -0.34$) for the interaction is a characteristic of the electrostatic interactions underlying many non-specific ligandprotein interactions, indicating that BCD destabilizes lysozyme structure. Destabilization of lysozyme by β CD indicates that β CD binds preferentially to the

unfolded lysozyme or to a partially folded intermediate form of lysozyme. Such effects are characteristic of nonspecific interactions, in that the nonspecific ligand binds weakly to many different groups at the protein/water interface. Therefore, the calorimetric results suggest that inhibition of lysozyme aggregation is the result of nonspecific interactions at the first set of binding sites. In the other words, the negative δ_A^{θ} values followed by positive value of δ_B^{θ} indicates that firstly, the non-specific binding of β CD to exposed side-chains on unfolded lysozyme will destabilize the native folded form of lysozyme. Alternatively, interactions with



Figure 1. Comparison between the experimental heats, q, (O), for lysozyme+ β CD interaction and calculated data (lines) via Eq. 1. [β CD] are concentrations of β CD solutions in MA at pH 12.

Table 1. Binding parameters for lysozyme+ β CD interaction at pH 12. *p*=1 indicates that the binding is non-cooperative in two sets of binding sites. Enthalpic force in the first binding sites is more important than entropic one, indicating that electrostatic interaction plays an important role in the interaction of lysozyme with β CD. The interaction in the second binding sites is stronger and both enthalpy and entropy driven but hydrophobic interaction has more important than electrostatic force

Parameters	First binding sites	Second binding sites
р	1	1
$g_{ m i}$	2.00 ± 0.03	5.11±0.08
$K_{\rm a}/{ m M}$	49967.89±21.15	130462.30±13.06
$\Delta H/kJ mol^{-1}site^{-1}$	-2.05±0.05	-0.08 ± 0.01
$\Delta G/kJ \text{ mol}^{-1} \text{ site}^{-1}$	-26.80±0.15	-29.18±0.25
Δ S/kJ mol ⁻¹ site ⁻¹	$0.09{\pm}0.01$	0.10±0.01
$\delta^{ heta}_{ m A}$	-0.36±0.03	
$\delta^{ heta}_{ ext{B}}$		0.09 ± 0.02

groups on oligomeric folded proteins can lead to dissociation of these protein aggregates. Finally, cyclodextrin interaction with unfolded proteins may enhance the solubility of partially denatured lysozyme by masking the exposed hydrophobic residues, thereby assisting the refolding of lysozyme molecule.

Acknowledgements

The financial support of Islamic azad university, Takestan branch is gratefully acknowledged.

References

- Pirzadeh, P. Moosavi-Movahedi, A.A. Hemmateenejad, B. Ahmad, F. and Shamsipur, M. Chemometric studies of lysozyme upon interaction with sodium dodecyl sulfate and β-cyclodextrin. Colloids and Surfaces B: Biointerfaces 52: 31-38 (2006).
- Tavornvipas, S. Hirayama, F. Takeda, S. Arima, H. and Uekama, K. Effects of cyclodextrins on chemically and thermally induced unfolding and aggregation of lysozyme and basic fibroblast growth factor. *J. Pharm Sci.* **95**: 2722-2729 (2006).
- 3. Desai, A. Lee, C. Sharma, L. and Sharma, A. Lysozyme refolding with cyclodextrins: structure-activity relationship. *Biochimie* **88**: 1435-1445 (2006).
- 4. Buswell, A.M. and Middelberg, A.P. Critical analysis of lysozyme refolding kinetics. *Biotechnol Prog.* 18: 470-5 (2002).
- Larreta-Garde, V. Xu, ZF. Lamy, L. Mathlouthi, M. and Thomas, D. Lysozyme kinetics in low water activity media. A possible hydration memory. *Biochem Biophys Res Commun.* 155: 816-822 (1988).
- Altobelli, G. and Subramaniam, S. Kinetics of association of anti-lysozyme monoclonal antibody D44.1 and hen-egg lysozyme. *Biophys* J. 79: 2954–2965 (2000).
- Hibbits, K.A. Gill, D.S. and Willson, R.C. Isothermal titration calorimetric study of the association of hen egg lysozyme and the anti-lysozyme antibody HyHEL-5. *Biochemistry* 33: 3584–3590 (1994).
- Newman, M.A. Mainhart, C.R. Mallett, C.P. Lavoie, T.B. and Smith-Gill, S.J. Patterns of antibody specificity during the BALB/c immune response to hen eggwhite lysozyme. *J. Immunol.* 149: 3260–3272 (1992).
- Schwarz, F.P. Tello, D. Goldbaum, F.A. Mariuzza, R.A. and Poljak, R.J. Thermodynamics of antigen-antibody binding using specific anti-lysozyme antibodies. *Eur. J. Biochem.* 228: 388–394 (1995).
- Tello, D. Goldbaum, F.A. Mariuzza, R.A. Ysern, X. Schwarz, F.P. and Poljak R.J. Three-dimensional structure and thermodynamics of antigen binding by anti-lysozyme antibodies. *Biochem Soc Trans.* 21: 943–946 (1993).
- Xavier, K.A. and Willson, R.C. Association and dissociation kinetics of anti-hen egg lysozyme monoclonal antibodies HyHEL-5 and HyHEL-10. *Biophys J.* 74: 2036–2045 (1998).
- 12. Szejtli, J. Introduction and General Overview of Cyclodextrin Chemistry. *Chem. Rev.* **98:** 1743-1753 (1998).
- Blake, C.C.F. Koenig, D.F. Mair, G.A. North, A.C.T. Phillips, D.C. and Sarma, V.R. Structure of Hen Egg-White Lysozyme: A Three-dimensional Fourier Synthesis

at 2 Å Resolution. Nature 206: 757-761 (1965).

- Kamatari, Y.O. Smith, L.J. Dobson, C.M. and Akasaka, K. Cavity hydration as a gateway to unfolding: An NMR study of hen lysozyme at high pressure and low temperature. *Biophys Chem.* 156: 24-30 (2011).
- Rezaei Behbehani, G. Saboury, A.A. and Fallah Baghery, A. A thermodynamic study on the binding of cobalt ion with myeline basic protein. *Bull. Kor. Chem. Soc.* 29: 736-740 (2008).
- 16. Rezaei Behbehani, G. Saboury, A.A. Mohebbian, M. and Ghammamy, S. Application of a simple calorimetric data analysis on the binding study of cyanide ions by Jack bean urease. *Chin. Chem. Lett.* **21**: 457-460 (2010).
- Rezaei Behbehani, G. Barzegar, L. Saboury, A.A. and Ghammami, S. A thermodynamic investigation on the binding of mercury ion with myelin basic protein at different temperatures. *Chin. Chem. Lett.* 22: 623-625 (2011).
- Rezaei Behbehani, G. Application of a New Method to Reproduce the Inthalpies of Transfer of Nal from Water to Aqueous Methanol Ethanol and iProH Solvent Systems at 298 K. Bull. Korean Chem. Soc. 2: 238-240 (2005).
- Rezaei Behbehani, G. Tetrabuty1Ammonium Bromide and tetrapentylamoni bromide from water to aqueous acetonitrile at 298K. Acta Chim. Slov. 52: 288-291 (2005).
- Rezaei Behbehani, G. and Ghamamy, S. A New Equation to Reproduce The Enthalpies of Transfer of Formamide, N-methyl Formamide and N, N- Dimethylformamidefrom water to aqueous methanol mixturs at 298K. *Thermochim. Acta* 444: 71-74 (2006).
- 21. Rezaei Behbahani, G. Saboury, A.A. and Tazikeh, E. Using the extension coordination model to reproduce the enthalpies of transfer of tetraethylurea from water to aqueous ethanol, propan -1-ol and acetonitrile at 298K. *Acta Chim. Slov.* **53**: 363-366 (2006).
- 22. Rezaei Behbahani, G. Saboury, A. A. and Divsalar, A. Thermodynamic study on the binding of calcium and magnesium ions with myelin basic protein Using the extended solvation theory. *Acta Biochimica et Biophysica Sinica* **40**: 964-969 (2008).
- Rezaei Behbahani, G. Saboury, A. A. and Taleshi, E. Determination of partial unfolding enthalpy for lysozyme upon interaction with dodecyltrimethylammoniumbromide using an extended solvation model. *J. Mol. Recognit.* 21: 132-135 (2008).
- 24. Rezaei Behbahani, G. Saboury, A. A. and Divsalar, A. Using the Extended Solvation theory for Thermodynamic Study on the interaction of Magnesium and Cobalt ions with human growth hormone. *J. Korean Chem. Soc.* **52**: 608-613 (2008).
- 25. Rezaei Behbahani, G. Saboury, A. A. Divsalar, A. Faridbod, F. and Ganjali, M.R. A Thermodynamic Study on the Binding of Human Serum Albumin with Lanthanum Ion. *Chinese Journal of Chemistry* 28: 159-163 (2009).
- 26. Rezaei Behbahani, G. Saboury, A. A. Divsalar, A. Hajiand, R. Rezaeid, Z. and Yahaghi, E. thermodynamic study on the binding of Cobalt and iron ions with bovine carbonic anhydrase II molecule at different temperatures. *J. Solution Chem.* **39**: 1142-1152 (2010).
- Rezaei Behbahani, G. Saboury, A. A. Barzegar, L. Zarean, O. Abedini, J. Payehghdr, M. A thermodynamic study on the interaction of nikel with mylin basic protein by isothermal titration calorimetry. *J. Therm. Anal. Cal.* 101: 379–384 (2010).