Pseudophillipsia (Carniphillipsia) (Trilobite) from the Permian Jamal Formation, Isfahan, Iran

H. Ameri^{*1}, M. Yazdi², and A. Bahrami²

 ¹Department of Ecology, Institute of Science, High Technology and Environmental Science, Graduate University of Advanced Technology, Kerman, Islamic Republic of Iran
 ²Department of Geology, Faculty of Sciences, University of Isfahan, Isfahan, Islamic Republic of Iran

Received: 24 July 2016 / Revised: 27 October 2016 / Accepted: 25 January 2017

Abstract

The aim of this study is to bridge a gap in the existence of Trilobites in the marine Permian successions of Iran by exploring a stratigraphic section of the Permian in Jamal Formation (Dizlu section). Based on comprehensive field studies and laboratory investigations three Trilobite specimens were identified in the study area including *Pseudophillipsia* (*Carniphillipsia*) dizluensis sp. nov., *Pseudophillipsia* (Carniphillipsia) sagittalis and Pseudophillipsia (Carniphillipsia) sp. A new occurrence of trilobites collected from the Permian (Wordian) Jamal Formation at a section at Dizlu, Isfahan Province, Iran, is documented. They belong to a new species of Pseudophillipsia (Carniphillipsia), which is described herein, and can be attributed to the a group of species centered on P.(C). paffenholzi (Weber). It is based on 30 specimens from a single horizon, most of which are enrolled. Co-occurrence of Pseudophillipsia (Carniphillipsia) sp. A. and stenohalin groups such as brachiopods. bryozoans, and crinoids are indicating that this species were preferred shallow and carbonate environments. This is the first time that the existences of these specimens are reported in Dizlu section.

Keywords: Trilobite; Pseudophillipsia (Carniphillipsia); Biostratigraphy; Permian; Iran.

Introduction

Late Paleozoic trilobites have been found in a few locations in Iran. The trilobites reported herein are discovered and collected by Dr. M. Yazdi during the recent last decade from a new and richly fossiliferous location in Central Iran. They occur together with divers' fauna of conodonts, ostracod, gastropods, brachiopods and bryozoans (Upper Permian). The Central Iran Upper Permian succession named as the Jamal Formation, a thick (480m) siliciclastic- carbonate sequence. The trilobites are well preserved in full relief in thin-bedded limestone matrix.

Previews Works

Post-Devonian trilobites in Iran were studied by many authors [3, 1, 2, 4 &5]. Besides, [3] were described two new taxa, *Acropyge lanceolata and Iranaspidion sagittalis*, from Abadeh [Central Iran) which are attributed to Late Guadeloupian. After *Iranaspidion* had been it put in synonymy with *Pseudophillipsia* [2,3, &4]. *I. sagittalis* referred to as

^{*} Corresponding author: Tel: +983426226611; Fax: +983426226617; Email: ameri.paleo@yahoo.com

Pseudophillipsia (Carniphillipsia) sagittalis [3]. Feist et al. (according to [5]) were described one wellpreserved specimen which collected from Chahriseh section (about 55km northeast of Isfahan). In addition, [6] were reported Permian trilobites (Pseudophillipsia sp.) in neighboring sections. The Wordian age (Murghabian) has been proposed for those trilobite bearing beds based on the bryozoan contents [6].Hahn [3] have reported three new species, namely Acropyge weggeni, Acropyge? sp. and Iranaspidion ? sp. From the Yush section (Central Alborz Mountains). Lerosey-Aubril [2] reviewed the Late Paleozoic Iranian and Armenian trilobites and recognized two new taxa including: Persia praecox and Pseudophillipsia (s.l) parvizii. Since that, several works conducted with regard to Permian Iran region; [13, 14, 15, 16, 17, 18, 19, 20, 21, 23 & 24].

In this paper, the trilobite fauna of the Dizlu section as a new trilobite locality are described. The Dizlu section is exposed at the north eastern flank of Lamar Mountain, nearly 60 km northeast of Isfahan [Fig. 1].

Geology Setting

Iran is divided into eight geological provinces. These provinces are (1) Zagros, (2) Sanandaj-Sirjan, (3) Urumieh-Dokhtar, (4) Central Iran, (5) Alborz, (6) Kopeh Dagh, (7) Lut, and (8) Makran accretionary prism. Each province has experienced distinct tectonic and sedimentary histories [7]. The Permian successions of Central Iran are subdivided into several formations based on lithological features. In Sanandaj-Sirjan belt, these strata are reported from three areas: Abadeh, Surmaq and, Shahreza areas.

The Abadeh - Shahreza region is well-known for their classic Permian- Triassic outcrops. Particularly, the Abadeh section is on the northern flank of the Hambast Mountain, about 60km SW of the Abadeh town. Likewise, the Shahreza section is located about 14km NE of the Shahreza town. This region was located in the southern part of the Sanandaj-Sirjan belt, as part of northern margin of the Neo-Tethyan Ocean [8]. In this region, the lithological subdivision of the Permian strata includes the Surmaq, the Abadeh, and the Hambast formations. The Hambast Formation is lithologically characterized by the alternation of shale and micritic grey limestone, followed by dark grey limestone beds of the Abadeh Formation. The succession was subdivided on the basis of conodonts biozons [9], representing the entire Clarkia nleveni and lowermost of the C. Transcaucasia zones belonging to Dzhulfian (Wuchiapingian). The upper part of the Hambast Formation is composed of thin-bedded grayish red nodular limestone with ammonoids, nautiloids, rare

brachiopods, rugose corals, crinoid ossicles and fishremains. The conodonts provided an age ranging from Late Dzhulfian- to Late Dorashamian based on C. transcaucasia to the Merrilina aultima – Stepanovites? mostleri zones [9]. The Shahreza section has been investigated in several studies for its facies and isotope geochemistry [9, 10, & 11]. Permian strata in Central Iran are known in several areas; Tabas (Bagh-e-Vang, Shesh Angosht, and Houze-Dorah sections), Kerman and Isfahan areas (Chahriseh and Bagher Abad sections). The Permian strata in Central Iran are known as the Jamal Formation. The Jamal Formation is Artinskian- Early Wuchiapingian in age based on its fossil contents (such as brachiopods, ammonoids, fusulinids and, conodonts). There are two stratigraphic sections of the Jamal Formation in Tabas region which are Bagh-e-Vang and Shesh Angosht sections. Bagh-e-Vang section is exposed nearly 50km north of Tabas town and the Shesh Angosht section is exposed about 4km east of the Bagh-e-Vang section. In both sections the Jamal Formation unconformeably overlies the Carboniferous sandstones of the Sardar Formation. The upper boundary of the Jamal Formation is conformable with the Lower Triassic Sorkh shale Formation [12].

Dizlu Section

The Dizlu section is located 60km northeast of Isfahan. This area constitutes part of the Central Iran zone and is confined to the east of the Sanandaj-Sirjan belt.

The oldest rock unites exposed in Dizlu region area series of dolomite, limestone, sandstone and sandy limestone dated as Devonian. Several faults have disrupted the rock sequences of the area, resulting in Paleozoic sediments being found adjacent to the Triassic deposits [22]. Permian succession in the Dizlu area is equivalent to the Jamal Formation (in Central Iran), Ruteh and Nesen formations (in Alborz and Kopeh Dagh), and Surmaq and Abadeh formations (in Sanandaj-Sirjan). In this section, the Permian strata (Jamal Formation) disconformably overlie on the Devonian deposits (the Bahram Formation). The upper boundary of the Jamal Formation is disconformable with the red bauxite and laterite of the Lower Triassic Sorkh Shale Formation. The base of the Jamal Formation in Dizlu section begins with grey sandstone and it is covered by grey to brown marly limestone. The middle part of the succession comprises thin bedded grey limestone with a few beds of sandy limestone, and sandstone, with gastropods, echinoids, brachiopods, bryozoan, conodonts, and ostracod. The upper part of the succession is composed of medium to thick bedded dark dolomite [Fig. 1-2].



Figure 1. Geological map of northeast of Isfahan, showing the position of the Dizlu section.

Materials and Methods

The Jamal Formation layers were investigated and their fossils content sampled systematically in the Dizlu section. The normal procedure of sampling was done to collect all possible identifiable fossils on the lithological log. Only the *in situ* fossils were collected from the study strata. The materials collected from this locality are more than 30 trilobite specimens. The collected trilobites are stored in the Department of Geology, University of Isfahan under acronym EUIC.



Figure 2. Lithological details of the Jamal Formation in the Dizlu section, northeast of Isfahan; the position of the sampled fossils is highlighted.

The examined and figured material was processed from marl limestone using a vibro tool. The specimens were coated with ammonium chloride to enhance the contrast before being photographed using ring-light illumination. In most cases, additional highlight from the NW quadrant was used. Photographs were taken using a Nikon D90 digital camera with a 105mm objective lens. A part from the digital sharpening, deepening of contrast, and blackening of the background of the photographs has generally not been retouched.

Results and Discussion

Systematic Paleontology

Terminology, morphological terms and abbreviations used herein follow those defined in part O of the Treatise on Invertebrate Paleontology. Main glabellar lobe designates that part of the glabella in front of the preoccipital, while anterior glabellar furrows refers to glabellar furrows S2to S4. The specimens discussed herein are deposited in the Isfahan University, Iran [EUIC].

Order PROETIDA Fortey& Owens 1975 Family PHILLIPSIIDAE Oehlert 1886 Subfamily DITOMOPYGINAE Hupé 1953 Genus *Pseudophillipsia* Gemmellaro 1892

Subgenus *Pseudophillipsia* (*Carniphillipsia*) Hahn &Brauckmann 1975.

Type Species. Pseudophillipsia ogivalis Gauri 1965, from the Lower Kasimovian (Pennsylvanian) of the Zoellner Ridge near Waidegger-Alm, Carnic Alps, Austria.

Pseudophillipsia (Carniphillipsia) dizluensis sp. nov. Plate 1, [a-j] and Figure 3 [a-d].

Holotypus (paratypes). Complete enrolled specimen, [pl. 1 [a-e], [EUIC 13282].

Etymology: the specific epithet *dizluensis* refers to the sample location that is Dizlu village.

Locus Typicus: Dizlu section, about 60 km northeast of Isfahan, Iran [Fig. 1].

Stratum Typicum: Marly limestones, upper part of the Jamal Formation, Wordian (Guadeloupian) [Fig. 2].

Material, locality and horizon: Complete enrolled holotype specimen [pl. 1 [a-e], [EUIC 13282], Complete enrolled specimen [pl. 1 [a-e], [EUIC 13283], large and well- preserved Complete specimen [pl. 1 [j], [EUIC 13284]; all from a dark-grey sandy wakestone with gastropods, brachiopods, bryozoan (fenestellids), crinoids, ostracods and rear trilobites; Jamal Formation, Wordian.

Diagnosis. one species of *Pseudophillipsia* (*Carniphillipsia*) displayed the following

characteristics: Cephalon of semi-circular out line, anterior border convex and narrow (sag.) in lateral view lateral margin is narrow but increasing backward (posterior), glabella convex and expand forward, glabellar furrows (S2-S4) are narrow and inflated posteriorly. So very convex forwards, So furrows deep and separated by sharp furrow with L1.Occipital ring is narrow and long (tr.); palpebral lobe convex and narrow (tr.) eyes are very convex and eye ring narrow. Preglabellar field is very narrow. Genal spin is short, pygidium with 13+/-1 axial ring and 10 pleural ribs

Description: Cephalon of strongly parabolic outline. Occipital ring high posteriorly and sloping forwards with sharp farrow; glabella moderately convex, higher than occipital ring posteriorly, increasingly sloping forwards. The widest part of glabella is across frontal lobe, this being about7 mm glabellar length, including occipital ring. Although the frontal glabellar profile is rounded across the midline, the anterolateral corner is slightly angulate where the axial furrows end, so that the overall shape of the composite anterior glabellar lobe is somewhat wedge shaped. Pygidium is triangular in outline [pl. 1 [j, g]. Pleural field representing about a third of the width (tr.) of the pygidium anteriorly and composed of 10, possibly 11 ribs; pleural furrows particularly deep, broad; in thorax axis is long, reaching border furrow; 20-22 axial rings which form two rows subdividing the axis into a wide (tr.) median lobe and two narrow lateral lobes; Border narrow anteriorly, slightly widening posteriorly. Axial rings and pleural ribs are particularly prominent.

Comparison: differs from *P. sagittalis* in its more distant cephalic border marked by a border furrow that continues around the anterior of the glabella, its smaller eyes which are distinctly less than half as long as the glabella. *P. dizluensis* differs from *P. sagittalis*, *P. armenia* and P. *paffenholzi* in its lesser number of Pygidial segments, and probably also in its much shorter



Figure 3. *Pseudophillipsia (Carniphillipsia) dizluensis* sp. nov. Reconstruction of (a), Cephalon in frontal view, (b), right lateral view, (c), cephalo-thorax in dorsal view, (d), thoraco-pygidium in dorsal view.

J. Sci. I. R. Iran

genal spine. The spins are broken from all of the available specimens, but the shape of libergena and the posteriorly directed course of the posterior border furrow are similar to those of *P. armenia*.

Table1 summarized the characteristics that displayed by *Pseudophillipsia* (*Carniphillipsia*) dizluensis sp. nov.to distinguish this species from other species of *Pseudophillipsia*.

Table 1. Summarized the characteristics that displayed	d by Pseudophillip	sia (Carniphillipsia) dizluen:	sis sp. nov. to distinguish thi	s species from other species of	f Pseudophillipsia.
--	--------------------	--------------------------------	---------------------------------	---------------------------------	---------------------

Taxa	Anterior	Cranidial	Posterior	Main glabellar	L1	Median	Occipital	Palpebral	Genal spine	Librigenal	Pygidial	Author/year
	margin of	anterior	glabella	lobe		preoccipital	ring	lobes		border	axial	
	cranidium	border				Lobe					rings	
Acropygelenceola		shorter (sag.)	parallel-		lower	lower					more	Kobayashi&
ta			sided				4					Hamada 1984
P.(s.l.) armenica	more	shorter (sag.)	parallel-	anteriorly narrow			convex	wider (tr.)			more	Weber 1944
	forwardly		sided	(tr.)								
	convex											
P. (c.) sagittalis	more	shorter (sag.)	parallel-	extending farther	lower	larger (sag.)	markedly	wider (tr.)			more	Kobayashi&
	forwardly		sided	forward								Hamada 1978
	convex											
P.(s.l.) caucasica			parallel-	narrower (tr.)		shorter(tr.)					loss	Weber 1944
D (1)			sided									
P.(s.l.) parvizii											more	Aubril 2012
P. (c.) paffenholzi	more	shorter (sag.)	parallel-	anterior glabellar	lower	larger (sag.)	longer (sag.)		wider base	mach high	more	Weber 1944
	forwardly		sided	furrows visible			no node			U		
	convex			smooth								
P. (c.) kemerensis	more	shorter (sag.)	parallel-	anterior glabellar	lower	shorter(tr.)	longer (sag.)	wider (tr.)			more	Aubril 2009
	forwardly		sided	furrows visible								
	convex			smooth								
P. (c.)		shorter (sag.)		anteriorly narrow	narrower (tr.)	lower	longer (sag.)					Qian, 1981
reggorcakaensis				(tr.) posteriorly								
				wider(tr.)								
<i>P</i> . (<i>c</i> .) sp. From				more forwardly			longer (sag.)					Stepanov et al.
Iran				convex, narrower			more convex					(1969
				(tr.) posteriorly			backwards					
P.(c.) sp. From				anterior glabellar			convex	more				Hahn
USA				furrows visible	1		forwards so	anteriorly				&Brauckmann
$\mathbf{D}(z)$				smooth		1		located				1975 Weber 1022
P. (C.)						larger (sag.)						weber 1955
$\mathbf{P}(\mathbf{a})$	more	shorter (seg.)		more forwardly		chortor (cog)	longor (sog)					Vin 1079
I. (C.)	forwardly	shorter (sag.)		convex		shorter (sag.)	more convey					1 11 1970
nuisnuiensis	convex			CONVEX			hackwards					
P(c)	more		narrower	narrower (tr.)	lower	wider (tr.)	longer (sag.)		wider base			I u 1974
chongaingensis	forwardly		(tr.)	nanower (u.)	narrower (tr.)	shorter (sag)	ionger (sag.)		wider base			Lu 17/T
enongquigensis	convex		(11.)	•	marrower (u.)	shorter (sug.)						
$P_{-}(c_{-})$ steatonyga						lower	less convex					Goldring 1957
(,							backwards					

330

$P_{-}(c_{-}) pryrifmis shorter (sag.) nor forwardly nor inflated nor inflated nor inflated Qian 1977 Qian 1977 Qian 1977 Qian 1977$													
$P_{-(c) intermedia} = \frac{1}{\text{forwardly}} = \frac{1}{\text{sided}} = \frac{1}{\text{triorly narrow}} + \frac{1}{\text{interiorly narrow}} + \frac{1}{interior$	P. (c.) pyrifrmis		shorter (sag.)		more forwardly	more inflated	more inflated						Qian 1977
$P_{-c, c) intermedia} converting conve$					convex wider								
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$					posteriorly (tr.)								
$ \begin{tabular}{ c c c } & \begin{tabular}{ c c c c } & \begin{tabular}{ c c c c } & \begin{tabular}{ c c c } & \begin{tabular}{ c c c c c } & \begin{tabular}{ c c c c c } & \begin{tabular}{ c c c c c c c } & \begin{tabular}{ c c c c c c c } & \begin{tabular}{ c c c c c c c c c c c } & \begin{tabular}{ c c c c c c c c c c c c c c c c c c c$	P. (c.) intermedia	more		parallel-	anteriorly narrow			longer (sag.)					Kobayashi and
convexfarther forward $P. (c.)$ more $$ parallelfarther forward $$ $$ wider (r.)wider base $$ <td></td> <td>forwardly</td> <td></td> <td>sided</td> <td>(tr.) extending</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>Hamada 1980</td>		forwardly		sided	(tr.) extending								Hamada 1980
$\begin{array}{cccccccccccccccccccccccccccccccccccc$		convex			farther forward								
	P. (c.)	more		parallel-	anteriorly narrow				wider (tr.)	wider base			Hahn & Hahn
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	javornikensis	forwardly		sided	(tr.) extending								1977
$\begin{array}{cccc} P. (c.) & \operatorname{more} & \cdots & \cdots & \operatorname{anteriorly narrow} & \operatorname{lower} & \operatorname{shorter} (\operatorname{sg.}) & \operatorname{lower} & \cdots & $		convex			farther forward								
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	P. (c.)	more			anteriorly narrow	lower longer	shorter (sag.)	longer (sag.)					Haas et al. 1980
$P_{i}(c_{i}) \text{more} \dots \text{parallel} \text{anteriorly narrow} \dots \dots \dots \dots \text{wider (tr.)} \text{wider base} \dots \dots \dots \dots \dots \dots \dots \dots \dots $	cooperationis	forwardly			(tr.) more	(exs.)	lower						
$\begin{array}{c c c c c c c c c c c c c c c c c c c $		convex			forwardly convex								
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	P. (c.)	more		parallel-	anteriorly narrow				wider (tr.)	wider base			Lin 1982
P. (c.) liparoides P. (c.)	mengshanensis	forwardly		sided	(tr.) extending								
P. (c.) liparoides parallel- sided anteriorly narrow (r.) posteriorly wider(tr.) more inflated wider (tr.) wider base mach high Hahn & Hahn, 1977 P. (c.) lipara more forwardly parallel- inflated anteriorly narrow (tr.) more lower shorter (sag.) Goldring 1957 P. (c.) lipara more forwardly sided (tr.) more (tr.) more interiorly narrow lower shorter (sag.) Goldring 1957 P. (c.) rakoveci more forwardly sided (tr.) lacks median shorter (sag.) markedly Gauri 1965 convex parallel- forwardly sided (tr.) lacks median shorter (sag.) wider (tr.) Gauri 1965 convex parallel- sided parallel- sided larger (sag.) wider (tr.) markedly shorter (exs.) markedly shorter (exs.) markedly shorter (exs.) shorter (sag.) wider base less Haas et al. 1980 P. (c.) loricata		convex			farther forward								
sided (tr.) posteriorly wider(tr.) more inflated 1977 P. (c.) lipara more parallel- anteriorly narrow lower shorter (sag.) Goldring 1957 forwardly sided (tr.) more lower no node Goldring 1957 econvex forwardly sided (tr.) lacks median shorter (sag.) shorter (sag.	P. (c.) liparoides			parallel-	anteriorly narrow		wider (tr.)		wider (tr.)	wider base	mach high		Hahn & Hahn,
P. (c.) lipara more parallel- anteriorly narrow of finded lower shorter (sag.) longer (sag.) Goldring 1957 forwardly convex sided (tr.) more (tr.) more lower no node Goldring 1957 P. (c.) rakoveci more parallel- anteriorly narrow (tr.) lacks median markedly Gauri 1965 ornvex posterior sukcus shorter (sag.) shorter (sag.) shorter (exs.) Gauri 1965 envex parallel- anteriorly narrow (tr.) lacks median shorter (sag.) shorter (exs.) Gauri 1965 envex parallel- Gauri 1965 envex parallel- Gauri 1965 envex parallel- markedly markedly markedly markedly <td></td> <td></td> <td></td> <td>sided</td> <td>(tr.) posteriorly</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>1977</td>				sided	(tr.) posteriorly								1977
P. (c.) lipara more parallel- anteriorly narrow lower shorter (sag.) longer (sag.) Goldring 1957forwardly sided (tr.) more lower no node Goldring 1957forwardly convex parallel- anteriorly narrow wider (tr.) markedly Gauri 1965forwardly sided (tr.) lacks median shorter (sag.) shorter (sag.) shorter (exs.)convex parallel posterior sukcus larger (sag.) longer (sag.) wider (tr.) Gauri 1965forwardly sided (tr.) lacks median shorter (sag.) shorter (exs.)convex parallel larger (sag.) longer (sag.) wider (tr.) more Zhang 1982shorter (exs.)shorter (exs.)shorter (exs.)shorter (exs.) shorter (sag.) wider base lless Haas et al. 1980(tr.) more forwardly convex higher posteriorly					wider(tr.) more								
P. (c.) lipara more parallel- anteriorly narrow lower shorter (sag.) longer (sag.) Goldring 1957 forwardly sided (tr.) more lower no node Goldring 1957 P. (c.) rakoveci more parallel- anteriorly narrow wider (tr.) markedly Gauri 1965 forwardly sided (tr.) lacks median shorter (sag.) shorter (sag.) shorter (ess.) Gauri 1965 convex posterior sukcus posterior sukcus parallel- Gauri 1965 econvex posterior sukcus posterior sukcus shorter (sag.) wider (tr.) Gauri 1965 P. (c.) loricata parallel- larger (sag.) lower higher posterior (sag.) markedly markedly Babxially ibabxially ibabxially <					inflated								
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	P. (c.) lipara	more		parallel-	anteriorly narrow	lower	shorter (sag.)	longer (sag.)					Goldring 1957
P. (c.) rakoveci more parallel- anteriorly narrow wider (tr.) markedly Gauri 1965 forwardly sided (tr.) lacks median shorter (sag.) shorter (sag.) shorter (exs.) convex parallel larger (sag.) longer (sag.) wider (tr.) more Zhang 1982 sided larger (sag.) longer (sag.) wider (tr.) more Zhang 1982 shorter (sag.) anteriorly narrow lower lower lower shorter (sag.) wider base less Haas et al. 1980 (r.) loricata shorter (sag.) anteriorly narrow lower lower backwards wider base less Haas et al. 1980 (r.) more forwardly convex higher posteriorly backwards less Haas et al. 1980		forwardly		sided	(tr.) more		lower	no node					
P. (c.) rakoveci more parallel- anteriorly narrow wider (tr.) markedly Gauri 1965 forwardly sided (tr.) lacks median shorter (sag.) shorter (sag.) shorter (exs.) Gauri 1965 P. (c.) schoeningi parallel- larger (sag.) longer (sag.) wider (tr.) more Zhang 1982 sided parallel- larger (sag.) longer (sag.) wider (tr.) more Zhang 1982 P. (c.) loricata shorter (sag.) anteriorly narrow lower lower shorter (sag.) more Zhang 1982 P. (c.) loricata shorter (sag.) wider tots.) abaxially lower shorter (sag.) less convex backwards less convex less convex backwards less convex backwards less convex backwards less convex backwards		convex			forwardly convex								
forwardly sided (tr.) lacks median posterior sukcus posterior sukcus posterior sukcus posterior sukcus posterior sukcus posterior sukcus abaxially P. (c.) schoeningi parallel parallel- sided larger (sag.) longer (sag.) wider (tr.) more Zhang 1982 markedly shorter (exs.) abaxially P. (c.) loricata shorter (sag.) anteriorly narrow lower lower (tr.) more (tr.) more (tr.) more forwardly convex higher posteriorly	P. (c.) rakoveci	more		parallel-	anteriorly narrow		wider (tr.)	markedly					Gauri 1965
convex posterior sukcus abaxially P. (c.) schoeningi parallel- sided larger (sag.) longer (sag.) wider (tr.) more Zhang 1982 P. (c.) loricata shorter (sag.) anteriorly narrow (tr.) more lower lower shorter (sag.) wider base less Haas et al. 1980 Haas et al. 1980 indiger posteriorly backwards wider base less Haas et al. 1980		forwardly		sided	(tr.) lacks median		shorter (sag.)	shorter (exs.)					
P. (c.) schoeningi parallel- larger (sag.) longer (sag.) wider (tr.) more Zhang 1982 sided sided shorter (exs.) abaxially shorter (exs.) abaxially P. (c.) loricata anteriorly narrow lower lower shorter (sag.) less Haas et al. 1980 (tr.) more (tr.) more less convex backwards backwards less Haas et al. 1980		convex			posterior sukcus			abaxially					
sided markedly shorter (exs.) abaxially P. (c.) loricata shorter (sag.) anteriorly narrow lower lower shorter (sag.) wider base less Haas et al. 1980 (tr.) more forwardly convex higher posteriorly	P. (c.) schoeningi			parallel-			larger (sag.)	longer (sag.)	wider (tr.)			more	Zhang 1982
P. (c.) loricata shorter (sag.) anteriorly narrow lower lower shorter (sag.) wider base less Haas et al. 1980 (tr.) more forwardly convex higher posteriorly				sided				markedly					
P. (c.) loricata shorter (sag.) anteriorly narrow lower lower shorter (sag.) wider base less Haas et al. 1980 (tr.) more forwardly convex higher posteriorly								shorter (exs.)					
P. (c.) loricata shorter (sag.) anteriorly narrow lower lower shorter (sag.) wider base less Haas et al. 1980 (tr.) more forwardly convex backwards higher posteriorly								abaxially					
(tr.) more less convex forwardly convex backwards higher posteriorly less convex	P. (c.) loricata		shorter (sag.)		anteriorly narrow	lower	lower	shorter (sag.)		wider base		less	Haas et al. 1980
forwardly convex backwards higher posteriorly					(tr.) more			less convex					
higher posteriorly					forwardly convex			backwards					
					higher posteriorly								

Pseudophillipsia (Carniphillipsia) sagittalis (Kobayashi & Hamada, 1978)

Plate 2, [a-1] and Plate 3, [g, h, j-1]

1978 Iranaspidion sagittalis Kobayashi& Hamada, pp. 157-60, figs 1-3. 1981 Iranaspidion sagittalis[15], p. 223.

1981 Iranaspidion sagittalis Kobayashi& Hamada, p. 56, pl. 1, figs 1-3. 1982 Iranaspidion sagittalis[16]; Owens, p. 29.

1984 Iranaspidion sagittalis[17], pp. 68-9, text-fig. 5k, pl. 14, figs 1-3.

2003 Iranaspidion sagittalis Kobayashi& Hamada;[18], p. 382.

2009 *Pseudophillipsia* (Carniphillipsia) sagittalis (Kobayashi & Hamada); Lerosey- Aubril& Angiolini, p.438.

2012 *Pseudophillipsia* (Carniphillipsia) sagittalis (Kobayashi & Hamada); Lerosey- Aubril, p.1033, fig. 4, k, n.

Material, locality and horizon: Complete enrolled specimen [EUIC 13285-6], pl. 2, [a-1], a large and well- preserved pygidium [EUIC 13291], pl. 3, [h] and a small, partially broken pygidium [EUIC 13291], pl. 3, [g]; both from a dark-grey sandy wakestone with gastropods, brachiopods, bryozoan (fenestellids], crinoids, ostracods and rear trilobites; Jamal



Plate 1. *Pseudophillipsia (Carniphillipsia) dizluensis* sp. nov., (a-e) (EUIC 13282), (a), holotypecephalo-thorax in dorsal view, (b), thoraco-pygidium in dorsal view, (c), thorax (d), Cephalon frontal view, (e), left lateral view. (f-i) *Pseudophillipsia (Carniphillipsia) dizluensis* sp. nov. (EUIC 13283), (f), cephalo-thorax in dorsal view, (g), pygidium in dorsal view, (h), thorax in dorsal view, (i), right lateral view, (j), *Pseudophillipsia (Carniphillipsia) dizluensis* sp. nov. (EUIC 13284), dorsal view. Scale bar =3 mm.

Formation, Wordian.

Diagnosis. Two species of *Pseudophillipsia* (*Carniphillipsia*), displaying the following characteristics: Cephalon of semi-circular out line

glabella short (sag.) and parallel-sided all along; anterior border convex backward cranidium particularly fornicated; frontal lobe inflated, short (sag.), covered with coarse tubercles along postero-lateral margin, and



Plate 2. (a-f), *Pseudophillipsia (Carniphillipsia) sagittalis* (Kobayashi & Hamada), (EUIC 13285), (a), Cephalon frontal view, (b), cephalo-thorax in dorsal view, (c), right lateral view, (d), cephalo-thorax in left lateral view, (e), thorax in dorsal view, (f), thoraco-pygidium in dorsal view. (g-k), *P. (C.) sagittalis*, (EUIC 13286), (g), Cephalon in dorsal view, (h), cephalo-thorax in left lateral view, (i), thorax in dorsal view, (j), thoraco-pygidium in dorsal view, (k), thorax in lateral view. (l), *P. (C.) sagittalis*, (EUIC 13287), pygidium in dorsal view. Scale bar =3 mm.

bearing weak sagittal furrow postero-medially; median occipital lobe trapezoid; Pygidium of roughly semicircular outline, with 11 pleural ribs.

Description. Cephalon of semi-circular out line cranidium elongate, glabella convex, elongate (sag.) and narrow (tr.) the widestpart of glabella is carouse frontal lobe (LA), this being about 9 mm glabellar length. Occipital ring high convex; anterior margin

narrow (sag.); Eye ridge and palpebral lobe narrow (tr.) fixgenal area narrow (tr.) S1 defining trapezoid and narrow (tr.) occipital lobe long (exs.) eye very sharped and long (exs.). Pygidium of roughly semi-circular outline [pl. 2, [g-h]. Axis particularly long, overhanging the border posteriorly and delimited laterally by straight axial furrows evenly converging backwards; 17 discernable axial rings which laterally corresponding to



Plate 3. (a-d), *Pseudophillipsia (Carniphillipsia)* sp. (EUIC 13288), (a), thoraco-pygidium in dorsal view, (b), thorax in dorsal view, (c), Cephalon in dorsal view, (d), thoraco-pygidium in left lateral view. (e), *Pseudophillipsia (Carniphillipsia)* sp. (EUIC 13289), pygidium in dorsal view. (f), *Pseudophillipsia (Carniphillipsia)* sp. (EUIC 13290), pygidium in dorsal view. (g), *Pseudophillipsia (Carniphillipsia) sagittalis*, (EUIC 13291), pygidium in dorsal view. (h), *Pseudophillipsia (Carniphillipsia) sagittalis*, (EUIC 13291), pygidium in dorsal view. (h), *Pseudophillipsia (Carniphillipsia) sagittalis*, (EUIC 13292), pygidium in dorsal view. (i), *Pseudophillipsia (Carniphillipsia)* sp. (EUIC 13293), free cheek, showing genal spine. (j-1), *Pseudophillipsia (Carniphillipsia) sagittalis*, (EUIC 13294-6), free cheek, showing genal spine. Scale bar =3 mm.

the six posterior most pleural ribs, pleural furrows deep but narrow, posterior part of axis strongly flexed downwards.

Pseudophillipsia (Carniphillipsia) sp. Plate 3, [a-d, e, f, i] 2000 *Pseudophillipsia (Carniphillipsia)* sp.; Feist et al. in Mistian et al., p. 99, pl. 8, figs. 2,3. 2009 *Pseudophillipsia (Carniphillipsia)* sp.; Lerosey-Aubril& Angiolini, p. 433, Table 1.

2012 *Pseudophillipsia (Carniphillipsia)* sp.; Lerosey-Aubril, figs. 6i, j, l, n, and o.

Material .locality and horizon. Strongly weathered complete enrolled specimen [EUIC 13288; pl.3 [a-d]].

Diagnosis. Cephalon with distinct border, slightly raised to form short flange; shallow border furrow continuing around anterior of cranidium; genal spines short; pygidium with 17 axial segments.

Description. Carapace essentially smooth, weak striations on cephalic border and underside of pygidial border and by transverse row of low sagittally elongate tubercules on crest of each axial ring of thorax and pygidium. Cephalon represents a parabolic outline. Glabella long; dorsal furrows broad and rather shallow, associated with deep fossulae opposite the mid-distance abaxially; median occipital lobe between and rectangular, wide (tr.) but rather short (exs. and sag.); S1 broad and deep near axial furrow, splitting in to shallow anterior and posterior branches rear wards; occipital ring with strength posterior margin; L1 ovoid, long (exs.), rather narrow (tr.) and inflated; main glabellar lobe somewhat eroded postero-sagittally, and bearing three short and shallow anterior glabellar furrows (S2-S4).

Thorax composed of nine segments. Dorsal furrows gently converging backwards and delimiting a wide (tr.) axial lobe. Axial ring wide (sag. and exs.) and composed of a post annulus of even width all along (sag. and exs.). Only the posteriormost segment can be studied entirely, although most characters seem common to the others. Pleura furrow apparently restricted to vicinity of fulcrum. In lateral view [pl.3 [a-d], axis and pleura of similar heights. In posterior view, axis rather high; pleura strongly flexed downwards, abaxial to fulcrum.

Pygidium apparently of strongly parabolic outline [pl.3 [e-f], subdivided into axial projection that receives the outer portion of the two posteriormost thoracic pleurae; 12 pleural ribs defined by broad and deep pleura furrows that interpleural furrows discernible. Axis delimited by deep axial furrows, subdivided into a median part and two abaxial parts, by two rows of muscle scars, and possibly bearing a faint sagittal sulcus posteriorly; 20 axial rings delimited by rather deep axial furrows that are straight medially but backwardly curved abaxial to muscle scars.

Discussion. Pseudophillipsia (Carniphillipsia) sp. A morphologically closed to Pseudophillipsia (Carniphillipsia) paffenholzi, P. (C.) chongyingensis and P. (C.) sagittalis. In addition to the characters common in the four species, exhibiting a narrow (sag.) anterior border furrow, 12 flat pleural ribs and a pygidial pleural field. Although, the parallel-sided portion of its glabella is longer compared with that of the P. (s.l.) armenica, P. (s.l.) caucasica, P. (C.) javornikensis and P. (C.) lipacoides its median preoccipital lobe is rectangular instead of trapezoid and it has fewer axial rings. Like P. (C.) chongyingensis, it also differs from P. (C.) paffenholzi in having a straight posterior margin of the occipital ring, a medially strongly forwardly curved SO.

Acknowledgement

We thank Fatane Zamani (Isfahan University) and Ebrahim Mohammadi (Graduate University of Advanced Technology, Kerman) for many helpful comments on the manuscript. We gratefully acknowledge the research funding provided for this project (No. 419406) by Institute of Science, High Technology and Environmental Sciences, Graduate University of Advanced Technology, Kerman.

References

- Taraz H., Golshani F., Nakazawa K., Shimizu D., Bando Y., Ishii K. I., Murata M., Okimura Y., Sakagami S. & Nakamura K. The Permian and the Lower Triassic systems in Abadeh region, central Iran. (1981).
- 2. Lerosey-Aubril R. The Late Palaeozoic trilobites of Iran and Armenia and their palaeogeographical significance. *Geological Magazine*, **149**:(06): 1023-1045 (2012)
- 3. Lerosey-Aubril R. & Angiolini L. Permian trilobites from Antalya Province, Turkey, and enrollment in Late Palaeozoic trilobites. *Turkish Journal of Earth Sciences*, **18**:(3): 427-448 (2009).
- Fortey R. A. & Heward A. P. A new, morphologically diverse Permian trilobite fauna from Oman. *Acta Palaeontol Pol.*, 60:(1): 201-216 (2015).
- Mistiaen B., Gholamalian H., Gourvennec R., Plusquellec Y., Bigey F., Brice D., Feist M., Feist R., Pour M. G. & Kebria-Ee M. Preliminary data on the Upper Devonian (Frasnian, Famennian) and Permian fauna and flora from the Chahriseh area (Esfahan Province, central Iran). *Annales-Societe Geologique Du Nord*, 8:(2): 93-102 (2000).
- Ernst A., Senowbari-Daryan B. & Hamedani A. Middle Permian Bryozoa from the Lakaftari area, northeast of Esfahan (central Iran). *Geodiversitas*, 28:(4): 543-590 (2006).
- Ameri H. Peri-Gondwana Late Early–Middle Cambrian trilobites from the Kuhbanan Formation in Dahu section, Kerman Province, Iran. *Arabian Journal of Geosciences*: 1-12 (2013).
- Stampfli G. & Borel G. A plate tectonic model for the Paleozoic and Mesozoic constrained by dynamic plate boundaries and restored synthetic oceanic isochrons. *Earth* and Planetary Science Letters, **196**:(1): 17-33 (2002).
- Kozur H. Correlation of the continental uppermost Permian and Lower Triassic of the Germanic Basin with the marine scale in the light of new data from China and Iran. *Albertiana*, 33: 48-51 (2005).
- Heydari E., Arzani N., Safaei M. & Hassanzadeh J. Ocean's response to a changing climate: Clues from variations in carbonate mineralogy across the Permian–Triassic boundary of the Shareza Section, Iran. *Global Planet. Change*, **105**: 79-90 (2013).
- 11. Kozur H. Biostratigraphy and event stratigraphy in Iran around the Permian–Triassic Boundary (PTB): implications for the causes of the PTB biotic crisis. *Global Planet. Change*, **55**:(1): 155-176 (2007).
- 12. Arefifard S. & Isaacson P. E. Permian Sequence stratigraphy in east-central Iran: Microplate records of Peri-Tethyan and

Peri-Gondwanan events. Stratigraphy, 8:(1): 61-83 (2011).

- Richoz S., Krystyn L., Baud A., Brandner R., Horacek M. & Mohtat-Aghai P. Permian–Triassic boundary interval in the Middle East (Iran and N. Oman): Progressive environmental change from detailed carbonate carbon isotope marine curve and sedimentary evolution. J. Asian Earth Sci. 39:(4): 236-253 (2010).
- 14. Liu X. C., Wang W., Shen S. Z., Gorgij M. N., Ye F. C., Zhang Y. C., Furuyama S., Kano A. & Chen X. Z. Late Guadalupian to Lopingian (Permian) carbon and strontium isotopic chemostratigraphy in the Abadeh section, central Iran. *Gondwana Research.*, 24:(1): 222-232 (2013).
- Shen S. Z. & Mei S. L. Lopingian (Late Permian) high-resolution conodont biostratigraphy in Iran with comparison to South China zonation. *Geological Journal.*, 45:(2-3): 135-161 (2010).
- 16. Ghaderi A., Leda L., Schobben M., Korn D. & Ashouri A. High-resolution stratigraphy of the Changhsingian (Late Permian) successions of NW Iran and the Transcaucasus based on lithological features, conodonts and ammonoids. -Mitteilungen aus dem Museum für Naturkunde in Berlin. *Fossil Record.*, **17**:(1): 41 (2014).
- Leda L., Korn D., Ghaderi A., Hairapetian V., Struck U. & Reimold W. U. Lithostratigraphy and carbonate microfacies across the Permian–Triassic boundary near Julfa (NW Iran) and in the Baghuk Mountains (Central Iran). *Facies.*, 60:(1): 295-325 (2014).
- Leven E. J. & Gorgij M. Fusulinids and stratigraphy of the Carboniferous and Permian in Iran. *Stratigraphy and Geological Correlation.*, 19:(7): 687-776 (2011).

- Zakharov Y. D. & Abnavi N. M. The ammonoid recovery after the end-Permian mass extinction: evidence from the Iran-Transcaucasia area, Siberia, Primorye, and Kazakhstan. *Acta Palaeontol. Pol.*, **58**:(1): 127-147 (2013).
- Davydov V. I. & Arefifard S. Middle Permian (Guadalupian) fusulinid taxonomy and biostratigraphy of the mid-latitude Dalan Basin, Zagros, Iran and their applications in paleoclimate dynamics and paleogeography. *GeoArabia*, *Journal of the Middle East Petroleum Geosciences.*, 18:(2): 17-62 (2013).
- 21. Moghadam H. S., Li X. H., Ling X. X., Stern R. J., Khedr M. Z., Chiaradia M., Ghorbani G., Arai S. & Tamura A. Devonian to Permian evolution of the Paleo-Tethys Ocean: New evidence from U–Pb zircon dating and Sr–Nd–Pb isotopes of the Darrehanjir–Mashhad "ophiolites", NE Iran. *Gondwana Research.*, 28:(2): 781-799 (2015).
- Ameri H. Peri-Gondwana Late Early–Middle Cambrian trilobites from the Kuhbanan Formation in Dahu section, Kerman Province, Iran. *Arabian Journal of Geosciences.*, 8:(3): 1467-1478 (2015).
- Hampe O., Hairapetian V., Dorka M., Witzmann F., Akbari A. M. & Korn D. A first late Permian fish fauna from Baghuk Mountain (Neo-Tethyan shelf, central Iran). *Bull. Geosci.*, 88:(1): 1-20 (2013).
- 24. Nejad M. E., Vachard D., Siabeghodsy A. A. & Abbasi S. Middle-Late Permian (Murgabian-Djulfian) foraminifers of the northern Maku area (western Azerbaijan, Iran). *Palaeontologia Electronica.*, 18:(1): 1-63 (2015).