

Original Article

Alteration of OGG1, MYH and MTH1 genes expression in relapsing-remitting multiple sclerosis patients

Roya Amirinejad¹, Mohammad Ali Sahraian², Bahram Mohammad Soltani¹, Mehrdad Behmanesh^{1*}

1. Department of Genetics, Faculty of Biological Sciences, Tarbiat Modares University, Tehran, Iran 2. MS Research Center, Neuroscience Institute, Tehran University of Medical Sciences, Tehran, Iran

Abstract

Introduction: Previous studies revealed that oxidative stress is elevated in multiple sclerosis (MS). It can harm to biological macromolecules such as DNA. However, the molecular mechanism in protection of genetic information from DNA damages is not clear in MS disease. In this study the expression level of some important genes of *OGG1* and *MYH* involved in base excision repair pathway and, *MTH1* and *ITPA* as main cleaning genes of nucleotide pool from rough nucleotides are examined in MS patients in compared to healthy group.

Methods: Peripheral blood mononuclear cells were isolated from relapsing-remitting-MS patients and healthy subjects. After RNA extraction and cDNA synthesis, the expression levels of target genes were examined by RT-qPCR technique.

Results: The level of the *MTH1* and *MYH* genes expression were decreased, but the level of *OGG1* mRNA was higher in patients in comparison to the control group. Obtained result did not shown any correlation between expression of examined genes and clinical features of patients such as MS severity and disease duration.

Conclusion: These preliminary results provide more supportive evidences for involvement of oxidative damage and variation in expression of DNA repair genes in MS. Significant increase of *OGG1* suggest that the development and progression of pathogenesis in Iranian MS can be related to chronic and direct oxidative damage of genomic DNA not nucleotide pools.

Keywords:

Multiple Sclerosis; DNA repair; Genes expression; Oxidative stress; OGG1

Received: 27 Jan 2017 Accepted: 23 May 2017

*Correspondence to: M. Behmanesh

Tel: +98-2182884451 **Fax:** +98-2182884717

Email: behmanesh@modares.ac.ir

Introduction

Multiple sclerosis (MS) is a chronic neurologic disorder that estimated affects 2.5 million peoples in the worldwide. Different types have been identified in MS including relapsing-remitting (RR), primary progressive and secondary progressive (Chen M et al., 2013) that are categorized based on patterns of progression, as well as the intensity and frequency of generated symptoms. RR-MS is the most common

form of MS disease. Environmental and genetic factors have been found to affect the development of MS. The disease is characterized by inflammation, damaged myelin sheaths in nerve cells, axonal degeneration, loss of oligodendrocyte and oxidative stress (Villoslada, 2016).

It has been suggested that oxidative stress and its by-products, reactive oxygen or nitrogen spices (ROS or RNS) are involved in pathogenesis of many types of human diseases such as cardiovascular, autoimmune and neurodegenerative diseases (Behmanesh et al, 2009, Adamczyk and Adamczyk-Sowa, 2016).

It has been shown that by establishing active demyelination process, ROS-mediated injury by peripheral immune cells and central counterparts promotes progressive tissue damage in MS (Kremer et al. 2004, Adamczyk and Adamczyk-Sowa, 2016). Also, microglia and infiltrated macrophages can generate proinflammatory mediators and oxidizing radicals (Colton and Gilbert, 1993).

Free radicals are normally produced in physiological conditions but their level increased in oxidative stress conditions. In all of eukaryotic cells, adjusted levels of free radicals have a crucial role in regulating various cellular functions such as inflammatory response in which production of ROS and RNS attribute to monocyte interactions with brain endothelium (Tasset et al., 2012).

Uncontrolled production of free radicals may cause oxidation of different types of biomolecules such as free nucleotides and genomic DNA. Oxidative modification of genetic material and accumulation of oxidized bases, cause different types of mutation in the DNA and instability in RNA molecules (Behmanesh et al., 2005). Among biomarkers of oxidative DNA damage, 8-oxo-guanine has the crucial role in DNA alteration. It has been shown that this rough base has different quantity in the genomic DNA of blood and brain patients with neurodegenerative diseases (Sliwinska et al., 2016).

To protect the genetic integrity of the cells from endogenous and exogenous source of free radicals, all of living cells are equipped with DNA repair systems. Nucleotide pool cleaners and base excision repair (BER) pathway are among the major mechanisms which are responsible for cell protection from accumulation of oxidized lesions in genomic DNA or nucleotide pools (Nakabeppu et al., 2014).

Several studies are shown that DNA damage and other oxidative stress biomarkers are increased in peripheral blood samples of MS patients (Grecchi et al., 2012; Ibitoye et al., 2016; Tasset et al., 2012). Satoh et al. (2005) identified an aberrant expression of some DNA repair genes in peripheral blood mononuclear cells (PBMC) of MS patients. The associations of some DNA pathway repair genes polymorphisms with MS have been verified (Briggs et al., 2010; Karahalil et al., 2015), but there is not enough information about the essence of DNA Physiol Pharmacol 21 (2017) 129-136 | 130

damages types which are involved in MS pathogenesis.

To get this point the expression of 8-oxoguanine DNA glycosylase (*OGG1*) and MUTYH glycosylase (*MYH*) involved in BER pathway as well as nudix hydrolase 1 (*MTH1*) and inosine triphosphatase (*ITPA*) as the main nucleotide pool cleaner genes compared in PBMC samples from RR-MS patients and healthy subjects.

We evaluated *OGG1*, *MYH*, *ITPA* and *MTH1* genes expression based on previous information about the possible association of DNA repair system with the pathological features of disease, elevated levels of oxidative stress (Miller et al., 2011; Tasset et al., 2012) and abnormal oxidized base (Haider et al., 2011) in MS patients.

Materials and methods

Patients and PBMC collection

Thirty one eligible patients were recruited for this study that diagnosed according to McDonald criteria (McDonald et al., 2001) and MRI test was performed for all of the patients. Patient's age was 22-52 years and The Expanded Disability Status Scale (EDSS) values of patients were between 0-5.5. They had presented to the MS Research Center of Sina Hospital of Tehran University of Medical Sciences, Tehran, Iran from November 2012 to October 2013 (Naghavi Gargari et al., 2015). All of the patients had no familial relation and no other inflammatory or autoimmune disease. They were RR-MS subtype. Also, 27 healthy samples were collected from volunteers with no history of hospitalization for neurological and autoimmune diseases. All of precipitants were selected after interviewed by an experienced neurologist. Written informed consent was obtained from all participants prior to the blood sampling. The Ethics Committees of Tarbiat Modares University approved this study. Three milliliters of whole blood was taken and collected in the anticoagulant EDTA tubes from each participant. PBMC were isolated by density gradient centrifugation on Ficoll-Paque solution (lympholyte, Cedarlane. Netherlands) according to manufacturer's instructions as described.

RNA extraction and cDNA synthesis

Total RNA of PBMC was extracted by acid

131 | Physiol Pharmacol 21 (2017) 129-136

guanidinium-phenol-chloroform procedure using RNXTM-plus solution (SinaClon co., Iran) according to the manufacturer's instructions. The isolated RNAs were treated with DNAasel (Fermentas, Lithuania) for 20 min at 37°C to eliminate any genomic DNA contamination. Integration, concentration and purity of RNAs were verified by agarose gel electrophoresis and spectrophotometry. Three micrograms of purified RNA was used for cDNA synthesis with random hexamer and oligo (dT)₁₈ primers (MWG, Germany) through M-MuLV reverse transcriptase (Thermo scientific, USA) in total 20 μ l reaction mixture according to manufacturer's instructions.

Real-time PCR analysis

The quantification of OGG1, MTH1, MYH and ITPA mRNAs expression was performed with Real-time PCR and Hot FIREPol EvaGreen qPCR Mix plus ROX (Solis BioDyne, Estonia). Relative levels of genes expression were detected by specific primers designed by Oligo software version 6 (USA). The sequence of used primers were: 5'-GTATATGGGCTGGCCTTGGAAG-3' 5'and CTGTTGGCCCTGATACACACG-3' for MYH, 5'-ACCCTGGCTCAACTGTATCACCAC-3' and 5'-CCGCTCCACCATGCCAGTGATG-3' for OGG1, 5'-GGGCCAGATCGTGTTTGAGTTCGT-3' and 5'-TCGTCGGGCCACATGTCCTTG-3' for MTH1 and 5'-AAGAAGCTGGAGGAGGTCG-3' and 5'-TCCAAGGGCATTGAAGCACA-3' for ITPA mRNAs amplification. The GAPDH primers sequences were 5'-CCATGAGAAGTATGACAAC-3' forward: and reverse: 5'-GAGTCCTTCCACGATACC-3' which was used as the internal control. Real-time PCR was carried out using Applied Biosystems StepOne™ Real-Time PCR Systems (Applied Biosystems/MDS SCIEX, Foster City, CA, USA) in a final reaction volume of 20 µl with 10 ng cDNA, 4 µl of EvaGreen 5X master mix and 200 nM of each forward and reverse primers. The PCR condition was as follows: an initial denaturation at 95°C for 5 min, followed by 40 cycles of denaturation at 95°C for 20 sec, annealing at 60°C for 15sec and extension at 72°C for 30sec. Specificity of PCR products were confirmed by running of qPCR products on 12% poly acrylamide gel electrophoresis and melting curve analysis. All experiments were done at least in duplicate. Each gene expression of the individual samples was normalized with the GAPDH gene expression and relative fold change was calculated by $2^{-\Delta\Delta Ct}$ formula (Livak and Schmittgen, 2001).

Statistical analysis

Data obtained from this study was tested for normal distribution by calculating Kolmogorov-Smirnov test and performing the Shapiro-Wilk normality test. Based on the normal distribution Pearson and spearman rank was carried out to analyze correlations between gene expression and clinical features. The expression level of genes were compared to controls via independent T-test and Mann-Whitney U test using SPSS software (Version 21; SPSS Inc, Chicago, IL) and GraphPad Prism Software (Version 5.0, Inc., San Diego, CA) to compare parametric and nonparametric continuous variables, respectively. Data are presented as mean ± standard deviation (SD) and p<0.05 was defined as the statistical significance.

Results

In this study we analyzed *OGG1, MTH1, MYH* and *ITPA* mRNAs expression in 31 MS patients and 27 healthy samples. The EDSS score of patients was a mean 2.16±0.97. The patients consisted of 25 females and 6 males with a mean age of 30.81±7.30 years old and the mean of disease duration was 4.64±3.87 years. The control group was included 17 females and 10 males with a mean age of 27.88±4.04 years old.

The relative expression of *MTH1*, *ITPA*, *MYH* and *OGG1* were determined using quantitative real-time PCR analysis. As shown in Figure 1A, although the level of *MTH1* transcripts in patients was significantly lower than healthy control (p=0.04, 2.6-fold), the expression of *ITPA* mRNA level was not significant difference between MS patients compared to control subjects (p=0.43, 1.4-fold, Fig. 1B). Moreover, results showed, *MYH* expression was decreased in MS patients compared to healthy control (p=0.025, 3.03-fold, Fig. 1C). Interestingly, the expression of *OGG1* was up-regulated in patients (p=0.001, 4.2-fold, Fig. 1D) compared with control samples.

In order to determine whether the disease severity affected on the DNA repair genes expression, the patients divided into two groups based on EDSS scores: EDSS \leq 2 (n=18) and EDSS > 2 (n=13). We observed that expression of *OGG1* (*p*=0.01, 5.2-fold)

DNA repair genes expression in multiple-sclerosis patients



Fig.1. Comparison of normalized genes expression levels between healthy controls and MS patients. Normalized expression of *MYH* (p=0.025), *MTH1* (p=0.04) and *OGG1* (p=0.001) mRNA levels were statistically different in RR-MS patients in comparison with healthy subjects, while the expression of *ITPA* (B) did not show any difference between two groups. The genes expression levels were normalized relative to the GAPDH expression as an internal control. p-value≤0.05 was considered statically significant. The expression shown as mean Δ Ct ± SED.

and MYH (p=0.03, 3.6-fold) were significantly increased and decreased, respectively in patients with EDSS \leq 2 compared with healthy control, but other studied genes have no statistically different from control samples (MTH1: p=0.06, ITPA: p=0.66). Also, in EDSS > 2 group, there were no significant differences between the mRNA level of interested genes in MS patients versus healthy controls (MYH: p=0.052, ITPA: *p*=0.24, OGG1: p=0.15) but interestingly normalized expression of MTH1 decreased in this group (p=0.04, fold change-2.1). Additionally, genes expression did not show any significant difference between two classified groups (EDSS > 2 and EDSS \leq 2 to each other) (*MYH*1: p=0.49, MTH1: p=0.59, OGG1: p=0.64 and ITPA: p=0.65). Obtained results did not reveal any correlations between the normalized expression of

OGG1, MTH1, MYH and *ITPA* and clinical features such as MS severity, disease duration and patient age (data not shown).

Correlation analysis was conducted that the expression level of *OGG1* was correlated with nucleotide pool enzymes (*ITPA*: p=0.01, r= 0.43, *MTH1*: p=0.01, r=0.45, Fig. 2). In addition, significant positive correlation was observed between *MYH* mRNA level with *ITPA* and *MTH1* (p<0.0001, r=0.7, p<0.0001, r= 0.8, respectively, Fig. 3).

Discussion

MS is a neurodegenerative disorder at which oxidative stress plays a major role in neuronal and axonal deterioration and inflammation in the CNS. Investigation of the molecular mechanism for



Fig.2. Correlation analysis of normalized expression levels of *OGG1* with *ITPA*(A) and *MTH1* (B) as the main nucleotide pool cleaner encoding genes. The result of Pearson correlation test showed a positive correlation between the expression levels of *OGG1* with *ITPA* (A) and *MTH1* (B) genes in PBMCs of MS patients.



Fig.3. Correlation analysis between normalized expression of MYH with *ITPA* and *MTH1* genes. The MYH level correlated significantly with *ITPA* (A) and *MTH1* (B) mRNA level in PBMCs of MS patient.

revealing the MS pathophysiology has been the aim of some researches. The presence of oxidative stress markers, including oxidative DNA damage and 8oxoG were reported in MS patients. Ljubisavljevic et al. (2016) demonstrated that the level of 8-oxoG was significantly higher in RR-MS patients in comparison to controls. Furthermore, Polachini et al. (2016) found a significant increase of damaged DNA in leukocytes of RR-MS patients. Although, Oxidative damage to DNA is not only a feature of MS disease, it can cause a pathological cascade of the damages which suggests the involvement of DNA repair system in the pathogenic mechanisms of this neurodegenerative disease. Concerning the potential role of oxidative stress and DNA repair system in neurodegenerative disease progression, we speculated whether the expression genes involved in DNA damage repairs (*MYTH, MTH1, OGG1* and *ITPA*) are implicated in MS progression. Over 150 genes are involved in the known nuclear DNA and mitochondrial DNA repair pathways (Briggs et al., 2010). The current study focused on expression of four genes derived from two DNA repair pathways: base excision repair and nucleotide pool cleaner. We observed that the expression levels of *MTH1* RNA was down-regulated in RR-MS patients in comparison to healthy control. Moreover, we found expression of *ITPA* mRNA was not significant difference in MS patient. Increased level of oxidative stress and oxidized bases (Miljković and Spasojević, 2013; Ohl et al.,

bases (Miljković and Spasojević, 2013; Ohl et al., 2015; Tasset et al., 2012) can cause misincorporation of modified bases into DNA/RNA during

DNA repair genes expression in multiple-sclerosis patients

replication and transcription that cause changing in the content of genetic information and functional abnormality in proteins level which may result in carcinogenesis or neurodegeneration (Nunomura et al., 2007; Gon et al., 2011). A possible explanation for our result might be that the substrate-level has significant effect on the cell economy (Peterson et al., 2016). So, reduction in expression may be related to absence of enzyme substrate. An alternative explanation for this result is that epigenetic modifications may regulate these gene expression in MS patients or dysregulation in upstream genes may affect studied genes expression. Deficiency of studied genes may lead to accumulation of mutant bases during replication and transcription in DNA and RNA respectively. Analysis of data demonstrated that the expression level of MYH mRNA was downregulated in RR-MS patients in comparison to healthy control.

In contrary, we noted the level of OGG1 in the PBMC of the RR-MS patients was significantly higher than the control group. Additionally, OGG1 mRNA expression was significantly up-regulated in EDSS ≤2 patients group versus healthy individuals. These observations suggest that decreased MYH and MTH1 expression may contribute to the elevated 8-oxodG level (Eshtad et al., 2016) in the genomic DNA not nucleotide pool of peripheral blood samples of MS patients that leads to OGG1 up-regulation for removing the mutant base, specifically in the early and middle stages of pathogenesis. The direct oxidation dame to the genomic DNA can be the other possible explanation for increscent of OGG1 expression. This result stays in agreement with significantly increased in DNA-damage biomarkers of MS PBMCs (Grecchi et al., 2012). Our finding is contrary to Tajouri et al. (2003) studies which have shown DNA repair genes such as OGG1 mRNA expression is down-regulated in MS patients. Moreover, a study involving 13 RR-MS and 5 systemic lupus erythematosus (SLE) patients demonstrated that DNA repair genes were deregulated only in SLE patients (Mandel et al., 2004). Differences in our results with other studies may be due to the disease subtype and sample resource or methods of measurement (Array versus Real-time PCR).

Additionally our result showed positive correlation between OGG1 and MYH with the expression of

MTH1 gene as a nucleotide pool cleaner. These results are in agreement with the study's findings that noted the hMYH and hMTH1 functionally cooperate for effective repair (Eshtad et al., 2016). Despite the positive correlation between OGG1 and MTH1, surprisingly, OGG1 expression was increased and versus MTH1 expression level was reduced in patients in comparison with control group. A possible explanation for this might be that the high OGG1 expression level could be а compensatory mechanism in the low expression of MTH1 for maintaining DNA integrity.

Gene mutations can cause hereditary-forms of the neurodegenerative diseases, although, the majority of the patients with neurodegenerative disorder have sporadic disease (Nunomura et al., 2007; Gon et al., 2011). Previously Studies have confirmed the association between single nucleotide polymorphisms of DNA repair genes and the risk of disease occurrence in the lymphocytes of MS patients (Briggs et al., 2010; Karahalil et al., 2015). Moreover, Selvaraj et al. (2009) observed a complete inhibition of PARP1 (DNA single-stranded break repair) function in EAE mice that led to an earlier onset and developed a more severe EAE in compared with wildtype animals. Based on indirect immunofluorescence technique, Grecchi et al. demonstrated that poly (ADP-ribose) (PAR) value, which occurs under stress conditions, increased in MS PBMCs, but PARP1 expression did not show any significant difference between MS patients and healthy control. Also, they found more positive cells for vH2AX as a DNAdamage biomarker in MS patients in comparison to healthy donors (Grecchi et al., 2012). These findings imply the efficacy of DNA repair to remove the mutation bases and may be an important factor for preventing the development of different types of diseases such as MS. Cell type, disease subtype and sample size can be limitation of our study and may affect experimental results. Evaluation of protein level in combination with mRNA level and oxidative biomarker such as oxidized base measurement in large-scale is suggested for robust assessment in future studies.

Conclusion

In conclusion, the present study indicated that the DNA repair genes might involve in the

pathophysiology of MS disease. This is a basis for future research and further studies are needed to confirm the function of several DNA repair systems in the progression of MS disease.

Acknowledgments

We sincerely acknowledge the patients and the MS research center of Sina Hospital. The Iran National Science Foundation and the Department of Research Affairs of Tarbiat Modares University provide the funding of this work.

Conflict of interest

The authors declare that they have no conflict of interest.

References

- Adamczyk B, Adamczyk-Sowa M. New insights into the role of oxidative stress mechanisms in the pathophysiology and treatment of multiple sclerosis. Oxid Med Cell Longev 2016; 2016: 1973834.
- Ahmadi A, Behmanesh M, Boroumand MA, Tavallaei M. Up-regulation of MSH2, XRCC1 and ATM genes in patients with type 2 diabetes and coronary artery disease. Diabetes Res Clin Pract 2015; 109: 500-506.
- Behmanesh M, Sakumi K, Abolhassani N, Toyokuni S, Oka S, Ohnishi YN, et al. ITPase-deficient mice show growth retardation and die before weaning. Cell Death Differ 2009; 16: 1315-1322.
- Behmanesh M, Sakumi K, Tsuchimoto D, Torisu K, Ohnishi-Honda Y, Rancourt D, et al. Characterization of the structure and expression of mouse ITPA gene and its related sequences in the mouse genome. DNA Res 2005; 12: 39-51.
- Briggs FB, Goldstein BA, McCauley JL, Zuvich RL, De Jager PL, Rioux JD, et al. Variation within DNA repair pathway genes and risk of multiple sclerosis. Am J Epidemiol 2010; 172: 217-224.
- Chen M, Carass A, Oh J, Nair G, Pham DL, Reich DS, et al. Automatic magnetic resonance spinal cord segmentation with topology constraints for variable fields of view. Neuroimage 2013; 83:1051-1062.
- Colton C, Gilbert DL. Microglia, an in vivo source of reactive oxygen species in the brain. Adv Neurol 1993; 59: 321-326.
- Eshtad S, Mavajian Z, Rudd SG, Visnes T, Boström J, Altun M, et al. hMYH and hMTH1 cooperate for survival in mismatch repair defective T-cell acute lymphoblastic leukemia. Oncogenesis 2016; 5: e275.
- Gon S, Napolitano R, Rocha W, Coulon S, Fuchs RP. Increase in dNTP pool size during the DNA damage response plays a key role in spontaneous and inducedmutagenesis in Escherichia coli. Proc Natl Acad Sci U S A 2011; 108: 19311-19316.

- Grecchi S, Mazzini G, Lisa A, Armentero MT, Bergamaschi R, Romani A, et al. Search for cellular stress biomarkers in lymphocytes from patients with multiple sclerosis: a pilot study. PLoS One 2012; 7: e44935.
- Haider L, Fischer MT, Frischer JM, Bauer J, Höftberger R, Botond G, et al. Oxidative damage in multiple sclerosis lesions. Brain 2011; 134: 1914-24.
- Ibitoye R, Kemp K, Rice C, Hares K, Scolding N, Wilkins A. Oxidative stress-related biomarkers in multiple sclerosis: a review. Biomark Med 2016; 10: 889-902.
- Karahalil B, Orhan G, Ak F. The impact of detoxifying and repair gene polymorphisms and the levels of serum ROS in the susceptibility to multiple sclerosis. Clin Neurol Neurosurg 2015; 139: 288-294.
- Kremer TM, Rinne ML, Xu Y, Chen XM, Kelley MR. Protection of pulmonary epithelial cells from oxidative stress by hMYH adenine glycosylase. Respir Res 2004; 5:16.
- Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-delta delta C (T)) method. Methods 2001; 25: 402-408.
- Ljubisavljevic S, Stojanovic I, Basic J, Pavlovic DA. The Validation Study of neurofilament heavy chain and 8hydroxy-2'-deoxyguanosine as plasma biomarkers of clinical/paraclinical activity in first and relapsingremitting demyelination acute attacks. Neurotox Res 2016; 30: 530-538.
- McDonald WI, Compston A, Edan G, Goodkin D, Hartung HP, Lublin FD, et al. Recommended diagnostic criteria for multiple sclerosis: guidelines from the international panel on the diagnosis of multiple sclerosis. Ann Neurol 2001; 50: 121–127.
- Mandel M, Gurevich M, Pauzner R, Kaminski N, Achiron A. Autoimmunity gene expression portrait: specific signature that intersects or differentiates between multiple sclerosis and systemic lupus erythematosus. Clin Exp Immunol 2004; 138: 164-170.
- Miljković D, Spasojević I. Multiple sclerosis: molecular mechanisms and therapeutic opportunities. Antioxid Redox Signal 2013; 19: 2286-2334.
- Miller E, Mrowicka M, Saluk-Juszczak J, Ireneusz M. The level of isoprostanes as a non-invasive marker for in vivo lipid peroxidation in secondary progressive multiple sclerosis. Neurochem Res 2011; 36: 1012-1016.
- Nakabeppu Y. Cellular levels of 8-oxoguanine in either DNA or the nucleotide pool play pivotal roles in carcinogenesis and survival of cancer cells. Int J Mol Sci 2014; 15: 12543-12557.
- Nakauchi A, Wong JH, Mahasirimongkol S, Yanai H, Yuliwulandari R, Mabuchi A, et al. Identification of ITPA on chromosome 20 as a susceptibility gene for youngonset tuberculosis. Hum Genome Var 2016; 3: 15067.
- Naghavi Gargari B, Behmanesh M, Sahraian MA. Effect of vitamin D treatment on interleukin-2 and interleukin-4 genes expression in multiple sclerosis. Physiol Pharmacol 2015; 19: 14-21.
- Nunomura A, Moreira PI, Takeda A, Smith MA, Perry G. Oxidative RNA damage and neurodegeneration. Curr

Med Chem 2007; 14: 2968-2975.

- Ohl K, Tenbrock K, Kipp M. Oxidative stress in multiple sclerosis: central and peripheral mode of action. Exp Neurol 2016; 277: 58-67.
- Papeo G. MutT Homolog 1 (MTH1): the silencing of a target. J Med Chem 2016; 59: 2343-2345.
- Peterson JR, Thor S, Kohler L, Kohler PR, Metcalf WW, Luthey-Schulten Z. Genome-wide gene expression and RNA half-life measurements allow predictions of regulation and metabolic behavior in Methanosarcina acetivorans. BMC Genomics 2016; 17: 924.
- Polachini CR, Spanevello RM, Zanini D, Baldissarelli J, Pereira LB, Schetinger MR, et al. Evaluation of deltaaminolevulinic dehydratase activity, oxidative stress biomarkers, and Vitamin D levels in patients with multiple sclerosis. Neurotox Res 2016; 29: 230-242.
- Prieto A, Díaz D, Barcenilla H, Castrillo C, Monserrat J, Merino AG, et al. Increased spontaneous ex vivo apoptosis and subset alterations in peripheral blood T cells from patients with multiple sclerosis. J Clin Immunol 2006; 26: 101-112.
- Satoh J, Nakanishi M, Koike F, Miyake S, Yamamoto T, Kawai M, et al. Microarray analysis identifies an aberrant expression of apoptosis and DNA damageregulatory genes in multiple sclerosis. Neurobiol Dis 2005; 18: 537-550.

Selvaraj V, Soundarapandian MM, Chechneva O, Williams

AJ, Sidorov MK, Soulika AM, et al. PARP-1 deficiency increases the severity of disease in a mouse model of multiple sclerosis. J Biol Chem 2009; 284: 26070-26084.

- Sliwinska A, Kwiatkowski D, Czarny P, Toma M, Wigner P, Drzewoski J, et al. The levels of 7, 8dihydrodeoxyguanosine (8-oxoG) and 8-oxoguanine DNA glycosylase 1 (OGG1)–A potential diagnostic biomarkers of Alzheimer's disease. J Neurol Sci 2016; 368: 155-159.
- Tajouri L, Mellick AS, Ashton KJ, Tannenberg AE, Nagra RM, Tourtellotte WW, et al. Quantitative and qualitative changes in gene expression patterns characterize the activity of plaques in multiple sclerosis. Brain Res Mol Brain Res 2003; 119: 170-183.
- Tasset I, Agüera E, Sánchez-López F, Feijóo M, Giraldo AI, Cruz AH, et al. Peripheral oxidative stress in relapsing– remitting multiple sclerosis. Clin Biochem 2012; 45: 440-444.
- Villoslada P. Neuroprotective therapies for multiple sclerosis and other demyelinating diseases. Multiple Sclerosis and Demyelinating Disorders 2016; 1: 1.
- Zhang Q, Fujino M, Xu J, Li XK. The role and potential therapeutic application of myeloid-derived suppressor cells in allo-and autoimmunity. Mediators Inflamm 2015; 2015: 421927.