

Attenuation of Cyclosporine-Induced Sperm Impairment and Embryotoxicity by *Crataegus monogyna* Fruit Aqueous Extract

Zahra Armand, M.Sc.¹, Gholamreza Najafi, Ph.D.², Farah Farokhi, Ph.D.¹, Ali Shalizar Jalali, Ph.D.²

1. Department of Biology, Faculty of Science, Urmia University, Urmia, Iran

2. Department of Basic Sciences, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

* Corresponding Address: P.O.Box: 571531177, Department of Basic Sciences, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Email: ali_shalizar@yahoo.com

Received: 6/Oct/2012, Accepted: 15/Jan/2013

Abstract

Objective: Cyclosporine (Cs), a cyclic undecapeptide with potent immunosuppressive activity, causes several adverse effects including reproductive toxicity. This study aims to examine the ability of *Crataegus monogyna* aqueous fruit extract as an antioxidant to protect against Cs-induced reproductive toxicity.

Materials and Methods: In this experimental study, 32 adult male Wistar rats were divided into four groups of eight animals each. Rats in two groups received 40 mg/kg/day Cs for 45 days by oral gavage. In addition, one of the two groups received *Crataegus monogyna* aqueous extract at a dose of 20 mg/kg/day orally four hours after Cs administration. The remaining two groups consisted of a vehicle treated control (Cont) group and a *Crataegus monogyna* control (Cr) group. Differences between groups were assessed by analysis of variance (ANOVA) using the SPSS software package for Windows.

Results: Cs treatment caused a significant decrease in sperm count and viability with an increase in DNA damage and protamine deficiency of the sperm cells. We observed significant decreases in fertilization rate and embryonic development, in addition to an increased rate of embryo arrest in Cs-treated rats. *Crataegus monogyna* co-administration attenuated all Cs-induced negative changes in the above-mentioned parameters.

Conclusion: Supplementation with *Crataegus monogyna* aqueous fruit extract could be useful against reproductive toxicity during Cs treatment in a rat model.

Keywords: Cyclosporine, Sperm, *In Vitro* Fertilization, *Crataegus monogyna*

Cell Journal(Yakhteh), Vol 15, No 3, Autumn 2013, Pages: 198-205

Citation: Armand Z, Najafi Gh, Farokhi F, Shalizar Jalali A. Attenuation of cyclosporine-induced sperm impairment and embryotoxicity by *crataegus monogyna* fruit aqueous extract. Cell J. 2013; 15(3): 198-205.

Introduction

Cyclosporine (Cs), a neutral lipophilic cyclic undecapeptide (C₆H₁₁N₁₁O₁₂), is an isolate of the *Topocladium inflatum gams* extract and a member of the Fungi imperfecti family (1). Since the early 1980s, Cs has been successfully used to prevent organ transplant rejection. It has significantly increased graft survival following renal, cardiac, pancreatic, bone marrow and hepatic transplantation and decreased patient mortality (2). In addition, Cs is used as a treatment for autoimmune diseases such as idiopathic nephritic syndrome (3),

uveitis (4), psoriasis (5) and rheumatoid arthritis (6). The immunosuppressant effects of Cs are associated with its specific ability to inhibit signal transductions through the T-cell receptor. This inhibition results in the prevention of cytokine production that would normally stimulate an immune response (7). Despite its therapeutic importance, it has been reported that Cs causes renal, hepatic and cardiac damage (8) as well as reproductive toxicity in humans and experimental animals (9-13). Although the precise mechanism by which Cs causes reproductive toxicity is still unclear, numerous

studies have shown that overproduction of reactive oxygen species (ROS) and lipid peroxidation are causative factors involved in Cs-induced adverse effects (14-16). Additionally, Cs induces direct damage to the hypothalamic-pituitary-gonadal axis (17-19) and Sertoli cell phagocytic function (20) which leads to reproductive toxicity. Hence, a combination of the drug delivery together with potent and safe antioxidant may be the appropriate approach to ameliorate Cs-induced reproductive toxicity.

Hawthorn (*Crataegus* spp.), a member of the Rosaceae family, is native to the Mediterranean region, North Africa, Europe and Central Asia and comprises more than 200 species worldwide (21). Hawthorn has been initially indicated as a drug in "Tang-Ben-Cao" (659A.D.), the world's oldest officially published pharmacopoeia. *Crataegus laevigata* (syn. *Crataegus oxyacantha*) and *Crataegus monogyna* are the two most common species used for medicinal purposes (22). Independent studies have shown that hawthorn fruit constituents are among the best antioxidants (23-26). Based on this concept, the present study seeks to determine if the aqueous extract of *Crataegus monogyna* fruit with antioxidant properties has a possible protective effect against reproductive toxicity during Cs treatment in a rat model.

Materials and Methods

Plant material

In this experimental study, we collected ripe *Crataegus monogyna* fruits from its natural habitat near the city of Urmia in West Azerbaijan Province, Northwestern Iran. The Research Laboratory of the Botany Department, Faculty of Science, Urmia University scientifically confirmed the identification of the collected plants.

Preparation of the aqueous extract

After collection, the fruits were dried for 7-10 days in the shade at room temperature. The dried fruits were subsequently ground and the powder stored in cloth bags at 5°C until transfer to the laboratory for extraction. The method for preparing dry water-soluble plant powders has been previously described (27). Briefly, dried plant material (25 g) was mixed in 250 mL of distilled water for 15 minutes at 100°C, followed by rapid filtration

through a crude cellulose filter, after which the solution was filtered through Whatman #1 filter paper. The resultant filtrate was freeze-dried and the powder stored at -18°C in a desiccant until use. The average (w/w) yield was 12.4%.

Animal model

Adult sexually mature male Wistar albino rats were obtained from the Animal Resources Center of the Science Faculty at Urmia University. All animals were housed in a specific pathogen-free environment under standard conditions of temperature ($22 \pm 2^\circ\text{C}$), relative humidity ($50 \pm 10\%$) and light (12 hour light/12 hour dark). A standard pellet diet and fresh drinking water were provided *ad libitum*. Animals were checked daily for occurrence of any toxic signs of drug delivery. All ethical considerations for the studies on animals were considered carefully and the experimental protocol was approved by the Ethics Committee for Research on Laboratory Animals at Urmia University.

Experimental protocol

After a seven-day acclimation period, we randomly divided the rats into four groups that consisted of eight animals each ($n=8$) as described below: control group (Cont), *Crataegus monogyna* group (Cr), Cs group (Cs) and Cs-*Crataegus monogyna* group (CsCr). The two experimental groups (Cs and CsCr) were gavaged with 40 mg/kg/day of Cs (Sandimmune®, Novartis Pharmaceutical Corp., Switzerland) that was dissolved in sesame oil. The controls received an equivalent amount of sesame oil. The Cr group was gavaged with *Crataegus monogyna* aqueous extract at a dose of 20 mg/kg/day.

The CsCr group also received the same dose of this extract four hours after Cs administration. The treatment period was 45 days. The protocol for this study, including doses and duration of treatment for Cs and *Crataegus monogyna*, was designed according to previous studies (28-30).

Sampling

Animals were euthanized by cervical dislocation following anesthesia with ketamine (75 mg/kg; IP) 24 hours after the last *Crataegus*

monogyna treatment. Following euthanasia, a ventral midline skin incision was made to access the abdomen and the epididymides were carefully separated from the testicles under 20x magnification with a stereo zoom microscope (Model TL2, Olympus Co., Tokyo, Japan).

Epididymal sperm characteristics

Epididymal sperm count

In order to evaluate the sperm parameters, we placed one caudal epididymis in 1 ml of rat 1-cell embryo culture medium (mR1ECM). The cauda was cut into 2-3 pieces and incubated at 37°C for 10 minutes in a CO₂ incubator to allow the sperm to swim out of the epididymal tubules. The epididymal sperm concentration was determined by hemocytometer as follows. We diluted the epididymal sperm to 1:20 in mR1ECM medium then transferred approximately 10 µl of the diluted specimen to each of the counting chambers of a hemocytometer, which was placed for 5 minutes in a humid chamber to prevent drying. The cells sedimented during this time and were counted with a light microscope at ×400. The sperm count was expressed as number of sperm per milliliter (31).

Epididymal sperm viability

A 20 µl of sperm suspension was mixed with an equal volume of 0.05% eosin-Y. After incubation for 2 minutes at room temperature, the slides were viewed by bright-field microscope with ×400 magnification. Dead sperm appeared pink whereas live sperms remained unstained. We counted 200 sperm for each sample then calculated the viability percentages (32).

Assessment of sperm DNA integrity and chromatin quality

Staining of spermatozoa with acridine orange

Acridine orange staining was used to assess the cauda epididymal sperm DNA integrity in the experimental groups. Thick smears were fixed in Carnoy's fixative that consisted of methanol: acetic acid in a 1:3 ratio for at least 2 hours. The slides were stained for 5 minutes and gently rinsed with deionized water. We evaluated 200 sperm under fluorescent microscope;

sperm cell heads with normal DNA integrity showed green fluorescence, whereas those with diminished DNA integrity stained orange-red (33).

Staining of spermatozoa with aniline blue

The acidic aniline blue (AAB) stain discriminates between lysine-rich histones and arginine/cysteine-rich protamines. This technique provides a specific positive reaction with lysine residues in nuclear histones and reveals differences in the basic nuclear protein composition of the sperm. Histone-rich nuclei of immature spermatozoa are rich in lysine and consequently absorb the blue stain. Protamine-rich nuclei of mature spermatozoa are rich in arginine and cysteine and contain relatively low levels of lysine which leads to a negative reaction to aniline blue. In this study, we stained the air-dried fixed smears for 7 minutes with 0.5% aniline blue in phosphate buffered saline. The pH was adjusted to 3.5 by acetic acid. Slides were gently rinsed in distilled water and air dried. The percentage of spermatozoa stained with aniline blue was determined by counting 200 spermatozoa per slide under bright field microscope (34).

Assisted reproductive technique (ART) procedure

Oocyte collection

Each female rat was injected subcutaneously with 30 IU pregnant mare's serum gonadotropin (PMSG, The Netherlands) 48 hours prior to receiving an intraperitoneal injection of 15 IU human chorionic gonadotropin (hCG, Folligon, The Netherlands). Rats were euthanized 16 hours after hCG administration. Their oviducts were removed and transferred to a petri dish that contained mR1ECM medium. Using a stereo zoom microscope (Model TL2, Olympus Co., Tokyo, Japan), the Ampullary portion was found and oocytes were dissected out.

Sperm preparation and insemination

Samples from each group were prepared from the sperm suspension as described earlier. Spermatozoa were obtained by swim-up and capacitated by incubation at 37°C and 5% CO₂ for at least 1 hour. The sperm that had a concentration of 1×10⁶ total sperm/ml were added to a 500 µl fertilization drop of oocyte-containing

mR1ECM medium. For each animal we divided a total of 20 oocytes into 10 drops. After six hours of incubation at 37°C under 5% CO₂, the cumulus cell-free fertilized oocytes were transferred to fresh drops of mR1ECM medium for embryo culture. We covered all of the medium droplets with mineral oil (35).

Assessment of fertilization and embryonic development

Fertilized oocytes were evaluated by appearance of the pronuclei and polar bodies under ×200 magnification with an inverted microscope.

After approximately 24 hours following the zygotes culture we assessed the two-cell embryo rate. *In vitro* embryonic development was evaluated at 120 hours under phase-contrast microscopy. Intact, fragmented and/or lysed embryos which did not develop were recorded as "arrested embryos". In the present study, the rate of cell lyses was recorded as follows: Type I: fully lysed, necrotic and/or fragmented embryos. Type II: embryos with partially lysed/fragmented blastomeres. Type III: embryos with some lysed/fragmented blastomeres and/or cytoplasmic vesicles (36).

Statistical analysis

Results are expressed as mean ± SD. Differences between groups were assessed by analysis of variance (ANOVA) using the SPSS software package for Windows. Statistical significance between groups was determined by Tukey multiple comparison post hoc test p values less than 0.05 were considered to be statistically significant.

Results

Epididymal sperm characteristics

As seen in table 1, Cs treatment by itself significantly decreased the mean percentage of sperm concentration and viability compared to the control group. Co-administration of *Crataegus monogyna* aqueous fruit extract significantly protected the Cs-induced negative changes in sperm quantity and viability versus the Cs group.

Administration of Cs alone led to a significant increase in the formation of DNA strand breaks in the epididymal sperm compared to the control group. Cs plus *Crataegus monogyna* aqueous fruit extract treatment markedly suppressed Cs-induced DNA damage. The percentage of immature spermin Cs-treated rats was higher than those of the controls. Administration of *Crataegus monogyna* aqueous fruit extract along with Cs improved the chromatin quality of epididymal sperm compared with the Cs-treated group (Table 1).

Fertilization and embryonic development

Table 2 shows the changes in the fertilization rate and embryonic development in response to various treatments. While Cs administration alone caused significant decrease in the rates of fertilization, two-cell embryos and blastocysts with an increase in percentage of arrested embryos compared to the control group, Cs plus *Crataegus monogyna* aqueous fruit extract treatment showed a marked amelioration in these parameters (Table 2).

Table 1: Effect of cyclosporine (Cs) and *Crataegus monogyna* aqueous fruit extract on epididymal sperm characteristics

	Control	Cs	<i>Crataegus monogyna</i>	Cs + <i>Crataegus monogyna</i>
Sperm count (10 ⁶ /ml)	51.75 ± 3.24	19.50 ± 3.16 ^a	52.62 ± 4.92 ^b	46.12 ± 3.94 ^{a,b}
Viability (%)	87.75 ± 2.60	68.87 ± 3.31 ^a	85.62 ± 5.20 ^b	83.62 ± 4.98 ^b
Positive for acridine orange stain (%)	6.50 ± 2.20	28.37 ± 3.92 ^a	5.87 ± 2.64 ^b	11.25 ± 4.02 ^{a,b}
Positive for aniline blue stain (%)	8.75 ± 2.71	19.50 ± 3.20 ^a	9.37 ± 3.33 ^b	15.12 ± 3.22 ^{a,b}

Values are expressed as mean ± SD (n=8).

^a; Significant differences as compared with the control group at p<0.05 and ^b; Significant differences as compared with the Cs group at p<0.05.

Table 2: Effect of cyclosporine (Cs) and *Crataegus monogyna* aqueous fruit extract on fertilization and embryonic development

	Control	Cs	<i>Crataegus monogyna</i>	Cs + <i>Crataegus monogyna</i>
Fertilized oocytes (%)	96.13 ± 0.45	62.49 ± 8.84 ^a	96.55 ± 0.59 ^b	82.07 ± 4.46 ^{a,b}
2-cell embryos (%)	94.87 ± 0.38	57.88 ± 10.36 ^a	92.17 ± 3.05 ^b	90.59 ± 2.78 ^b
Blastocysts (%)	72.66 ± 1.66	24.49 ± 4.01 ^a	67.60 ± 1.06 ^b	70.51 ± 1.60 ^b
Arrested embryos (%)	27.32 ± 1.66	75.49 ± 4.01 ^a	32.39 ± 1.05 ^b	29.48 ± 1.60 ^b
Arrest type I (%)	4.74 ± 0.27	54.83 ± 8.03 ^a	6.78 ± 2.32 ^b	4.06 ± 1.11 ^b
Arrest type II (%)	5.15 ± 1.37	11.83 ± 2.12 ^a	7.48 ± 1.22 ^b	7.16 ± 0.44 ^b
Arrest type III (%)	17.42 ± 2.76	8.82 ± 2.77 ^a	18.11 ± 2.39 ^b	18.24 ± 2.21 ^b

The values are expressed as mean ± SD (n=8).

^a;Significant differences as compared with the control group at $p < 0.05$ and ^b;Significant differences as compared with the Cs group at $p < 0.05$.

Discussion

Cs is the most frequently used medication for the management of solid organ transplantation and immune diseases because of its potent immunosuppressive effect (37). However, effective clinical use of Cs is limited due to adverse effects such as reproductive toxicity as documented in humans and experimental animals (9-13). Hence, a strategy that diminishes the side-effects of Cs but preserves its therapeutic efficacy is necessary. Previous studies have shown that ROS-induced oxidative stress and lipid peroxidations may be involved in the pathophysiology of Cs reproductive toxicity (13, 38).

Spermatozoa are germ cells that have a fundamental role in fertilization. Damaged sperm are one of the factors that cause infertility. Mammalian spermatozoa are particularly vulnerable to excessive ROS-induced oxidative damage because of the molecular anatomy of their plasma membranes (39). The sperm cell membrane contains large quantities of polyunsaturated fatty acids which are involved in regulation of sperm maturation, spermatogenesis, capacitation, acrosome reaction and membrane fusion (40). ROS are highly reactive and can attack the unsaturated bonds of the membrane lipids, leading to lipid peroxidation (41). Peroxidation of sperm lipids demolishes the structure of the lipid matrix in the membranes of spermatozoa, which leads to decreased sperm viability and impaired spermatogenesis (42). In the present study Cs treatment has significantly decreased epididymal sperm count and viability

compared to the control group, which confirmed previous reports that Cs induced spermatozoal damage (11, 17-19, 43).

It has been well documented that oxidative damage to mature sperms during their migration from the seminiferous tubules to the epididymides forms immature sperms as an important factor involved in reproductive disorders (44). It has been shown that oxidative stress affects the integrity of the sperm's genome by causing single and double-strand DNA break increase. As shown in the present study, exposure to Cs resulted in a significant increase in spermatozoa DNA strand breaks and in the percentage of immature sperm compared to the control. The results of the current study supported the above mentioned findings.

There is increasing evidence that failures in fertilization and embryo development are associated with reduced sperm quality and increased rate of DNA damage in spermatozoa (46-48). Previous studies have shown a correlation between poor blastocyst formation and increased sperm DNA damage (49). Protamine deficiency and increased histone remnants in sperm cells result in an impaired fertilization process and embryo development (50-52).

Damaged spermatozoa are the source of free radicals (53), therefore Cs-induced embryo damage during *in vitro* fertilization could be related to the deleterious effect of ROS-producing dam-

aged spermatozoa during *in vitro* insemination of oocytes. It has been shown that *in vitro* incubation of oocytes with ROS-producing spermatozoa lead to oxidative damage of the oocytes or pronucleate embryos, eventually resulting in impaired embryo development (54, 55).

To date a number of studies have shown the benefits of anti-oxidants in protecting the reproductive system from the deleterious effects of free radicals generated during Cs exposure. Ellagic acid, a phytochemical with radical scavenging capacity, has recently been shown to ameliorate Cs-induced testicular and spermatozoal damages associated with oxidative stress (56). Lycopene has a potential protective effect against Cs-induced reproductive toxicity due to its highly efficient anti-oxidant properties (13). According to evidence, supplementation with an antioxidant food such as black grapes is protective against Cs-induced oxidant stress and peroxidation in rat ovary tissues (12).

In the present study, co-administration of *Crataegus monogyna* aqueous fruit extract provided effective protection against sperm impairment and embryo toxicity in Cs-treated rats. This status corroborated the fact that *Crataegus monogyna* was protective where free radicals had a crucial role in toxicity. Previous studies have shown that hawthorn extract can effectively lessen the extent of oxidative stress in mouse bone marrow cells and the rat brain due to its strong antioxidant activity (57, 58). Recent reports have also demonstrated that *Crataegus monogyna* fruit aqueous extract can effectively prevent oxidative damage to the reproductive system (59, 60).

Conclusion

The findings of our study have indicated that Cs-induced oxidative stress can lead to sperm impairment and embryotoxicity. Concurrent administration of *Crataegus monogyna* aqueous fruit extract effectively counteracted Cs-related oxidative injury through restoration of an anti-oxidant defense system.

Acknowledgements

The authors gratefully acknowledge the financial assistance of Urmia University in performing this investigation. The authors declare that

there are no conflicts of interest in this study.

References

1. Kahan BD. Cyclosporine: a revolution in transplantation. *Transplant Proc.* 1999; 31(1-2A): 14S-15S.
2. Kahan BD, Flechner SM, Lorber MI, Golden D, Conley S, Van Buren CT. Complications of cyclosporine-prednisone immunosuppression in 402 renal allograft recipients exclusively followed at a single center for from one to five years. *Transplantation.* 1987; 43(2), 197-204.
3. Berg KJ, Forre O, Bjerkhoe F, Amundsen E, Djoseland O, Rugstad HE, et al. Side effects of cyclosporine A treatment in patients with rheumatoid arthritis. *Kidney Int.* 1986; 29(6): 1180-1187.
4. Kaçmaz RO, Kempen JH, Newcomb C, Daniel E, Gangaputra S, Nussenblatt RB, et al. Cyclosporine for ocular inflammatory diseases. *Ophthalmology.* 2010; 117(3): 576-584.
5. Mihatsch MJ, Wolff K. Consensus conference on cyclosporine A for psoriasis February 1992. *Br J Dermatol.* 1992; 126(6): 621-623.
6. Yoshinoya S, Yamamoto K, Mitamura T, Aikawa T, Takeuchi A, Takahashi K, et al. Successful treatment of rheumatoid arthritis with low-dose cyclosporine A. *Transplant Proc.* 1988; 20(3 Suppl 4): 243-247.
7. Nabel GJ. A transformed view of cyclosporine. *Nature.* 1999; 397(6719): 471-472.
8. Yüce A, Ateşşahin A, Çeribaşı AO. Amelioration of cyclosporine A-induced renal, hepatic and cardiac damages by ellagic acid in rats. *Basic Clin Pharmacol Toxicol.* 2008; 103(2): 186-191.
9. Srinivas M, Agarwala S, Datta Gupta S, Das SN, Jha P, Misro MM, et al. Effect of cyclosporine on fertility in male rats. *Pediatr Surg Int.* 1998; 13(5-6): 388-391.
10. Misro MM, Chaki SP, Srinivas M, Chaube SK. Effect of cyclosporine on human sperm motility *in vitro*. *Arch Androl.* 1999; 43(3): 215-220.
11. Xu LG, Xu HM, Zhang JR, Song QZ, Qi XP, Wang XH. Effects of different dosages of cyclosporine A on the semen parameters of renal transplant patients. *Zhonghua Nan Ke Xue.* 2003; 9(9): 679-680, 683.
12. Ergüder IB, Çetin R, Devrim E, Kiliçoğlu B, Avci A, Durak I. Effects of cyclosporine on oxidant/antioxidant status in rat ovary tissues: protective role of black grape extract. *Int Immunopharmacol.* 2005; 5(7-8): 1311-1315.
13. Türk G, Ateşşahin A, Sönmez M, Yüce A, Çeribaşı, AO. Lycopene protects against cyclosporine A-induced testicular toxicity in rats. *Theriogenology.* 2007; 67(4): 778-785.
14. Wang C, Salahudeen AK. Lipid peroxidation accompanies cyclosporine nephrotoxicity: effect of vitamin E. *Kidney Int.* 1995; 47(3): 927-934.
15. Parra T, de Arriba G, Conejo JR, Cantero M, Arribas I, Rodríguez-Puyol D, et al. Cyclosporine increases local glomerular synthesis of reactive oxygen species in rats: effect of vitamin E on cyclosporine nephrotoxicity. *Transpl.* 1998; 66(10): 1325-1329.
16. Durak I, Kaçmaz M, Cimen MY, Büyükköçak S, Elagün S, Öztürk HS. The effect of cyclosporine on antioxidant enzyme activities and malondialdehyde levels in rabbit hepatocytes. *Transpl Immunol.* 2002; 10(4): 255-258.
17. Seethalakshmi L, Menon M, Malhotra RK, Diamond DA. Effect of cyclosporine A on male reproduction in rats. *J Urol.* 1987; 138(4 Pt 2): 991-995.
18. Seethalakshmi L, Flores C, Diamond DA, Menon M. Reversal of the toxic effects of cyclosporine on male reproduction and kidney function of rats by simultaneous administration of hCG + FSH. *J Urol.* 1990; 144(6): 1489-

- 1492.
19. Seethalakshmi L, Flores C, Khauli RB, Diamond DA, Menon M. Evaluation of the effect of experimental cyclosporine toxicity on male reproduction and renalfun-
ction. Reversal by concomitant human chorionic gonadotropin administration. *Transplantation*. 1990; 49(1): 17-19.
 20. Masuda H, Fujihira S, Ueno H, Kagawa M, Katsuo-
ka Y, Mori H. Ultrastructural study on cytotoxic effects of cyclo-
sporine A in spermiogenesis in rats. *Med Electron Microsc-*
sc. 2003; 36(3): 183-191.
 21. Rigelsky JM, Sweet BV. Hawthorn: pharmacology and
therapeutic uses. *Am J Health Syst Pharm*. 2002; 59(5):
417-422.
 22. Yao M, Ritchie HE, Brown-Woodman PD. A reproduc-
tive screening test of hawthorn. *J Ethnopharmacol*. 2008;
118(1): 127-132.
 23. Kéry A, Verzárné Petri G, Incze I. Comparative study of
flavonoids from *Crataegus oxyacantha* L. and *Crataegus*
monogyna Jacq. *Acta Pharm Hung*. 1977; 47(1): 11-23.
 24. Bahorun T, Trotin F, Pommery J, Vasseur J, Pinkas M.
Antioxidant activities of *Crataegus monogyna* extracts.
Planta Med. 1994; 60(4): 323-328.
 25. Bahorun T, Greiser B, Trotin F, Brunet C, Dine T, Luyckx
M, et al. Oxygen species scavenging activity of phenolic
extracts from hawthorn fresh plant organs and pharma-
ceutical preparations. *Arzneimittelforschung*. 1996; 46(11):
1086-1089.
 26. Rakotoarison DA, Greissier B, Trotin F, Brunet C, Dine
T, Luyckx M, et al. Antioxidant activities of polyphenolic
extracts from flowers, in vitro callus and cell suspension
cultures of *Crataegus monogyna*. *Pharmazie*. 1997; 52(1):
60-64.
 27. Ljubuncic P, Azaizeh H, Portnaya I, Cogan U, Said O,
Saleh KA, et al. Antioxidant activity and cytotoxicity of
eight plants used in traditional Arab medicine. *J Ethno*
pharmacol. 2005; 99(1): 43-47.
 28. Reichenspurner H, Hildebrandt A, Human PA, Boehm DH,
Rose AG, Odell JA, et al. 15-Deoxyspergualin for induction
of graft non reactivity after cardiac and renal allotransplan-
tation in primates. *Transplantation*. 1990; 50(2): 181-185.
 29. Dehpour AR, Shirzad N, Ghafourifar P, Abdollahi M. Ef-
fects of cyclosporine A on the functions of submandibular
and parotid glands of rats. *Gen Pharmacol*. 1996; 27(5):
887-890.
 30. Khalil R, Abuharfeil N, Shabsoug B. The effect of *Cratae-*
gus aronica aqueous extract in rabbits fed with high cho-
lesterol diet. *Eur J Sci Res*. 2008; 22(3): 352-360.
 31. Zambrano E, Rodriguez-Gonzalez GL, Guzman C, Gar-
cia-Becerra R, Boeck L, Menjivar M, et al. A maternal
low protein diet during pregnancy and lactation in the rat
impairs male reproductive development. *J Physiol*. 2005;
563(1): 275-284.
 32. Wyrobek AJ, Gordon LA, Burkhart JG, Francis MW, Kap-
p Jr RW, Letz G, et al. An evaluation of the mouse sperm
morphology test and other sperm tests in nonhuman
mammals. A report of the US environmental protection
agency gene-tox program. *Mutat Res*. 1983; 115(1): 1-72.
 33. Erenpreiss J, Bars J, Lipatnikova V, Erenpreisa J, Zalkalns
J. Comparative study of cytochemical tests for sperm chro-
matin integrity. *J Androl*. 2001; 22(1): 45-53.
 34. Hammadeh ME, al-Hasani S, Stieber M, Rosenbaum P,
Küpker D, Diedrich K, et al. The effect of chromatin con-
densation (aniline blue staining) and morphology (strict
criteria) of human spermatozoa on fertilization, cleavage
and pregnancy rates in an intracytoplasmic sperm injec-
tion programme. *Hum Reprod*. 1996; 11(11): 2468-2471.
 35. Toyoda Y, Chang MC. Fertilization of rat eggs in vitro by
epididymal spermatozoa and the development of eggs fol-
lowing transfer. *J Reprod Fertil*. 1974; 36(1): 9-22.
 36. Cebal E, Carrasco I, Vantman D, Smith R. Preimplanta-
tion embryotoxicity after mouse embryo exposition to re-
active oxygen species. *Biocell*. 2007; 31(1): 51-59.
 37. Kahan BD. Immunosuppressive therapy. *Curr Opin Immu-
nol*. 1992; 4(5): 553-560.
 38. Sikka SC, Coy DC, Lemmi CA, Rajfer J. Effect of cyclo-
sporine on steroidogenesis in rat leydig cells. *Transplanta-
tion*. 1988; 46(6): 886-890.
 39. Vernet P, Aitken RJ, Drevet JR. Antioxidant strategies in
the epididymis. *Mol Cell Endocrinol*. 2004; 216(1-2): 31-
39.
 40. Agarwal A, Makker K, Sharma R. Clinical relevance of
oxidative stress in male factor infertility: an update. *Am J*
Reprod Immunol. 2008; 59(1): 2-11.
 41. Sanocka D, Kurpisz M. Reactive oxygen species and
sperm cells. *Reprod Biol Endocrinol*. 2004; 2(12): 1-7.
 42. Türk G, Ateşşahin A, Sönmez M, Çeribaşı AO, Yüce A. Im-
provement of cisplatin-induced injuries to sperm quality,
the oxidant-antioxidant system, and the histologic struc-
ture of the rat testis by ellagic acid. *Fertil Steril*. 2008; 89(5
Suppl): 1474-1481.
 43. Iwasaki M, Fuse H, Katayama T. The effects of cyclo-
sporine azathioprine and mizoribine on male reproduction
in rats. *Nihon Hinyokika Gakkai Zasshi*. 1996; 87(1): 42-
49.
 44. Agarwal A, Said TM. Role of sperm chromatin abnormali-
ties and DNA damage in male infertility. *Hum Reprod Up-
date*. 2003; 9(4): 331-345.
 45. Moustafa MH, Sharma RK, Thornton J, Mascha E, Abdel-
Hafez MA, Thomas AJ Jr, et al. Relationship between
ROS production, apoptosis and DNA denaturation in
spermatozoa from patients examined for infertility. *Hum*
Reprod. 2004; 19(1): 129-138.
 46. Ron-el R, Nachum H, Herman A, Golan A, Caspi E, Soffer
Y. Delayed fertilization and poor embryonic development
associated with impaired semen quality. *Fertil Steril*. 1991;
55(2): 338-344.
 47. Høst E, Lindenberg S, Smidt-Jensen S. The role of DNA
strand breaks in human spermatozoa used for IVF and
ICSI. *Acta Obstet Gynecol Scand*. 2000; 79(7): 559-563.
 48. Erenpreiss J, Spano M, Erenpreisa J, Bungum M, Gi-
wercman A. Sperm chromatin structure and male fertility:
biological and clinical aspects. *Asian J Androl*. 2006; 8(1):
11-29.
 49. Meseguer M, Martínez-Conejero JA, O'Connor JE, Pel-
licer A, Remohí J, Garrido N. The significance of sperm
DNA oxidation in embryo development and reproductive
outcome in an oocyte donation program: a new model to
study a male infertility prognostic factor. *Fertil Steril*. 2008;
89(5): 1191-1199.
 50. Sakkas D, Urner F, Bizzaro D, Manicardi G, Bianchi PG,
Shoukir Y, et al. Sperm nuclear DNA damage and altered
chromatin structure: effect on fertilization and embryo de-
velopment. *Hum Reprod*. 1998; 13 Suppl 4: 11-19.
 51. Cho C, Jung-Ha H, Willis WD, Goulding EH, Stein P, Xu
Z, et al. Protamine 2 deficiency leads to sperm DNA dam-
age and embryo death in mice. *Biol Reprod*. 2003; 69(1):
211-217.
 52. Oliva R. Protamines and male infertility. *Hum Reprod Up-
date*. 2006; 12(4): 417-435.
 53. Said TM, Agarwal A, Sharma RK, Thomas AJ Jr, Sikka
SC. Impact of sperm morphology on DNA damage caused
by oxidative stress induced by beta-nicotinamide adenine
dinucleotide phosphate. *Fertil Steril*. 2005; 83(1): 95-103.
 54. Nasr-Esfahani MH, Aitken JR, Johnson MH. Hydrogen
peroxide levels in mouse oocytes and early cleavage
stage embryos developed in vitro or in vivo. *Development*.

C. Monogyna Attenuates Cs-Induced Reproductive Toxicity

- 1990; 109(2): 501-507.
55. Paszkowski T, Clarke RN. The graafian follicle is a site of L-ascorbate accumulation. *J Assist Reprod Genet.* 1999; 16(1): 41-45.
56. Türk G, Sönmez M, Ceribaşı AO, Yüce A, Ateşşahin A. Attenuation of cyclosporine A-induced testicular and spermatozoal damages associated with oxidative stress by ellagic acid. *Int Immunopharmacol.* 2010; 10(2): 177-182.
57. Hosseinimehr SJ, Azadbakht M, Abadi AJ. Protective effect of hawthorn extract against genotoxicity induced by cyclophosphamide in mouse bone marrow cells. *Environ Toxicol Pharmacol.* 2008; 25(1): 51-56.
58. Elango C, Jayachandaran KS, Niranjali Devaraj S. Hawthorn extract reduces infarct volume and improves neurological score by reducing oxidative stress in rat brain following middle cerebral artery occlusion. *Int J Dev Neurosci.* 2009; 27(8): 799-803.
59. Jalali AS, Hasanzadeh S, Malekinejad H. Chemoprotective effect of *Crataegus monogyna* aqueous extract against cyclophosphamide-induced reproductive toxicity. *Veterinary Research Forum.* 2011; 2(4): 266-273.
60. Jalali AS, Hasanzadeh S, Malekinejad H. *Crataegus monogyna* aqueous extract ameliorates Cyclophosphamide-induced Toxicity in rat testis: Stereological evidences. *Acta Med Iran.* 2012; 50(1): 1-8.
-

Archive of SID