

Comment on: Anti-Oxidative Stress Activity of *Stachys lavandulifolia* Aqueous Extract in Humans

Azar Baradaran, Ph.D.¹, Hamid Nasri, M.D.², Mahmoud Rafeiean-Kopaei, P.h.D.^{3*}

1. Department of Pathology, Isfahan University of Medical Sciences, Isfahan, Iran

2. Department of Nephrology, Division of Nephropathology, Isfahan University of Medical Sciences, Isfahan, Iran

3. Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

* Corresponding Address: P.O.Box: 8136893669, Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran
Email: rafeiean@yahoo.com

Received: 11/May/2013, Accepted: 29/Jun/2013

A recent article by Rahzani et al. (1) published in the esteemed Cell Journal reported the anti-oxidative stress activity of *Stachys lavandulifolia* aqueous extract in humans and suggested its consumption as a supplement in the management of diseases related to oxidative stress. We would like to emphasize some of the limitations regarding antioxidant supplementation, in general, and *Stachys lavandulifolia*, in particular.

It has been established that oxidative stress is involved in the development of a wide variety of chronic and degenerative diseases such as cancer, Parkinson and Alzheimer's (2-5). Antioxidants are also effective in the prevention or reduction of adverse effects related to medication usage (5-10). However, they may potentially have deleterious effects. A major concern of antioxidant supplementation is their harmful effect on reactive oxygen species (ROS) production (pro-oxidant action), particularly when precise modulation of ROS levels are necessary for normal cell function (4-10). In fact, it has been reported that antioxidants may exhibit pro-oxidant activity under specific conditions. Of particular importance are redox conditions the dosage and the presence of free transition metals at cellular sites. For example, the antioxidant vitamin C in the presence of ferric iron may act as a potent mediator of lipid peroxidation. It has been suggested that β -carotene sometimes acts as a pro-oxidant in the lungs of smokers and similarly vitamin C may increase DNA damage in humans (11, 12). Therefore,

it is necessary to take into account the bioavailability and differential activities of antioxidant compounds before their administration.

Other than general considerations for antioxidant consumption, the aspects of each particular antioxidant should also be considered (3, 13, 14). Recently, in a preclinical study we reported the renal toxicity of hydroalcoholic extract of *Stachys lavandulifolia* Vahl in Wistar rats (15). In this experimental study we randomly assigned 100 male Wistar rats to five equal groups, one control and four experimental. Animals received intraperitoneal injections of saline or *Stachys lavandulifolia* extract (50, 100, 150, 200 mg/kg) for one month after which blood samples were collected from half of the animals from each group. Other animals received no injections for one additional month, then blood samples were obtained. In the groups that *Stachys lavandulifolia* Vahl extracts were used for one month we observed mild degeneration of renal tubular epithelial cells (6, 9). In the second month of the study these histologic lesions significantly increased ($p < 0.05$). We concluded that hydroalcoholic extract of *Stachys lavandulifolia* has renal tubular toxicity which might continue following drug discontinuation (6, 9, 15).

Therefore, although antioxidant supplements generally have beneficial effects, as a caution it is advised to only consume such supplements under medical supervision in order to avoid any potential negative effects.

Keywords: *Stachys lavandulifolia*, Toxicity, Nephrotoxicity, Antioxidant

Cell Journal (Yakhteh), Vol 15, No 3, Autumn 2013, Pages: 272-273

Citation: Baradaran A, Nasri H, Rafeiean-Kopaei M. Comment on: anti-oxidative stress activity of *stachys lavandulifolia* aqueous extract in humans. Cell J. 2013; 15(3): 272-273.

References

1. Rahzani K, Malekiran AA, Zeraatpishe A, Hosseini N, Seify SMR, Abdollahi M. Anti-oxidative stress activity of *Stachys lavandulifolia* aqueous extract in humans. *Cell J*. 2013; 14(4): 314-317.
2. Tavafi M. Diabetic nephropathy and antioxidants. *J Nephropathology*. 2013; 2(1): 20-27.
3. Khajehdehi P. Turmeric: reemerging of a neglected Asian traditional remedy. *J Nephropathology*. 2012; 1(1): 17-22.
4. Valko M, Leibfritz D, Moncol J, Cronin MTD, Mazur M, Telser J. Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol*. 2007; 39 (1): 44-84.
5. Ghorbani A, Omidvar B, Parsi A. Protective effect of selenium on cisplatin induced nephrotoxicity: a double-blind controlled randomized clinical trial. *J Nephropathology*. 2013; 2(2): 129-134.
6. Rafeian-Kopaei M, Nasri H, Nematbakhsh M, Baradaran A, Gheissari A, Rouhi H, et al. Erythropoietin ameliorates gentamicin-induced renal toxicity: a biochemical and histopathological study. *J Nephropathology*. 2012; 1(2): 109-116.
7. Shirzad H, Taji F, Rafeian-Kopaei M. Correlation between antioxidant activity of garlic extracts and WEHI-164 fibrosarcoma tumor growth in BALB/c mice. *J Med Food*. 2011; 14: 969-974.
8. Tavafi M. Inhibition of gentamicin-induced renal tubular cell necrosis. *J Nephropathology*. 2012; 1(2): 83-86.
9. Nematbakhsh M, Ashrafi F, Pezeshki Z, Fatahi Z, Kianpoor F, Sanei MH, et al. A histopathological study of nephrotoxicity, hepatotoxicity or testicular toxicity: which one is the first observation as side effect of Cisplatin-induced toxicity in animal model. *J Nephropathology*. 2012; 1(3): 190-193.
10. Rafeian Kopaei M, Baradaran A, Merrikhi A, Nematbakhsh M, Madihi Y, Nasri H. Efficacy of co administration of garlic extract and metformin for prevention of gentamicin-renal toxicity in wistar rats: a biochemical study. *Int J Prev Med*. 2013; 4(3): 258-264.
11. Galati G, Sabzevari O, Wilson JX, O'Brien PJ. Prooxidant activity and cellular effects of the phenoxyl radicals of dietary flavonoids and other polyphenolics. *Toxicology*. 2002; 177(1): 91-104.
12. Fox RB. Prevention of granulocyte-mediated oxidant lung injury in rats by a hydroxyl radical scavenger, dimethyl thiourea. *J Clin Invest*. 1984; 74(4): 1456-1464.
13. Rafeian-Kopaei M, Baradaran A. Teucrium polium and kidney. *J Ren Inj Prev*. 2013; 2(1): 3-4.
14. Kadkhodae M. Erythropoietin; bright future and new hopes for an old drug. *J Nephropathology*. 2012; 1(2): 81-82.
15. Taghikhani M, Nasri H, Asgari A, Afrough H, Namjoo AR, Ansari-Samani R, et al. The renal toxicity of hydroalcoholic extract of *Stachys lavandulifolia* vahl in wistar rats. *Life Sci J*. 2012; 9(4): 3025-3031.