

# Isolation and characterization of p-Coumaric acid from *Diospyros melanoxylon* medicinal plant endemic to Western Ghats, India

Anil Kumar K. M<sup>1,2</sup>, Pallavi S<sup>1,2</sup>, Shivaraju Harikaranahalli Puttaiah<sup>1,✉</sup>

1. Department of Water and Health, JSS Academy of Higher Education and Research, Mysore, India
2. Center for Water, Food and Energy, GREENS trust, Harikaranahalli, Dombaranahalli Post, Turuvekere Taluka, Tumkur District, Karnataka, India

**Date of submission:** 31 Dec 2018, **Date of acceptance:** 12 Jun 2019

## ABSTRACT

*Diospyros melanoxylon* has been traditionally for the treatment of fistula, relieving arthritis, and skin care. The bark extracts of the plant have been investigated in terms of phytochemical and pharmacological potential, while the leaf extract has been untapped. The present study aimed to evaluate the methanol extract of *D. melanoxylon* (DMM) in terms of the antibacterial ( $P < 0.05$ ), antioxidant (1.2-1.6-fold) and anti-inflammatory potential ( $IC_{50}$ : 80  $\mu$ g/ml). DMM exhibited effective antibacterial, antioxidant, and anti-inflammatory activities at significantly higher levels than the standards. In addition, the HR-LCMS analysis of MBI revealed the presence of a few active compounds, which belonged to the class of phenolic acids and flavonoids at greater concentrations than other phytochemicals ( $n > 20$ ). The activity-guided repeated fractionation of the methanol extract using silica gel column chromatography yielded a single compound, which exhibited remarkable antioxidant activity. The physicochemical and spectroscopic analyses (UV, IR, <sup>1</sup>H NMR, <sup>13</sup>C NMR, and MS) indicated that the bioactive isolated compound was p-coumaric acid, the effect of which was on par with the standard antioxidant, antibacterial, and anti-inflammatory drugs. Conversely, the effects of the extract on these pharmacological attributes enhanced, confirming that the better activity observed in the study was mainly due to the synergistic effects exerted by various compounds in the extract. *In-silico* studies have also confirmed the potential of the compound in these effective antibacterial properties. Therefore, the *D. melanoxylon* extract is a strong therapeutic agent with pharmacological potential.

**Keywords:** Phenolic acids, p-Coumaric acid, Methanol extract, Free radicals, Anti-inflammatory

## Introduction

*Diospyros melanoxylon* is an endemic species of the Western Ghats region in India and Sri Lanka, which belongs to the Ebenaceae family. This plant was first described by William Roxburgh in his book Flora Indica.<sup>1</sup> Most of the species in this family are a valuable source of wood and ornamental flowers. Most species of the *Diospyros* genus are mainly confined to the tropics, with approximately 300 species found in Asia.<sup>2</sup> India alone houses 66 species belonging to this genus, which are

known for their therapeutic potential.<sup>3</sup> The World Conservation Monitoring Centre has recorded 12 major biodiversity hotspots, among which India contains two, including Western Ghats and Eastern Ghats, containing 15,000-18,000 flowering plants.

Among various regions of the Western Ghats, the Shimoga district of the Malnad ranges is less exploited, which is located in the heart of the Western Ghats region in the Karnataka state; it is one of the biodiversity hotspots in India. In our previous study, a thorough screening of various plants in this region was performed, providing evidence on the least exploited species. In the current research, we aimed to evaluate the therapeutic potential of *D. melanoxylon*, which is extensively found in the Western Ghats region

✉ Shivaraju Harikaranahalli Puttaiah  
shivarajuenvi@gmail.com

**Citation:** Anil Kumar K.M, Pallavi S, Puttaiah Sh.H. Isolation and characterization of p-Coumaric acid from *Diospyros melanoxylon* medicinal plant endemic to Western Ghats, India. J Adv Environ Health Res 2019; 7(4): 269-282

of the Southern India. Apart from its use as timber and a material to wrap beedi, this endemic species is known for its numerous pharmacological properties (antioxidant, antidiabetic, and nephroprotective properties), which are mainly attributed to the presence of polyphenols, flavonoids, alkaloids, and anthocyanins.<sup>4</sup> Traditionally, *D. melanoxydon* has been used for the treatment of arthritis, relieving fistula and abdominal pain, and skin care.

Large amounts of phenolics, alkaloids, steroids, ascorbic acid, tannins, and saponins have been observed in the bark extract of *D. melanoxydon*.<sup>5</sup> Formulations prepared from the leaves of the plant have also been examined in terms of the effects on the improvement of intestinal flatulence and digestion issues. In the present study, the optimum activity among various bioactive compounds in the extracts of *D. melanoxydon* was observed in the extract containing p-coumaric acid. This secondary metabolite is a phenolic acid belonging to the hydroxycinnamic acid family, and p-coumaric acid is essential to secondary metabolism since it could be converted into phenolic acid, flavonoid, and lignin derivatives.<sup>6,7</sup>

The chemoprotectant and the antioxidant potential of p-coumaric acid have attracted the attention of researchers to identify the plants that could be used as potential sources for the extraction of p-coumaric acid.<sup>8</sup> Furthermore, p-coumaric acid is known for its potent antibacterial activity against a broad spectrum of pathogens in synergy with other compounds.<sup>9</sup>

Based on the previous reports and a preliminary phytochemical screening in our study, the present study aimed to describe the isolation and characterization of p-coumaric acid from the bark of *D. melanoxydon* and assess its potential as an antimicrobial, antioxidant, and anti-inflammatory agent.

## Materials and Methods

### Plant materials and extraction

Fresh, healthy leaves of *D. melanoxydon* were collected from the Western Ghats in Shimoga district, located in Karnataka, India during May 2017. The leaves were washed,

shade dried, and milled using a 50 mesh.

### Isolation and identification of the bioactive compound

At this stage, 500 grams of the dried leaves were used to extract the bioactive compound sequentially using the solvents of increasing polarity viz., petroleum ether, chloroform, ethyl acetate, acetone, methanol, and water. Afterwards, the diverse extracts were filtered, and the solvent was excluded using a rotary evaporator (model: Rotavapor R-200, Buchi, Switzerland). The concentrated extracts were assayed in terms of the pharmacological activities viz. and antibacterial, anti-inflammatory, and radical-scavenging properties (DPPH, ABTS, and superoxide). Since the yield (145 g/kg) and radical scavenging activities of the methanol extract were higher, the extract was selected for the isolation of the bioactive compound. The phytochemical screening for the methanol extract was also performed in accordance with the standard protocols as described by Harborne *et al.*<sup>10</sup>. In order to isolate the bioactive compound, 50 grams of *D. melanoxydon* methanol extract (DMM) was dispersed in water and successively extracted twice, each time using petroleum ether, ethyl acetate, and n-butanol to obtain the yield of petroleum ether (6.3 g), ethyl acetate (7.9 g), n-butanol (3.7 g), and H<sub>2</sub>O-soluble (15 g) fractions. The subsequent fractions were subjected to pharmacological activities, while the activities of ethyl acetate soluble fraction (ESF) were applied for the isolation of the bioactive compound.

At the next stage, the ESF was subjected to silica gel column chromatography (100-200 mesh, 1.5 kg; length: 80 cm, diameter: 7 cm; elution rate: 2 ml/min, flow total elution: 500 ml) and eluted using a gradient of dichloromethane (methanol 100:1 [5 l], 90:10 [10 l], 80:20 [5 l], 70:30 [10 l], 60:40 [5 l], 50:50 [5 l], 25:75 [5 l], 0:100 [3 l]) in order to acquire fractions ESF 1 (2.2 g), ESF 2 (4.0 g), ESF 4 (6.6 g), ESF 4 (3.5 g), ESF 5 (6.9 g), ESF 6 (5.3 g), ESF 7 (4.4 g), and ESF 8 (5.7 g). Fractions ESF 1, ESF 2, and ESF 5 exhibited potent

biological activities with the following order: ESF 5>ESF 1>ESF 2. As a result, fraction ESF 5 was subjected to the silica gel column chromatography (length: 50 cm, diameter: 3 cm; elution rate: 1 ml/min; flow total elution: 100 ml) and eluted with the linear gradients of petroleum ether (acetone [95:5, 90:10, 85:15, 80:20, 70:30, and 60:40 v/v]) to obtain six major sub-fractions. Sub-fraction three was further separated using the silica gel CC and petroleum ether acetone (85:15), followed by re-chromatography on a Sephadex LH-20 column with acetone as the eluting solvent to obtain p-coumaric acid as the single compound (24.5 mg).

### ***Thin-layer Chromatography***

The collected fractions were spotted onto silica gel F254 plates (25×25 cm; Merck, Germany). The solvent system contained chloroform, methanol, and formic acid (85:10:5 v/v/v), revealing optimum separation. Following that, the developed plates were air dried, and the spots were visualized by spraying a ferric chloride solution (0.5% w/v). The retention factor ( $R_f$ ) of the isolated compound ( $R_f=0.79$ ) and standard value ( $R_f=0.79$ ) were observed to be similar when calculated and compared.

### ***Identification of the bioactive compound using analytical methods***

The melting points were determined on an electrically heated VMP-III melting point apparatus in an uncorrected manner. The infrared (IR) spectra were recorded on the NICOLET 380 FT IR spectrometer (Thermo Fisher Scientific, France) using potassium bromide (KBr) pellets. In addition, the nuclear magnetic resonance (NMR) spectra was recorded on a Bruker DRX-400 spectrometer (Bruker Biospin Co., Karlsruhe, Germany) with  $^1\text{H}$  NMR at 400 MHz and  $^{13}\text{C}$  NMR at 100 MHz. Following that, the isolated compound was prepared using deuterated methanol (99.8 atom% of deuterium) and tetramethylsilane (TMS) as an internal standard in five-millimeter NMR tubes.

Data were measured in  $\text{CDCl}_3$  with

chemical shifts based on the TMS signal and expressed in parts per million ( $\delta$ ). The ultraviolet spectrum of p-coumaric acid in methanol was also recorded using a Shimadzu UV-1800 spectrophotometer. The elemental analysis of the compound was performed using the Perkin-Elmer 2400 elemental analyzer, and the mass spectrum was recorded using the Q-TOF Waters Ultima instrument (QTOF GAA 082, Waters, Manchester, UK) with an electron spray ionization source. The positive ion mode with the spray voltage of 3.5 kV at the source temperature of  $80^\circ\text{C}$  was set to obtain the spectra, and the mass spectra were recorded under electron impact ionization at the energy of 70 eV. The sample was prepared within the concentration range of 0.25-0.50 mg/ml and injected using the flow analysis at the flow rate of  $10 \mu\text{Lmin}^{-1}$ ; the recorded mass was within the range of 100-500 m/z.

### ***Determination of total phenol***

The total phenolic content (TPC) of the methanol extract and fractions was determined using the Folin-Ciocalteu method<sup>11</sup> and expressed as gallic acid equivalents (GAE) in milligrams per gram of the sample.

### ***Determination of total flavonoids***

The total flavonoid content (TFC) of the methanol extract and fractions was determined using the aluminum chloride technique<sup>12</sup> and expressed as quercetin equivalents (QE) in milligrams per gram of the sample.

### ***Determination of total proanthocyanidin***

The total proanthocyanidin (TCC) content of the methanol extract and fractions was determined using the method described in<sup>13</sup> and expressed as GAE in milligrams per gram of the sample.

### ***High-resolution Liquid Chromatography and Mass Spectrometry (HR-LCMS) analysis***

The fresh, healthy leaves of *D. melanoxylon* were extracted using methanol and subjected to high-resolution liquid chromatography and mass spectrometry (HR-LCMS) analysis, which was carried out at the

of the Southern India. Apart from its use as timber and a material to wrap beedi, this endemic species is known for its numerous pharmacological properties (antioxidant, antidiabetic, and nephroprotective properties), which are mainly attributed to the presence of polyphenols, flavonoids, alkaloids, and anthocyanins.<sup>4</sup> Traditionally, *D. melanoxylon* has been used for the treatment of arthritis, relieving fistula and abdominal pain, and skin care.

Large amounts of phenolics, alkaloids, steroids, ascorbic acid, tannins, and saponins have been observed in the bark extract of *D. melanoxylon*.<sup>5</sup> Formulations prepared from the leaves of the plant have also been examined in terms of the effects on the improvement of intestinal flatulence and digestion issues. In the present study, the optimum activity among various bioactive compounds in the extracts of *D. melanoxylon* was observed in the extract containing p-coumaric acid. This secondary metabolite is a phenolic acid belonging to the hydroxycinnamic acid family, and p-coumaric acid is essential to secondary metabolism since it could be converted into phenolic acid, flavonoid, and lignin derivatives.<sup>6,7</sup>

The chemoprotectant and the antioxidant potential of p-coumaric acid have attracted the attention of researchers to identify the plants that could be used as potential sources for the extraction of p-coumaric acid.<sup>8</sup> Furthermore, p-coumaric acid is known for its potent antibacterial activity against a broad spectrum of pathogens in synergy with other compounds.<sup>9</sup>

Based on the previous reports and a preliminary phytochemical screening in our study, the present study aimed to describe the isolation and characterization of p-coumaric acid from the bark of *D. melanoxylon* and assess its potential as an antimicrobial, antioxidant, and anti-inflammatory agent.

## **Materials and Methods**

### ***Plant materials and extraction***

Fresh, healthy leaves of *D. melanoxylon* were collected from the Western Ghats in Shimoga district, located in Karnataka, India during May 2017. The leaves were washed,

shade dried, and milled using a 50 mesh.

### ***Isolation and identification of the bioactive compound***

At this stage, 500 grams of the dried leaves were used to extract the bioactive compound sequentially using the solvents of increasing polarity *viz.*, petroleum ether, chloroform, ethyl acetate, acetone, methanol, and water. Afterwards, the diverse extracts were filtered, and the solvent was excluded using a rotary evaporator (model: Rotavapor R-200, Buchi, Switzerland). The concentrated extracts were assayed in terms of the pharmacological activities *viz.* and antibacterial, anti-inflammatory, and radical-scavenging properties (DPPH, ABTS, and superoxide). Since the yield (145 g/kg) and radical scavenging activities of the methanol extract were higher, the extract was selected for the isolation of the bioactive compound. The phytochemical screening for the methanol extract was also performed in accordance with the standard protocols as described by Harborne *et al.*<sup>10</sup>. In order to isolate the bioactive compound, 50 grams of *D. melanoxylon* methanol extract (DMM) was dispersed in water and successively extracted twice, each time using petroleum ether, ethyl acetate, and n-butanol to obtain the yield of petroleum ether (6.3 g), ethyl acetate (7.9 g), n-butanol (3.7 g), and H<sub>2</sub>O-soluble (15 g) fractions. The subsequent fractions were subjected to pharmacological activities, while the activities of ethyl acetate soluble fraction (ESF) were applied for the isolation of the bioactive compound.

At the next stage, the ESF was subjected to silica gel column chromatography (100-200 mesh, 1.5 kg; length: 80 cm, diameter: 7 cm; elution rate: 2 ml/min, flow total elution: 500 ml) and eluted using a gradient of dichloromethane (methanol 100:1 [5 l], 90:10 [10 l], 80:20 [5 l], 70:30 [10 l], 60:40 [5 l], 50:50 [5 l], 25:75 [5 l], 0:100 [3 l]) in order to acquire fractions ESF 1 (2.2 g), ESF 2 (4.0 g), ESF 4 (6.6 g), ESF 4 (3.5 g), ESF 5 (6.9 g), ESF 6 (5.3 g), ESF 7 (4.4 g), and ESF 8 (5.7 g). Fractions ESF 1, ESF 2, and ESF 5 exhibited potent

biological activities with the following order: ESF 5>ESF 1>ESF 2. As a result, fraction ESF 5 was subjected to the silica gel column chromatography (length: 50 cm, diameter: 3 cm; elution rate: 1 ml/min; flow total elution: 100 ml) and eluted with the linear gradients of petroleum ether (acetone [95:5, 90:10, 85:15, 80:20, 70:30, and 60:40 v/v]) to obtain six major sub-fractions. Sub-fraction three was further separated using the silica gel CC and petroleum ether acetone (85:15), followed by re-chromatography on a Sephadex LH-20 column with acetone as the eluting solvent to obtain p-coumaric acid as the single compound (24.5 mg).

#### **Thin-layer Chromatography**

The collected fractions were spotted onto silica gel F254 plates (25×25 cm; Merck, Germany). The solvent system contained chloroform, methanol, and formic acid (85:10:5 v/v/v), revealing optimum separation. Following that, the developed plates were air dried, and the spots were visualized by spraying a ferric chloride solution (0.5% w/v). The retention factor ( $R_f$ ) of the isolated compound ( $R_f=0.79$ ) and standard value ( $R_f=0.79$ ) were observed to be similar when calculated and compared.

#### **Identification of the bioactive compound using analytical methods**

The melting points were determined on an electrically heated VMP-III melting point apparatus in an uncorrected manner. The infrared (IR) spectra were recorded on the NICOLET 380 FT IR spectrometer (Thermo Fisher Scientific, France) using potassium bromide (KBr) pellets. In addition, the nuclear magnetic resonance (NMR) spectra was recorded on a Bruker DRX-400 spectrometer (Bruker Biospin Co., Karlsruhe, Germany) with <sup>1</sup>H NMR at 400 MHz and <sup>13</sup>C NMR at 100 MHz. Following that, the isolated compound was prepared using deuterated methanol (99.8 atom% of deuterium) and tetramethylsilane (TMS) as an internal standard in five-millimeter NMR tubes.

Data were measured in CDCl<sub>3</sub> with

chemical shifts based on the TMS signal and expressed in parts per million ( $\delta$ ). The ultraviolet spectrum of p-coumaric acid in methanol was also recorded using a Shimadzu UV-1800 spectrophotometer. The elemental analysis of the compound was performed using the Perkin-Elmer 2400 elemental analyzer, and the mass spectrum was recorded using the Q-TOF Waters Ultima instrument (QTOF GAA 082, Waters, Manchester, UK) with an electron spray ionization source. The positive ion mode with the spray voltage of 3.5 kV at the source temperature of 80°C was set to obtain the spectra, and the mass spectra were recorded under electron impact ionization at the energy of 70 eV. The sample was prepared within the concentration range of 0.25-0.50 mg/ml and injected using the flow analysis at the flow rate of 10  $\mu\text{lmin}^{-1}$ ; the recorded mass was within the range of 100-500 m/z.

#### **Determination of total phenol**

The total phenolic content (TPC) of the methanol extract and fractions was determined using the Folin-Ciocalteu method<sup>11</sup> and expressed as gallic acid equivalents (GAE) in milligrams per gram of the sample.

#### **Determination of total flavonoids**

The total flavonoid content (TFC) of the methanol extract and fractions was determined using the aluminum chloride technique<sup>12</sup> and expressed as quercetin equivalents (QE) in milligrams per gram of the sample.

#### **Determination of total proanthocyanidin**

The total proanthocyanidin (TCC) content of the methanol extract and fractions was determined using the method described in<sup>13</sup> and expressed as GAE in milligrams per gram of the sample.

#### **High-resolution Liquid Chromatography and Mass Spectrometry (HR-LCMS) analysis**

The fresh, healthy leaves of *D. melanoxylon* were extracted using methanol and subjected to high-resolution liquid chromatography and mass spectrometry (HR-LCMS) analysis, which was carried out at the

using the activity-guided repeated column chromatography (Table 1). Fig. 1 shows elution with solvents such as methanol and dichloromethane, yielding a bioactive compound in the petroleum ether in the form of acetone fraction (85:15 v/v). The eluted fractions were analyzed using TLC and HPLC,

showing 85% purity of the bioactive compound. Moreover, the resilient biological activities of the fraction led to structural elucidation by various spectroscopic methods (UV, IR,<sup>1</sup>HNMR,<sup>13</sup> CNMR, and MS). The elucidation of the structures was as follows:




Fig. 1. Separation scheme of p-Coumaric acid from methanol extract of *Diospyros melanoxylon* and its structure

Table 1. Antibacterial activity by disc diffusion for methanol extract of *Diospyros melanoxylon* leaves (DMM), diverseethyl acetate soluble fractions (ESF) and isolated compound p-Coumaric acid (PC)

	Zone of inhibition* (in “mm”)					
	Std.	DMM	ESF	ESF5	ESF5-c	PC
<b>Gram positive</b>						
<i>Bacillus cereus</i>	11.02 ± 0.09	16.05 ± 1.33	11.34 ± 1.04	09.64 ± 1.11	07.35 ± 0.13	05.67 ± 2.23
<i>Micrococcus luteus</i>	14.25 ± 0.06	10.13 ± 1.05	07.86 ± 0.36	07.19 ± 0.31	05.04 ± 0.03	03.19 ± 3.16
<i>Staphylococcus aureus</i>	18.67 ± 0.09	20.13 ± 0.86	18.43 ± 1.07	18.00 ± 2.22	14.61 ± 0.49	11.22 ± 0.33
<b>Gram negative</b>						
<i>Klebsiella pneumoniae</i>	11.08 ± 0.15	10.27 ± 1.17	10.01 ± 1.14	05.88 ± 0.61	5.67 ± 0.15	4.98 ± 0.15
<i>Enterobacter aerogenes</i>	15.25 ± 0.88	10.21 ± 0.91	08.84 ± 0.18	07.54 ± 0.28	06.56 ± 0.90	03.33 ± 0.25
<i>Escherichia coli</i>	32.04 ± 0.53	34.31 ± 1.78	33.16 ± 1.25	33.03 ± 0.18	28.83 ± 0.42	25.15 ± 1.17
<i>Pseudomonas fluorescens</i>	18.09 ± 0.31	08.51 ± 2.69	08.04 ± 2.03	07.66 ± 0.09	05.50 ± 3.10	03.17 ± 1.12
<i>Pseudomonas aeruginosa</i>	25.06 ± 0.80	10.06 ± 3.46	08.55 ± 0.10	05.65 ± 0.16	05.58 ± 0.25	03.95 ± 0.10
<i>Salmonella enteritidis</i>	18.40 ± 0.32	20.91 ± 1.55	20.81 ± 1.57	17.01 ± 0.44	16.24 ± 0.24	16.01 ± 0.17

\* Values are expressed as mean ± SE. (Std.): Amoxicillin; (-): inactive

**p-coumaric acid: (E)-3-(4-hydroxyphenyl) acrylic acid**

buff crystalline solid, melting point: 208-210 °C; IR (KBr): 1680 (C=C), 1,750 (COOH), 3,650 cm<sup>-1</sup> (phenolic-OH); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 4.5 (bs, 1H, Ar-OH), 6.5 (d, J=7.5 Hz, 1H, COCH), 6.8-7.2 (m, 4H, Ar-H), 7.4 (d, J=7 Hz,

1H, Ar-H);<sup>13</sup> C NMR (DMSO-d<sub>6</sub>): δ 115.4, 115.8 (2), 127.9 (3), 148.1, 157.6, 170.5; LC-MS: m/z 165 (M+1); analytical calculated data for C<sub>9</sub>H<sub>8</sub>O<sub>3</sub>(164): C, 65.85; H, 4.91; found: C, 65.89; H, 4.96%

The identity of the compound was deciphered based on the mentioned results and

after comparison with the NMR and MS data as reported in the literature.<sup>19-21</sup>

### Antibacterial activity

Tables 1-2 show the antibacterial activity (disc-diffusion) and MIC values exhibited by the DMM, diverse ESF, and p-coumaric acid isolated compound against nine food pathogens. Methanol was considered as the negative control and could not inhibit the growth of the tested bacterial strains. The tested samples revealed dissimilar antibacterial activity, and the potency of the antibacterial activity varied depending on the species of the bacterial strains, diverse ethyl acetate soluble fractions, methanol extract, and isolated compound. In general, DMM exhibited potent inhibitory activity compared to the diverse soluble fractions and isolated

compounds. Moreover, the antibacterial activity of MBI (1 mg/ml) was on par with the standard antibiotic activity of amoxicillin, which was used as the positive control.

The results obtained from the agar well-diffusion method were used to determine the MIC values (Table 2; Fig. 2. In terms of the MIC values (mg/ml), DMM evidently possessed potent inhibition capacity in case of the gram-positive bacteria (*Bacillus cereus* and *Staphylococcus aureus*) and gram-negative bacteria (*Escherichia coli* and *Salmonella enteritidis*), and the effect was more significant compared to amoxicillin ( $P<0.05$ ). In general, the MIC values of DMM, isolated p-coumaric acid, and all the tested fractions were within the range of 0.80-7.99 mg/ml.

Table 2. The minimum inhibitory concentration (MIC) for methanol extract of *Diospyros melanoxylon* leaves (DMM), diverse ethyl acetate soluble fractions (ESF) and isolated compound p-Coumaric acid (PC)

	MIC* (in "mg/ml")					
	Std.	DMM	ESF	ESF5	ESF5-c	PC
<b>Gram positive</b>						
<i>Bacillus cereus</i>	1.34 ± 0.61	1.05 ± 0.25	1.30 ± 1.10	1.77 ± 0.13	3.78 ± 0.65	3.85 ± 0.55
<i>Micrococcus luteus</i>	2.50 ± 0.26	2.67 ± 0.21	3.05 ± 1.08	4.50 ± 0.84	4.61 ± 3.44	6.28 ± 1.63
<i>Staphylococcus aureus</i>	3.10 ± 0.12	2.77 ± 2.47	2.98 ± 0.41	3.33 ± 2.19	3.25 ± 1.54	3.90 ± 1.88
<b>Gram negative</b>						
<i>Klebsiella pneumoniae</i>	4.05 ± 0.38	4.58 ± 1.89	4.98 ± 0.02	6.06 ± 1.14	7.02 ± 0.82	7.76 ± 1.40
<i>Enterobacter aerogenes</i>	1.75 ± 0.25	1.85 ± 0.43	2.01 ± 0.48	2.69 ± 0.20	4.44 ± 2.31	5.15 ± 5.65
<i>Escherichia coli</i>	1.00 ± 0.20	0.80 ± 0.35	0.94 ± 0.58	1.34 ± 2.27	1.47 ± 0.15	1.77 ± 0.38
<i>Pseudomonas fluorescens</i>	3.08 ± 0.95	4.74 ± 1.06	5.31 ± 2.32	6.59 ± 0.32	7.86 ± 1.24	7.99 ± 1.05
<i>Pseudomonas aeruginosa</i>	2.50 ± 0.04	3.61 ± 0.24	4.83 ± 0.40	5.55 ± 0.58	5.97 ± 0.46	6.56 ± 0.71
<i>Salmonella enteritidis</i>	1.95 ± 0.45	1.34 ± 0.79	1.55 ± 1.22	2.54 ± 1.11	2.64 ± 1.24	3.00 ± 2.50

\* Values are expressed as mean ± SE. (Std.): Amoxicillin


Previous studies have clearly demonstrated that the antibacterial effects of p-coumaric acid are exerted through the dual damage mechanism, which involves the increased permeability of the membrane, along with binding to the phosphate group of DNA.<sup>22</sup> In addition, some studies have reported that the derivatives of p-coumaric acid have higher antibacterial activity compared to the compound alone.<sup>23-25</sup> According to the results of the present study, the activity of the extract was more prominent compared to the compound alone, which suggested that the collective effects of various phytochemicals in the extract are responsible for its potent antibacterial

activity.


In the current research, the antibacterial activity of *D. melanoxylon* was recorded against *Klebsiella*, *Pseudomonas*, *E. coli*, and *S. aureus*, and the plant extract that was dissolved in bloodless and warm water showed variable antibacterial effects against in these bacterial pathogens. The leaf extract of *D. melanoxylon* changed *in-vitro* against *Klebsiella*, *Pseudomonas*, *E. coli*, and *S. aureus*. In addition, the exhibited inhibition zones (mm) were listed in the inhibition sector using the methanol herbal extract, becoming subtracted from the inhibition zone. The leaf extract of *D.*

water showed the maximum inhibition zone in case of *Klebsiella* (17 mm), *Pseudomonas s*


(16 mm), *E. coli* (15 mm), streptomycin (19 mm), and *S. aureus* (16 mm).




Inhibition of klebsiella by leaf extract with solvents (A) methanol; (B) methanolic extract ; (C) Streptomycin



Inhibition of Pseudomonas by leaf extract with solvents (A) methanol; (B) methanolic extract; (C) Streptomycin



Inhibition of E-coli by leaf extract with solvents (A) methanol; (B) Methanolic extract; (C) Streptomycin



Inhibition of Staphylococcus aureus by leaf extract with solvents (A) methanolic extract; (B) Methanol; (C) Streptomycin

Fig. 2. Antibacterial activities for methanol extract of *Diospyros melanoxylo*n leaves

**Antioxidant activity**

Table 3 shows the comparative radical scavenger activity of DMM, diverse ESF, and isolated p-coumaric acid in terms of the EC<sub>50</sub> values. Accordingly, the methanol extract of *D. melanoxylo*n showed maximum radical scavenging activity when tested with DPPH, ABTS<sup>+</sup>, and superoxide radical scavenging assays. Conversely, the diverse soluble fractions and isolated p-coumaric acid exhibited low activity (1.2-1.6 times lower than the methanol extract). Moreover, DMM was more effective than BHA (positive control) in the scavenging

of free (DPPH), cation (ABTS), and anion (superoxide) radicals. In general, the tested samples showed more significant inhibitory effects in the following order: DMM>BHA>ESF>ESF5>ESF5-c>p-coumaric acid.

Interestingly, the antioxidant activity of the DMM fraction was more significant, which also provided maximum phenolic compounds during the preliminary phytochemical screening. This is consistent with the previous studies in this regard, suggesting the higher antioxidant potential of phenolic acids. Although high



phenolic content does not necessarily indicate greater antioxidant potential, assessment of the individual components of herbal extracts could provide evidence on their antioxidant capacity.<sup>26</sup> Nevertheless, the observed effects in the present

study could be attributed to the redox potential of the phenolic acids found in the extract, rendering it a reducing agent and hydrogen donor.

Table 3. Total phenolic (TPC), flavonoid (TFC) proanthocyanidin (TCC) contents and antioxidant activity of methanol extract of *Diospyros melanoxylon* leaves (DMM), diverse ethyl acetate soluble fractions (ESF) and isolated compound p-Coumaric acid (PC)

Sample	TPC (mg GAE/g)	TFC (mg QE/g)	TCC (mg GAE/g)	EC <sub>50</sub> <sup>x,y</sup> (µg/ml)		
				Radical scavenging activities		
				DPPH	ABTS	Superoxide
DMM	234.15±0.05 <sup>d</sup>	96.12±0.14 <sup>d</sup>	157.15±0.80 <sup>d</sup>	30.90±0.21 <sup>a</sup>	25.15±0.89 <sup>a</sup>	64.02±0.17 <sup>a</sup>
ESF	165.23±1.00 <sup>c</sup>	42.23±0.55 <sup>c</sup>	86.53±0.25 <sup>c</sup>	37.16±2.14 <sup>c</sup>	30.53±0.48 <sup>b</sup>	79.09±0.05 <sup>c</sup>
ESF5	98.54±0.82 <sup>b</sup>	30.64±2.02 <sup>b</sup>	55.95±0.49 <sup>b</sup>	41.03±0.23 <sup>d</sup>	37.77±0.50 <sup>c</sup>	85.46±1.11 <sup>d</sup>
ESF5-c	66.45±0.20 <sup>a</sup>	19.19±0.75 <sup>a</sup>	30.04±0.02 <sup>a</sup>	49.30±0.22 <sup>e</sup>	45.00±0.19 <sup>d</sup>	92.70±2.02 <sup>e</sup>
PC	-	-	-	55.55±0.12 <sup>f</sup>	53.01±0.86 <sup>e</sup>	98.89±2.22 <sup>f</sup>
BHA	-	-	-	35.55±0.01 <sup>b</sup>	30.16±1.28 <sup>b</sup>	66.57±0.34 <sup>b</sup>

<sup>x</sup> Values are expressed as mean ± SE. Means in the same column with distinct superscripts are significantly different (p ≤ 0.05) as separated by Duncan multiple range test.

<sup>y</sup> The EC<sub>50</sub> value is defined as the effective concentration of the test samples to show 50% of antioxidant activity under assay conditions

Table 4. Correlation between EC<sub>50</sub> of radical scavenging activities and total phenolic, flavonoid, proanthocyanidin content of methanol extract of *Diospyros melanoxylon* leaves (DMM), diverse ethyl acetate soluble fractions (ESF) and isolated compound p-Coumaric acid (PC)

	Correlation (R)*		
	phenolic	flavonoid	proanthocyanidin
<b>DPPH</b>			
DMM	0.956	0.912	0.856
ESF	0.912	0.867	0.825
ESF5	0.825	0.756	0.741
ESF5-c	0.793	0.700	0.666
PC	0.702	0.654	0.614
<b>ABTS</b>			
DMM	0.889	0.875	0.875
ESF	0.826	0.854	0.814
ESF5	0.745	0.702	0.733
ESF5-c	0.667	0.643	0.694
PC	0.606	0.608	0.662
<b>Superoxide</b>			
DMM	0.812	0.713	0.785
ESF	0.756	0.695	0.745
ESF5	0.701	0.630	0.674
ESF5-c	0.645	0.542	0.620
PC	0.599	0.505	0.544

\* Values are expressed as mean ± SE

#### Correlation between antioxidant activity and TPC, TFC, and TCC

We performed the correlation analysis of TPC, TFC, and proanthocyanidin (TCC) contents with the EC<sub>50</sub> values of the radical

scavenging ability of the methanol extract of *D. melanoxylon*, isolated compound, and diverse soluble fractions. According to the findings, TPC, TFC, and TCC were significantly correlated (R<sup>2</sup>=0.505-0.956) with the DPPH, ABTS, and superoxide radical scavenging (Table 4). Several studies have confirmed that the presence of alkaloids and phenolic acids could enhance the antioxidant ability of herbal extracts.<sup>27-29</sup> Previous findings on *D. melanoxylon* have also confirmed the potent antioxidant activity of its bark extract.<sup>30, 31</sup> However, this was the first report regarding the methanol leaf extract of this plant, which exhibited promising antioxidant activity in line with the previous findings regarding the potential of phenols and flavonoids.

#### Anti-inflammatory activity

As part of the study on the mechanism of anti-inflammation activity, the ability of methanol extract of *D. melanoxylon*, isolated compound, and diverse soluble fractions to inhibit protein denaturation was assessed in the present study. According to the findings, the

extract could effectively inhibit albumin denaturation with the  $IC_{50}$  of 69  $\mu$ g/ml (Table 5). On the other hand, aspirin was considered as the standard anti-inflammatory drug (positive control) in our research, which could inhibit albumin denaturation with the  $IC_{50}$  of 80  $\mu$ g/ml. Additionally, p-coumaric acid ( $IC_{50}$ =119  $\mu$ g/ml) isolated from DMM was observed to be a potent inhibitor, while inhibition was slightly lower compared to DMM, diverse ESF, and aspirin. Albumin denaturation is an indicator of inflammation, which was assessed in the present

study in order to determine the anti-inflammatory potential of the herbal extract.

In the current research, the RBC membrane stabilization method was applied for DMM, diverse ESF, and the isolated compound in order to investigate the *in-vitro* anti-inflammatory activity as the stabilization of the lysosomal membrane prevents inflammation and further damage to the surrounding tissues by preventing the release of its enzymes. In the present study, DMM could effectively inhibit the heat-induced hemolysis with the  $IC_{50}$  of 80  $\mu$ g/ml (Table 5).

Table 5. Albumin denaturation, membrane protection/stabilization and proteinase inhibition potential of methanol extract of *Diospyros melanoxylon* leaves (DMM), diverse ethyl acetate soluble fractions (ESF) and isolated compound p-Coumaric acid (PC)

Anti-inflammatory	$IC_{50}^{x,y}$ ( $\mu$ g/ml)					
	DMM	ESF	ESF5	ESF5-c	PC	Aspirin <sup>#</sup>
Albumin denaturation	69.12±0.24 <sup>a</sup>	81.06±0.93 <sup>b</sup>	97.85±1.13 <sup>c</sup>	106.66±1.05 <sup>d</sup>	119.20±3.75 <sup>e</sup>	80.11±0.11 <sup>b</sup>
Membrane protection	80.21±2.52 <sup>a</sup>	89.95±1.01 <sup>b</sup>	96.65±0.24 <sup>c</sup>	133.32±0.71 <sup>d</sup>	144.41±0.27 <sup>e</sup>	90.80±0.40 <sup>b</sup>
Proteinase inhibition	91.19±1.36 <sup>a</sup>	101.00±0.05 <sup>c</sup>	137.52±1.28 <sup>d</sup>	143.56±0.27 <sup>e</sup>	149.95±0.28 <sup>f</sup>	100.42±0.17 <sup>b</sup>

<sup>x</sup> Values are expressed as mean  $\pm$  SE. Means in the same row with distinct superscripts are significantly different ( $p \leq 0.05$ ) as separated by Duncan multiple range test.

<sup>y</sup> The  $IC_{50}$  value is defined as the inhibitor concentration to inhibit 50% under assay conditions.

<sup>#</sup>Aspirin was used as positive control.

In the current research, DMM and ESF ( $IC_{50}$ =89  $\mu$ g/ml) exhibited slightly higher inhibition potential than the standard drug ( $IC_{50}$ =90  $\mu$ g/ml), while p-coumaric acid ( $IC_{50}$ =144  $\mu$ g/ml) had slightly lower inhibitory activity. In addition, proteinase was observed to be instrumental in the repair of tissue damage during inflammation. DMM ( $IC_{50}$ =91  $\mu$ g/ml) exhibited substantial anti-proteinase activity compared to the standard drug ( $IC_{50}$ =100  $\mu$ g/ml), while the inhibitory effects of proteinase based on the  $IC_{50}$  values of ESF ( $IC_{50}$ =137  $\mu$ g/ml), ESF 5 ( $IC_{50}$ =143  $\mu$ g/ml), and ESF5-c ( $IC_{50}$ =149  $\mu$ g/ml) were comparatively less significant ( $P < 0.05$ ) compared to the therapeutic drug aspirin ( $IC_{50}$ =100  $\mu$ g/ml). Therefore, it could be concluded that the anti-inflammatory

potential of the leaf extract of *D. melanoxylon* should be further evaluated in terms of the mechanism of action. These findings are consistent with the previous studies in this regard, suggesting that anti-inflammatory activity of the extract was directly correlated with its phytochemical constituents.<sup>32, 33</sup>

#### HR-LCMS profile study of the active methanol extract

The potent antibacterial, antioxidant, and anti-inflammatory effects of the methanol extract were investigated using the HR-LCMS analysis, which revealed approximately 20 compounds (Table 6), five of which were identified as phenolic acids (p-coumaric acid, homoveratric acid, chlorogenic acid, eudesmic acid, and tuberonic acid) and one flavonoid

(rutin). Furthermore, the HR-LCMS analysis of the methanol extract of *D. melanoxylon* leaves demonstrated nine and six principal peaks,

respectively, which confirmed the presence of diverse phytochemical materials.

Table 6. Chemical profile of the methanol extract of *Diospyros melanoxylon* leaves by HR – LCMS in +ESI mode

Compound detected	Molecular formula	DB Diff (ppm)
<b>+ESI mode</b>		
Chlorogenic acid	C <sub>16</sub> H <sub>18</sub> O <sub>9</sub>	9.21
7-[2-Trifluoromethyl-4- (2-Hydroxyphenyl)-1,3- Dioxan-cis-5-YL]-HEPT-5Zenoic acid	C <sub>18</sub> H <sub>21</sub> F <sub>3</sub> O <sub>5</sub>	25.02
Eudesmic acid	C <sub>10</sub> H <sub>12</sub> O <sub>5</sub>	7.72
Homoveratric acid	C <sub>10</sub> H <sub>12</sub> O <sub>4</sub>	8.08
Deacetyl-Nmonodemethylidiltiazem	C <sub>19</sub> H <sub>22</sub> N <sub>2</sub> O <sub>3</sub> S	14.37
12R-acetoxy-punaglandin 3	C <sub>27</sub> H <sub>35</sub> Cl O <sub>9</sub>	21.01
p-Coumaric acid	C <sub>9</sub> H <sub>8</sub> O <sub>3</sub>	8.10
Tuberonic acid	C <sub>12</sub> H <sub>18</sub> O <sub>4</sub>	9.23
Rutin	C <sub>27</sub> H <sub>30</sub> O <sub>16</sub>	9.89
5-Phenylvaleric acid	C <sub>11</sub> H <sub>14</sub> O <sub>2</sub>	8.41
<b>-ESI mode</b>		
Piceid	C <sub>21</sub> H <sub>24</sub> O <sub>8</sub>	15.80
3,9,15-Docosatriynoic acid	C <sub>22</sub> H <sub>32</sub> O <sub>2</sub>	15.64
4,7,10,13,16- docosapentaenoic acid	C <sub>22</sub> H <sub>34</sub> O <sub>2</sub>	15.44
7-[2-Trifluoromethyl-4- (2-Hydroxyphenyl)-1,3- Dioxan-cis-5-YL]-HEPT-5Zenoic acid	C <sub>18</sub> H <sub>21</sub> F <sub>3</sub> O <sub>5</sub>	6.67

In the assessment of the excessive resolution liquid chromatography and mass spectra of the constituents with the primary library, many of these compounds were characterized and identified, including dihydromyricetin, dihydrorobinetin, rutin, cosmosiin, barbituric acid, 5-ethyl-five-(2-hydroxyethyl), 2,2,nine,nine- tetramethyl-undecan-1,10-diol,sinomenine, dihydrodeoxy-streptomycin, hexadecanediocacid, thosuximide M5, hydroxy anastrozole, 7-desmethylpapaverine, lyxosylamine, isovaleric acid, taurine, minoxidil, 4-trimethyl aminobutanal, 6-beta naltrexol-3-glucuronide, and glucosylgalactosyl hydroxylysine.

Among the compounds detected in the extract were tannins, alkaloids, saponins, glycosides, flavonoids, and steroids using the simple phytochemical technique. The antibacterial activity was also terminated using poison techniques, while notable results have been observed in case of *Colletotrichum capsici*, while excellent results have been yielded toward *Fusarium oxysporum* compared to fungicides. In the current research, the plant extract play a pivotal role in controlling plant diseases, while

the phytochemical properties are also effective in the treatment of various diseases. Therefore, new herbal medicines could be proper therapeutic alternatives for chronic diseases.

#### Molecular modeling

As is depicted in Figs. 3-5, p-coumaric acid could bind to  $\beta$ -lactamase at the MEG binding site, interacting with the binding site residues ASN109, PHE91, PRO90, TYR89, TRP94, and LYS87. Furthermore, p-coumaric acid had good affinity toward the MEG binding site with the binding energy of -4.6 Kcal. It is also notable that p-coumaric acid acted as a competitive inhibitor of  $\beta$ -lactamase by binding to the MEG binding site, while preventing MEG as a natural substrate of  $\beta$ -lactamase, thereby inhibiting the action of the  $\beta$ -lactamase enzyme. The inhibition of  $\beta$ -lactamase limits the ability of the organism to degrade the penicillin class of antibiotics. Therefore, the antibiotics that are fortified with p-coumaric acid could contribute to the treatment of drug-resistant microorganisms through the dual damage effect exerted by the compound on the microbial membrane.




Fig. 3. Image showing p-Coumaric acid bound to  $\beta$ -Lactamase at MEG binding site




Fig. 4. Image showing interaction of p-Coumaric acid with the binding site residues ASN109, PHE91, PRO90, TYR89, TRP94 and LYS87




Fig. 5. Image showing hydrophobic interaction of p-Coumaric acid with MEG binding site residue of  $\beta$ -Lactamase

## Conclusion

According to the results, the analytical activity guided the characterization of the leaf extract of *D. melanoxylon*, confirming the

presence of various compounds, such as p-coumaric acid, homoveratric acid, chlorogenic acid, eudesmic acid, and tuberonic acid. Moreover, the antibacterial, antioxidant, and anti-inflammatory activity of the extract was more prominent compared to the isolated compound, which suggested the synergistic effects of the components of the extract to have higher potential as a therapeutic agent compared to the p-coumaric acid extract from the fraction. The results uphold the positive effects of the extract, highlighting the higher *in-vivo* potential of the extract to reconfirm its potential health benefits.

## References

1. Roxburgh W. Flora Indica (Eds Carey W and WallichN). Serampore. 2: 1820–1824.
2. Wallinofer B. The biology and systematics of Ebenaceae: A review. Ann Naturlist Mus Wien 2001; 103(2): 485–512.
3. Rathore JS. Diospyros melanoxylon, a breadwinner tree of India. Econ Bot 1972; 26(4): 333–339.
4. Palaparthi S. Pharmacological Profile of Diospyros melanoxylon Methanolic Extract. J Pharmacogn Nat Prod 2019; 5: 157.
5. Rath S K, Mohapatra N, Dubey D, Panda S K, Thatoi HN, Dutta SK. Antimicrobial activity of Diospyros melanoxylon bark from Similipal Biosphere Reserve, Orissa, India. Afr J Biotechnol 2009; 8 (9): 1924–1928.
6. El-Seedi HR, El-Said AM, Khalifa SA, Göransson U, Bohlin L, Borg-Karlson AK, et al. Biosynthesis, natural sources, dietary intake, pharmacokinetic properties, and biological activities of hydroxycinnamic acids. J Agric Food Chem 2012; 60:10877–10895.
7. Pragasaam SJ, Venkatesan V, Rasool M. Immunomodulatory and anti-inflammatory effect of p-coumaric acid, a common dietary polyphenol on experimental inflammation in rats. Inflammation 2013; 36: 169–176.
8. Torres y Torres J L, P.N Rosazza J. Microbial transformations of p-coumaric acid by Bacillus megaterium and Curvularia lunata. J Nat Prod 2001; 64: 1408–14.
9. Lou Z, Wang H, Rao S, Sun J, Ma C, Li J. p-Coumaric acid kills bacteria through dual damage mechanisms. Food Control 2012; 25(2): 550–554.
10. Harborne A J. Phytochemical methods: A guide to modern techniques of plant analysis. Chapman

- and Hall Ltd, London, Springer Netherlands 1998.
11. Kujala TS, Loponen JM, Klika KD, Pihlaja K. Phenolic and betacyanins in red beetroot (*Beta vulgaris*) root: Distribution and effects of cold storage on the content of total phenolics and three individual compounds. *J Agric Food Chem* 2000; 48: 5338–5342.
  12. Ordóñez AAL, Gomez JD, Vattuone MA & Isla MI. Antioxidant activities of *Sechium edule* (Jacq.) Swart extracts. *Food Chem* 2006; 97: 452–458.
  13. Sun BS, Ricardo-da-Silva JM, Spranger I. Quantification of catechins and proanthocyanidins in several Portuguese grapevine varieties and red wines. *Ciencia T ec Vitiv* 2001; 16:23–34.
  14. Nurhanani R, Rasyidah R, Sarni MJ, Azlina AA. Radical scavenging and reducing properties of extracts of cashew shoots (*Anacardium occidentale*). *Food Chem* 2008; 111: 38–44.
  15. Ramith Ramu, Prithvi S Shirahatti, Farhan Zameer, Dhananjaya B Lakkapa, MN Nagendra. Evaluation of Banana (*Musa sp. var. Nanjangud Rasa bale*) flower and pseudostem extracts on antimicrobial, cytotoxicity and thrombolytic activities. *Int J Pharm Pharm Sci* 2015; 7 (1): 136-140.
  16. Mizushima Y, Kobayashi M. Interaction of anti-inflammatory drugs with serum preteins, especially with some biologically active proteins. *J Pharm Pharmacol* 1968; 20(3): 169–73.
  17. Rabia Naz, Hafsa Ayub, Sajid Nawaz, Zia Ul Islam, Tayyaba Yasmin, Asghari Bano, et al. Antimicrobial activity, toxicity and anti-inflammatory potential of methanolic extracts of four ethnomedicinal plant species from Punjab, Pakistan. *BMC Complement Altern Med* 2017; 17(1): 302.
  18. Oyedepo OO, Femurewa AJ. Anti-protease and membrane stabilizing activities of extracts of *Fagrazanthoxiloides*, *Olax subscorpioides* and *Tetrapleuratetraptera*. *Int J Pharm Cogn* 1995; 33: 65–9.
  19. Katsuko Inamatsu. Isolation of P-coumaric acid and sterols from rice straw compost. *Soil Sci Plant Nutr* 1968; 14(2): 62-67.
  20. Elgizawy H, Hussein M A. Isolation, Structure Elucidation of Ferulic and Coumaric acids from *Fortunella japonica* Swingle leaves and their Structure Antioxidant activity relationship. *Free Radic Antioxidant* 2017; 7(1): 23-30.
  21. Feng Guo, Shuoyang Zhang, Xiuying Yan, Yuhan Dan, Jian Wang, Yunli Zhao, et al. Bioassay-guided isolation of antioxidant and  $\alpha$ -glucosidase inhibitory constituents from stem of *Vigna angularis*. *Bioorganic Chem* 2019; 87: 312-320.
  22. Zhao Y, Chen MS, Zhao ZG, Yu SJ. The antibiotic activity and mechanisms of sugarcane (*Saccharum officinarum* L.) bagasse extract against food-borne pathogens. *Food Chem* 2015; 185: 112–118.
  23. Pereira JA, Oliveira I, Sousa A, Valentao P, Andrade PB, Ferreira ICFR. et al. Walnut (*Juglans regia* L.) leaves: phenolic compounds, antibacterial activity and antioxidant potential of different cultivars. *Food Chem Toxicol* 2007; 45(11): 2287–2295.
  24. Madikizela B, Aderogba MA, Finnie JF, Van Staden J. Isolation and characterization of antimicrobial compounds from *Terminalia phanerophlebia* Engl. & Diels leaf extracts. *J Ethnopharmacol* 2014; 156: 228–234.
  25. Tawata S, Taira S, Kobamoto N, Zhu J, Ishihara M, Toyama S. Synthesis and antifungal activity of cinnamic acid esters. *Biosci Biotechnol Biochem* 1996; 60(5): 909–910.
  26. Rice Evans CA, Miller NJ, Paganga G. Structure-antioxidant activity relationships of flavonoids and phenolic acids. *Free Radical Biol Med* 1996; 20(7): 933-56.
  27. Pu F, Ren XL, Zhang XP. Phenolic compounds and antioxidant activity in fruits of six *Diospyros kaki* genotypes. *Eur Food Res Technol* 2013; 237(16): 923–932.
  28. Casselst BK, Asencio M, Conget P, Speisky H, Videla LA, Lissi EA. Structure-antioxidative activity relationships in benzyloisoquinoline alkaloids. *Pharmacol Res* 1995; 31(2): 103–107.
  29. Yin WQ, Duan SQ, Zhang Y, Zeng L, Song XM. Antioxidant activities of different solvents extracts and alkaloids of *Uncaria hynchophylla* (Miq.) Jacks. *J Guangxi Normal University* 2010; 28(1): 31–34.
  30. Al Rashid MH, P.v.p DB, Majumder S, Mandal V, Pal M, Mandal SC, et al. Antioxidant and anticancer activity of extract and fractions obtained from *Diospyros Melanoxylon* Roxb. leaves and correlation with their polyphenolic profiles. *Int J Pharm Pharm Sci* 2018; 10(11): 6-16.
  31. Hmar BZ, Mishra S, Pradhan RC. Physico-Chemical, mechanical and antioxidant properties of Kendu (*Diospyros Melanoxylon* Roxb.). *Curr*

- Res Nutr Food Sci 2017; 5 (3):214-222.
32. Bhuvana P, Anuradha V, Syed Ali M, Suganya V, Sangeetha P. Drug standardization and invitro anti-inflammatory and antibacterial activity of Chlorella vulgaris. Int J Pharm Biol Sci 2018; 8 (1): 103-110.
  33. Bigoniya P, Rana AC. Protective Effect of Wrightiatinctoria Bark Triterpenoid 342 Fraction on Carbon Tetrachloride Induced Acute Rat Liver Toxicity. Iran J Pharm Ther 2010; 9(2): 55-62.