



A review on effect of air pollutants on fetal development and pregnancy outcomes

Rekha Kashyap, Rajnish Gupta*, Mahamad Asif Mansuri, Rupal Thasale, Zarana Rathod, Ravi Patni

Air Pollution Division, ICMR-National Institute of Occupational Health, Meghani Nagar, Ahmedabad-016, Gujarat, India

ARTICLE INFORMATION

Article Chronology:

Received 02 July 2019

Revised 14 August 2019

Accepted 21 September 2019

Published 29 September 2019

Keywords:

Fetal development; Maternal health;
Prenatal exposure; Reproductive health

ABSTRACT:

Developing fetus is connected to the mother by placenta to receive oxygen and nutrients. Maternal exposure to ambient air pollution not only affects maternal health but also provides a pathway for many toxic pollutants to cross the placental barrier and interrupt biochemical milieu. There are numerous scientific studies available describing possible negative health impact of air pollutants on reproductive health, pregnancy outcomes and fetal development yet no toxic effect is available recently. Studying the pollutants exposure effect on fetal development is crucial for the underlying mechanism between prenatal exposure and pregnancy outcomes. Present review meticulously provides the compiled data of 40 most recent studies with possible action mechanism of air pollutants on pregnancy outcomes and fetal development to find a better solution to exterminate or reduce the problem.

CORRESPONDING AUTHOR:

rajnish_gupta1985@hotmail.com

Tel: (+91 79) 22688866

Fax: (+91 79) 22688866

Review

Air pollution in the present scenario is considered the major inducer of harmful health effects, especially by emitting harmful gases (SO_2 , NO_x , NH_3 , CO , and CO_2), polycyclic aromatic hydrocarbons (Naphthalene, Anthracene, Phenanthrene, Fluoranthene, Benzo(b)fluoranthene, Benzo(k)fluoranthene, Benzo(a)pyrene, Dibenzo(a,h)anthracene, and Indeno(1, 2, 3-cd) pyrene), Volatile Organic Compounds (Benzene, Toluene, *o-m-p* Xylene, Ethylbenzene etc.), Particulate Matter (PM_{10}

and $\text{PM}_{2.5}$, $\text{PM}_{0.1}$) as well as combustion-derived nanoparticles (both Petrol Exhaust Nanoparticles, PENPs and Diesel Exhaust Nanoparticles, DENPs) [1] causing not only environmental issues [2] but also cause health hazard *via* blood circulation, inhalation and cause asthma, pharyngeal and laryngeal cancer, tuberculosis, nutritional deficit, cardiovascular and other endocrine disorders, adverse pregnancy outcomes, neonatal abnormality and mortality too [1, 3].

Pregnancy outcomes are important indicators

of the maternal as well as newborn and infant health. Stillbirth, low birth weight (LBW) and impaired growth directly reflect the abnormal maternal health [4]. In the last few years, there is an increasing research trend towards exploring the possible adverse effects of ambient air pollution on maternal health and birth outcomes. Previous attempts were made to critically review the correlation between air pollutant exposure and

maternal health but the problem is increasing exponentially day by day and there is no recent data is available to scientifically validate the same and as the problem is increasing very speedily there is an urgent need to monitor the risky consequences of pollutant exposure. This review compiled the data of 40 recent most population-based pregnancies and fetal growth studies worldwide (Table 1).

Table 1. Effect of different air pollutants on pregnancy and fetal outcome

S. No.	Objective of study	Pollutant name	Effect	References
1	Traffic-related air pollution (TRAP) and childhood obesity	NO _x	Increase in prenatal non-freeway NO _x was associated with 33% higher leptin and 9% high molecular weight adiponectin levels in cord blood. Leptin levels were found 71% higher in mothers who resided near to major roadways (75 m). Increased leptin level and altered infant weight was related to only female infants not males	[5]
2	Continuous 24-48 hours exposures of PM _{2.5} and other air toxicants on maternal birth weight, child acute respiratory infections as well as adult chronic respiratory and lung impairments	PM _{2.5} and air toxicants	The association between PM _{2.5} and air toxicants exposure, maternal birth weight, child acute respiratory infections as well as adult chronic respiratory and lung impairments were examined. Peripheral and cord blood samples were also collected to investigate gene-environment interactions	[6]
3	24 h household PM _{2.5} exposure during pregnancy	PM _{2.5}	PM _{2.5} exposure in pregnancy was associated with 4 g decrease in birth weight while 2% increase in its prevalence.	[7]
4	Association between prenatal exposure to PM _{2.5} and its constituents and preterm delivery	PM _{2.5} and PM ₁₀ , ammonium, chlorine, sodium, nitrate, vanadium and bromine	Increased risk of preterm delivery with full gestational exposure to PM _{2.5} and its constituents; preterm delivery were observed for blacks, Asians and older mothers	[8]
5	Impacts of air pollution on small gestational age (SGA) birth weight, LBW and preterm birth	CO, NO ₂ , NO, O ₃ , SO ₂ , PM _{2.5} and PM ₁₀	Exposure to all air pollutants was associated with SGA; proximity to highways was related with 26% increase in SGA and 11% increase in LBW. PM _{2.5} exposure with preterm births was observed at <37 weeks and for other pollutants at <30 weeks of gestation	[9]
6	Exposure to environmental contaminants during pregnancy	-	Environmental contaminants' exposure during pregnancy resulted in deregulation of genetic and epigenetic mechanisms of prenatal growth. Alteration of fetal development signaling led to various transgenerational disorders; necrotic, apoptotic and inflammatory reactions to fetal nucleic acids	[10]

7	Effect of high ambient air pollution exposure on low fertility rate and early pregnancy loss	PM ₁₀ , NO ₂ , SO ₂ , O ₃ and CO	Cumulative pregnancy rate in multiple IVF cycles was 51.3% per person. Air pollution exposure during periods 1 (start of COS to oocyte retrieval) and 3 (embryo transfer to hCG test) was generally associated with IVF outcomes. Increased NO ₂ and CO during period 1 were associated with decreased intrauterine pregnancy probability. PM ₁₀ , NO ₂ and CO levels during period 3 were also inversely related to intrauterine pregnancy.	[11]
8	Exposure to air pollutants and inhalable environmental elements during pregnancy and after birth and childhood-onset systemic lupus erythematosus (cSLE) diagnosis	PM ₁₀ , SO ₂ , NO ₂ , O ₃ and CO	An increased frequency of maternal occupational exposure during pregnancy to volatile vapor and smoking (maternal and/or secondhand) was related to cSLE development.	[12]
9	Relation of prenatal exposure to air pollutants, smoking with pregnancy complications and adverse birth outcomes	NO _x	NO ₂ and CALINE4-derived NO _x exposure increased the risk of high uterine artery resistance in later pregnancies. Smoking also increased the risk of higher uterine resistance and contributed to bilateral notching in mid-pregnancy	[13]
10	Placental biomarkers of PAH exposure and glutathione-S-transferase biotransformation enzymes in obstetric population	PAHs and Pb	Maternal serum folate levels were inversely related to PAHs-DNA adducts. Maternal Pb exposure was significantly correlated with cord-blood Pb levels. Maternal serum folate levels were negative predictor of total DNA-PAH adducts levels in placenta.	[14]
11	Associations between cause-specific stillbirth and prenatal exposure to air chemical pollutants and PM _{2.5}	Air chemical pollutants and PM _{2.5}	Stillbirths due to impaired fetal growth were increased following gestational exposure to PM _{2.5}	[15]
12	Residential pollutant exposure and pregnancy outcomes	NO _x , O ₃ , PM _{2.5} and PM ₁₀	Traffic exhaust were related to 2-6% increase in low birth weight, 1-3% increase in small gestational age affecting fetal growth	[16]
13	Pollutant exposure risk for neonatal development, intrauterine growth retardation, preterm birth, low birth weight, reduced birth size and congenital malformation	CO, NO ₂ , PM ₁₀ and PM _{2.5}	Exposure-mediated pregnancy complication is associated with reduced birth weight and increased probabilities of low birth weight, chromosomal aberrations, decreased neurological development as well as high risk of chronic diseases	[16,17]
14	Effect of estimated daily air pollutants exposure, temperature and greenness on newborns	-	Higher environmental pollutants exposure led to poorer fetal birth-weight, head-circumference and gestational age at birth	[18]
15	Association between air pollution exposure and birth weight	NO ₂ and PM ₁₀	Non-significant variations were observed in birth weight	[19]
16	Exposure effect of PM _{2.5} on preterm birth	PM _{2.5}	8.3% preterm births were observed during the 3 rd trimester	[20]

17	To identify association of pregnancy loss with common pollutant exposure	PM _{2.5} , O ₃ and sulfate compounds	Increased incidence of pregnancy loss was 28% were identified; increase in average whole pregnancy ozone and PM _{2.5}	[21]
18	Associations between air pollution and various birth outcomes	NO ₂ and PM _{2.5}	PM _{2.5} was not significantly associated with spontaneous preterm birth. NO ₂ in second trimester was negatively associated with spontaneous preterm birth. Spatial exposures brought about significant reduction in OR of PM _{2.5} and NO ₂	[22]
19	Evaluation of mother-infant pairs using spatio-temporal models and stationary monitors for exposure to ambient air pollution	PM _{2.5}	Increased risk of preterm birth and reduced fetal growth in an urban area. PM _{2.5} increase during pregnancy was associated with preterm birth. Increased lower birth weight, an IQR increase in modeled and monitored PM _{2.5} was 12.1 g and 15.9 g	[23]
20	Air pollution effects in pregnancy	-	Adverse effect of air pollution on respiratory and cardiovascular systems, also associated with type 2 diabetes; Embryonic and fetal organs are vulnerable to air pollutants mediated damage; increased risk of negative gestational outcomes, preterm delivery and low fetus birth weight	[24]
21	Correlation between household cooking fuels and risk of stillbirth	Household cooking fuels	Characteristic risk for stillbirths of firewood as cooking fuel in India was 11% and 1% for kerosene cooking. Women who cook with firewood, risk factors were significantly associated with occurrence of stillbirth	[25]
22	Air pollutant exposure during pregnancy	PM _{2.5} , NO ₂ and O ₃	Exposure was responsible for maternal asthma, gestational diabetes, heart disease, hypertension, low birth weight, pre-existing diabetes, preeclampsia, preterm birth, and small gestational age	[26]
23	Exposure to high pollution fuels	-	Use of high pollution fuels resulted 92 g reduction in mean neonatal birth weight as compared to low pollution fuel. Full adjusted OR for below average birth size risk was 1.29	[23]
24	Pollutant exposure during pregnancy and effect on intrauterine environment and fetal heart rate	-	Change in fetal heart rate, increased cesarean sections, neonatal asphyxia, cerebral palsy and even fetal death during delivery following pollutant exposure	[27]
25	To evaluate secondary health data for neonatal mortality rate (NMR) following exposure to cow dung/crop residues/firewood as fuel	Cow dung/crop residues/firewood	Mean rural NMR was 4.2% live births and urban NMR was 3.31%. Household air pollution (HAP) was 92.2% in rural areas as compared to 40.8% in urban areas. HAP was strongly associated with NMR for urban and rural areas combined.	[28]
26	Association between maternal exposures to environmental SO ₂ and total suspended particulates and risk of low birth weight	SO ₂ and total suspended particulates	Exposures to SO ₂ and total suspended particulates yielded adjOR between VLBW and maternal exposures to high levels of air pollution	[29]
27	Air pollution exposure during pregnancy; maternal asthma and neonatal respiratory outcomes	PM ₁₀ , PM _{2.5} , SO ₂ , NO ₂ and CO	Transient tachypnea of the newborn (TTN), asphyxia, and respiratory distress syndrome risk were increased during preconception and I trimester after PM ₁₀ , CO, NO _x , O ₃ and SO ₂ exposure	[30]

28	Pregnancy trimester-specific exposure to ambient air pollutants and respiratory health outcomes in the first two years of life	CO and PM _{2.5}	Trimester-specific air pollution modified the BMI and respiratory health in growing up; children of first trimester PM _{2.5} exposed overweight/obese mothers had an adjusted incidence rate ratio 1.85, 1.76 and 1.90 in quartile 2–4 for bronchiolitis/bronchitis (term ii), ear infection in the first year of life	[31]
29	Relationship between PM _{2.5} exposure during pregnancy and post-neonatal mortality	PM _{2.5}	PM _{2.5} exposure increased the risk factor for total mortality, respiratory syndrome and sudden infant death syndrome (SIDS) mortality	[32]
30	To evaluate adverse pregnancy outcome, infant respiratory and circulatory problems	PM ₁₀ and PM _{2.5}	Maternal serum, urine as well as umbilical cord blood, tissue and placental tissue analyze using 8-hydroxydeoxyguanosine (8-OHdG), a biological indicator of oxidative stress	[33]
31	<i>In utero</i> particulate matter exposure in adulthood	PM ₁₀ and PM _{2.5}	<i>In utero</i> PM _{2.5} exposure caused acute inflammation, chronic matrix remodeling, as well as alterations of Ca ²⁺ handling proteins, resulting in cardiac dysfunction	[34]
32	Association between exposure to indoor biomass fuel sources, second-hand tobacco smoke and adverse health outcomes in early infancy	Wood and/or dung as primary indoor biomass fuel sources	Exposure to wood and/or dung as primary fuel source was associated with 49% increase in LBW risk, 34% increase in respiratory illness incidence and 21% increase in six-month infant mortality risk. Exposed neonatal also had 45% and 30% respective increase in risks of underweight and stunting.	[35]
33	Exposure effect of PM and other ambient pollutants on 200 pregnancy outcomes	PM and other ambient pollutants	Maternal biological samples i.e. hair samples, urine samples (for 8-oxo-deoxyguanosine), blood samples for transcript markers and biomarkers of pre-eclampsia as well as neonatal blood samples for transcript markers were analyzed as predictors for pregnant women and neonatal at risk	[36]
34	Maternal exposure to PM ₁₀ in association with birth weight, fetus growth, shape and gestational age at birth	PM ₁₀	PM ₁₀ exposure caused significant lower birth weight (39.0g, preterm births) and (24.0g, term births); reduction in fetal growth among term and moderately preterm births were also observed	[37]
35	To compare the effect of traffic-related air pollution exposure and preeclampsia, preterm birth and very preterm birth	CO, NO, NO ₂ , NO _x and PM ₁₀ and PM _{2.5}	Pollutant exposure elevated risks for preeclampsia, preterm birth, and very preterm birth from maternal exposures. Increased risk of preterm birth and very preterm birth were also positively associated with PM ₁₀ and PM _{2.5} exposure	[38]
36	PM _{2.5} exposure in pregnant females and adverse birth outcome using a gap-filled satellite	PM _{2.5}	PM _{2.5} exposure was associated with a 12.85 g decrease in term birth weight, increased risk of preterm birth and term LBW.	[39]
37	Effects of prenatal and postnatal exposure to PM on Kawasaki disease (KD)	Suspended particulate matter	The Odd ratios of PM exposure were 1.59 for prenatal exposure and 1.41 for postnatal exposure suggestive of an increased risk of Kawasaki disease	[40]
38	Exposure effect of traffic air pollutants during pre-pregnancy or early pregnancy and birth defects	NO ₂ , PM ₁₀ and PM _{2.5}	Exposure to high level of NO ₂ and PM ₁₀ increased the risk on birth defects (P<0.05) during pre-pregnancy or early pregnancy	[41]

39	Air pollution exposure during pregnancy and risk assessment for next generation	PM _{2.5}	Increased offspring child systolic BP percentile and risk for elevated BP were observed following PM _{2.5} concentration at $\geq 13 \mu\text{g}/\text{m}^3$ during third-trimester (4.85 percentile increase in child SBP).	[42]
40	Air pollutant exposure during three months preconception, gestational weeks 3-8 and orofacial defects	PM ₁₀ , PM _{2.5} , CO, SO ₂ and NO _x	Preconception, CO and PM ₁₀ were significantly associated with cleft palate, while SO ₂ was associated with cleft palate \pm without cleft palate; NO _x and PM _{2.5} were also linked with risk.	[43]

Inclusion Criteria

Original, full-text articles, clinical studies, and reviews were considered in the study. Studies were included if women were exposed with environmental toxic pollutants during pregnancy (e.g., maternal age and gestational age (preterm and very preterm birth), pre-eclampsia, season of birth, birth weight and size, head circumference). Relationship between toxic environmental pollutants (particulate matter, SO₂, NO_x, O₃, volatile organic compounds, polycyclic aromatic hydrocarbons and HCHO) released from vehicular emission, industrial processes, tobacco smoke/e-cigarette, incense burning and frequency of cesarean sections (C-sections), preterm birth, neonatal asphyxia, cerebral palsy and fetal death were studied.

Exclusion Criteria

Studies which were not in English language OR of irrelevant population OR Irrelevant exposures OR irrelevant outcomes OR of patients with previously reported risk factors were excluded in the study.

Therefore, from 108 records identified through database synthesis, we included 40 studies in the qualitative synthesis.

Air pollutants induced cellular toxicity, DNA damage, and apoptosis

Carbon monoxide (CO)

CO toxicity occurs due to the synergistic effect of tissue hypoxia as well as direct CO-mediated cell

damage [44]. The affinity of CO to hemoglobin is approximately 210 times more than O₂. Exposure to the small environmental concentration of CO may form carboxyhemoglobin and compromises cellular oxygen availability and subsequent lethality. CO poisoning has been reported to develop long-term neurological effects as CO perturb the mitochondrial cytochrome-*c*-oxidase or Complex IV by binding to prosthetic group heme a₃, resulting in shut down of oxidative phosphorylation, similarly to cyanide and nitric oxide and causing abnormal ATP synthesis rate and cell signaling [45-50].

Ozone (O₃)

O₃ is a powerful oxidant, invades an unpaired electron of polyunsaturated fatty acids of the cell membrane to form ozonides and unstable zwitterions or trioxolane which further generate lipohydroperoxides, aldehydes, and H₂O₂. This pathway commences generation of lipid radicals and auto-oxidation of membranes and further causes DNA damage leading to impaired cellular function [51]. High-level ozone (O₃) exposure is a possible risk factor for late Alzheimer's disease onset [52]. Ambient ozone and fine particulate matter exposure is a risk factor for developing asthma, chronic obstructive pulmonary disease, atherosclerosis and diabetes [53, 54]. Ambient O₃ exposure is directly associated with oxidative-damage, diminished antioxidant defence and pro-inflammatory response in skin [55].

Sulfur dioxide (SO₂)

SO₂, a sensory irritant mainly penetrates the upper airways and can cause oxidative stress and DNA damage in multiple organs (brain, heart, kidney, liver, spleen, and testis) including lungs by disintegrating cell membrane [56-58].

Nitrogen oxide (NO_x)

Exposure to elevated NO_x leads to inflammation of the mucous membrane and rigidity of respiratory tracts. Exposure has reported to adversely affect natural killer cells and T-lymphocytes (mainly CD8⁺ determinant cells) causing weakened adaptive immunity against viral infection [59].

Particulate Matter (PM)

Nuclear membrane translocation of PM though is difficult to damage nuclear DNA due to size-specific nuclear transportation; by this non-oxidative mechanism several aberrant DNA adducts are generated. Lesion generated by polycyclic aromatic hydrocarbons (PAHs) and oxidative stress cause oxidative DNA damage, strand breaks (SB) and cytogenetic markers [60, 61] directly associated with increased inflammatory responses (GM-CSF, IL-6, IL-8, TNF- α), lipid peroxidation rate and alterations in cellular metabolism accompanied by activation of phospholipase A2, DNase II, the caspase-activated DNase (CAD) [62], endonuclease G (EndoG) [63], apoptosis-inducing factor (AIF) [64] leading to cell death [65-69].

Oxidative stress and ROS production following pollutant exposure

Oxidative stress is referred to as a disproportion between the appearance of ROS in the biological system and system's capacity to completely balance redox potential, repair the damage and/or detoxify the reactive intermediates. The repair

mechanism includes decreased oxidant species production or increased free radical scavenger levels (vitamin C or glutathione) or antioxidant enzymes *i.e.* catalase (CAT), glutathione peroxidase (GPx) and superoxide dismutase (SOD). Air Pollutants are responsible for ROS production and degenerating intracellular compartment as well as the fluid of respiratory tract, epithelial lining fluid, cytochrome P₄₅₀, cytoplasm, and mitochondria. Pollutants exposure is associated with oxidative damaged deoxynucleotides in blood and urine. Increased physicochemical surface properties of PM causes increased ROS-oxidative stress. High surface area to volume ratio of PM provide more surface for transition metals or organic compounds to get attach and causing aberration mitochondrial signaling mediators and enzymes, misbalanced intracellular Ca²⁺ homeostasis and activation of activator protein-1 (AP-1), mitogen-activated protein kinases (MAPK) and nuclear factor κ B (NF- κ B), upregulation of pro-inflammatory gene expression and cytokines [70]. Acute pollutant exposure diminishes the anti-oxidative status either by depleting the free radical scavengers or by decreasing antioxidant enzymes activity [71-74].

Effect of pollutants on pregnancy and fetal growth

International scenario

High ambient air pollution exposure was found to be linked with a low fertility rate and early pregnancy loss in women. Increased exposure to PM₁₀, CO, and NO₂ during Controlled Ovarian Stimulation (COS) and after embryo transfer is inversely proportional to decreased intrauterine pregnancy. 6621 cycles of 4581 patients with one or more fresh IVF cycles were analyzed and found cumulative pregnancy in multiple IVF was 51.3% per person. To evaluate patients' individual exposure to air pollution four exposure peri-

ods were measured including: 1): start of COS to oocyte retrieval, 2): oocyte retrieval to embryo transfer, 3): embryo transfer to hCG test and 4): start of COS to hCG test. Cox-proportional hazard ratios (HRs) were evaluated probabilities of pregnancy loss and intrauterine pregnancy for an IQR increase in each air pollutant concentration during each period. Increased NO_2 and CO during period 1 were associated with decreased intrauterine pregnancy. PM_{10} , NO_2 and CO levels during period 3 were negatively linked with intrauterine pregnancy. Both PM_{10} and NO_2 exposure during period 3 showed pregnancy loss [11]. Associations between cause-specific stillbirth and prenatal exposure to air chemical pollutants and $\text{PM}_{2.5}$ were quantified using Positive Matrix Factorization by Ebisu et al. 2018 on nested case-control study design (n=32262). The odds ratio per interquartile range for stillbirths due to impaired fetal growth was increased following gestational exposure to $\text{PM}_{2.5}$ though; no association was established between stillbirths caused by maternal complications and pollutant exposure [15]. To explore the exposure effect of $\text{PM}_{2.5}$ on preterm birth in China 426246 pregnant women were enrolled for birth outcome study in National free pre-pregnancy checkups project. The gestational exposure to $\text{PM}_{2.5}$ was alienated into the first trimester, second trimester, third trimester and entire pregnancy. Cox proportional hazards regression was used for evaluation of exposure effect by considering maternal age, alcohol use, education level, occupation, second-hand smoking, pre-pregnancy BMI, number of previous pregnancies, coastal areas and season of conception. The study found that 35261 (8.3%) preterm births were most significant during the third trimester and $\text{PM}_{2.5}$ exposure at $10 \mu\text{g}/\text{m}^3$ was significant during the first trimester and second trimester. Subgroup analysis revealed that women

who were having age more than 30 years, had low education level, worked as planters, had the male baby, had previous pregnancies, not living in coastal areas and conceived baby in winter was more sensitive to $\text{PM}_{2.5}$ exposure [20].

Exposure effect of air pollutants on the timing of incident pregnancy loss (from ovulation) was evaluated in 343 singleton pregnancies in a multisite prospective cohort study. The total evaluated pregnancy loss was 98 out of 343. Adjusted Cox proportional hazards models revealed an interquartile range increase in average pregnancy, ozone (HR-1.12), $\text{PM}_{2.5}$ (HR-1.13) and sulfate compounds (HR-1.58) exposure caused faster pregnancy loss. It was evidenced that pollutant exposure throughout pregnancy was associated with loss, but outlining specific heightened vulnerability period provides larger preconception [21].

Mother-infant pairs (n=61640) were evaluated using spatial-temporal models and stationary monitors for exposure to ambient air pollution and found associated with a higher risk of preterm birth and reduced fetal growth in an urban area. For average $\text{PM}_{2.5}$ increased IQR ($2.5 \mu\text{g}/\text{m}^3$) during pregnancy was associated with 1.04 ORs of preterm birth. For lower birth weight, an IQR increase in modeled and monitored $\text{PM}_{2.5}$ was 12.1 g and 15.9 g [23].

Exposure to $\text{PM}_{2.5}$, NO_2 , and O_3 in 818400 singleton live births during pregnancy was evaluated and was found that exposure was responsible for modification of maternal asthma, gestational diabetes, heart disease, hypertension, pre-existing diabetes, pre-eclampsia, preterm birth, low birth weight, and small gestational age. Interquartile range increases in $\text{PM}_{2.5}$, NO_2 , and O_3 during entire pregnancy were associated with 4%, 8.4% and 2% increase in the OR of preterm birth. For pregnancy exposure to $\text{PM}_{2.5}$ and NO_2 among

women with pre-existing diabetes, increases of 10.6% and 23.8% in the OR of preterm birth were detected while the increases of 3.8% and 6.5% ($P_{\text{int}} < 0.01$) were observed among women without this condition. For exposure to $\text{PM}_{2.5}$ during pregnancy, increase in OR of preterm birth was 8.3% more among women with pre-eclampsia than among women without preeclampsia (3.6%) ($P_{\text{int}} = 0.04$). Exposure to O_3 during pregnancy among asthmatic women caused a stronger increase in the OR of preterm birth (12.0%) as compared to non-asthmatic women (2.0%) [26]. Pollutant exposure during pregnancy affects the intrauterine milieu further causing a change in fetal heart rate (FHR) resulting in increased cesarean sections (C-sections) [27]. The incidence of neonatal asphyxia, cerebral palsy and even fetal death during delivery is also attributed to increased pollutant contact.

Son et al. 2017 investigated the relationship between $\text{PM}_{2.5}$ exposure during pregnancy and post-neonatal mortality using model adjusted by sex, birth weight, gestational length, maternal characteristics, temperature and season of birth and relative humidity. The study included extended Cox proportional hazards modeling with duration-dependent covariates in 465682 births with 385 mortalities. Neonatal exposure to $\text{PM}_{2.5}$ was measured from birth to death. Risk factor for total mortality, respiratory syndrome and Sudden Infant Death Syndrome (SIDS) mortality per-interquartile-range increase were 2.66, 3.14 and 2.50 respectively [32].

To evaluate adverse pregnancy outcomes and infant respiratory and circulatory problems following PM_{10} and $\text{PM}_{2.5}$ exposures, 2500 pregnant women with their infants were studied using 8-hydroxydeoxyguanosine (8-OHdG), a biological indicator of oxidative stress. Maternal serum, urine, umbilical cord blood and tissue and pla-

cental tissue were analyzed during all three trimesters [33].

To compare the effect of traffic-related air pollution exposure and pre-eclampsia, preterm birth (gestational age <37 weeks) and very preterm birth (<30 weeks) 81186 singleton births were studied and found at elevated risks from maternal exposure to traffic air pollutants (CO , NO , NO_2 , NO_x and $\text{PM}_{2.5}$ and PM_{10}) [38].

To understand the exposure effect of traffic air pollutants including NO_2 , PM_{10} and $\text{PM}_{2.5}$ during pre-pregnancy or early pregnancy and birth defects, a case-crossover study involving 4235 pregnant women were employed. The average ambient NO_2 , $\text{PM}_{2.5}$, and PM_{10} respective concentrations appeared as $60.83 \mu\text{g}/\text{m}^3$, $103.88 \mu\text{g}/\text{m}^3$ and $104.94 \mu\text{g}/\text{m}^3$. It was observed that exposure to a high level of NO_2 and PM_{10} increased the risk of birth defects ($P < 0.05$) during pre-pregnancy or early pregnancy [41].

Pollutant exposure during pregnancy is considered as a potential risk issue for neonatal development, intrauterine growth retardation, preterm birth, low birth weight, reduced birth size and congenital malformations [16, 17]. Epidemiological research on maternal exposure to air pollutants and birth outcomes are suggestive of exposure derived pregnancy problems which are further associated with chromosomal aberrations, decreased neurological development as well as high risk of chronic diseases. The toxicity of pollutants also causes an inflammatory response to weaken both innate as well as adaptive immune response [72].

18-34 week pregnant women were involved in a questionnaire and household chemical (indoor pollutants, formaldehyde, NO_2 , and VOCs) usage survey. In addition, gestational age, birth weight, and length and head circumference were collected from birth records. The association between

pollutants and pollution surrogates were detected using general linear models, controlling for maternal age, parity, maternal health, and season of birth. Only HCHO was associated with any of the birth outcomes. There was a 0.044 decrease in BW z-score ($P = 0.033$) and 0.05 decrease in HC z-score ($P = 0.06$) for each unit increase in HCHO. Although HCHO concentrations were very low, this finding is consistent with other studies of formaldehyde and poor birth outcomes [75].

Residential pollutant exposure was found directly related to pregnancy outcomes by producing small size for gestational age, preterm birth and low birth weight in a population-based cohort study. Pollution exposure was negatively associated with term birth weight [75]. In a similar study conducted by Smith et al. 2017 on 540365 singleton term live births and found average air exposure of NO_2 ($41 \mu\text{g}/\text{m}^3$), nitrogen oxides NO_x ($73 \mu\text{g}/\text{m}^3$), O_3 ($32 \mu\text{g}/\text{m}^3$), $\text{PM}_{2.5}$ ($14 \mu\text{g}/\text{m}^3$) and PM_{10} ($23 \mu\text{g}/\text{m}^3$). Interquartile range increases in NO_2 , NO_x , O_3 , $\text{PM}_{2.5}$ and PM_{10} from traffic exhaust were related to 2% to 6% increase in odds of term low birth weight, 1% to 3% increase in odds of term small for gestational age suggesting adversely affecting fetal growth [77].

A cross-sectional study performed by Milanzi and Namacha, 2017 using secondary data of 9124 respondents revealed exposure to high pollution fuels resulted in a 92 g reduction in mean birth weight [78].

Haifa Pregnancy Cohort Study (HPCS) conducted by Golan et al. 2018 on nearly 750000 newborns and estimated daily air pollutants exposure, temperature, and greenness. It was found that higher environmental pollutants exposure led to poorer fetal growth parameters likewise birth-weight, head-circumference and gestational age at birth [18].

Maternal exposure to incense burning and hypertensive disorders as well as blood pressure of 10563 pregnant women were evaluated by He et al. 2018. It was found that incense smelling during late pregnancy was associated with increased hypertensive risk (relative risk, 1.84) and increased blood pressure (1.6 mmHg systolic blood pressure increase). Incense burning is found a well-established modifiable risk factor for hypertensive disorders [79].

Prenatal air pollution exposure is responsible for early age respiratory problems and is associated with increased maternal pre-pregnancy BMI. Children of first-trimester pregnant obese females (exposed to $\text{PM}_{2.5}$) had an adjusted wheezing incidence ratio of 1.85 (1.23-2.78), 1.76 (1.08-2.85) and 1.90 (1.10-3.27) in quartile 2-4. Additionally, second-trimester exposure was associated with bronchiolitis/bronchitis. First-trimester exposure to $\text{PM}_{2.5}$ was associated with ear infection in children with adjusted Odds Ratio (adjOR) 7.64 (1.18-49.37), 11.37 (1.47-87.97) and 8.26 (1.13-60.29) for quartile 2-4 and second-year with adjOR 3.28 (1.00-10.73) and 4.15 (1.05-16.36) for quartile 2-3 [32].

Van den Eeden et al. 2018 evaluated exposure effect of PM and other ambient pollutants on 200 pregnancy outcomes by biochemical changes in maternal, placental and cord blood. Samples i.e. maternal hair samples, urine samples for 8-oxo-deoxyguanosine, maternal blood samples for transcript markers and biomarkers of pre-eclampsia, neonatal blood samples for transcript markers were analyzed [36].

Xiao et al. 2018 evaluated the relationship $\text{PM}_{2.5}$ exposure in pregnant females and adverse birth outcomes using a gap-filled satellite in Shanghai, China. 132783 live birth records singleton evaluated for association with term birth weight and term low birth weight (LBW) and found whole-pregnancy $\text{PM}_{2.5}$ exposure was associated with a

12.85 g decrease in term birth weight, increased risk of preterm birth (OR 1.27) and increased risk of term LBW (OR 1.22). The scale of associations between $PM_{2.5}$ exposure during pregnancy and adverse birth outcomes was high [39].

Effects of prenatal and postnatal exposure to PM on Kawasaki Disease (KD) was studied using nation population data of 30367 participants (age group 6 to 30 months) at municipality levels. A multilevel logistic regression analysis for individual and municipality level was performed and found that children, who were exposed to higher Suspended Particulate Matter (SPM) level during pregnancy, were hospitalized for KD more than the others. The Odd ratios of PM exposure were 1.59 for prenatal exposure and 1.41 for postnatal exposure suggesting that early-life exposure to PM is associated with an increased risk of Kawasaki disease [40].

Air pollution exposure during pregnancy and risk assessment for the next generation was studied in 1293 mothers and their children of age between 3 and 9 years of age. Multivariable-adjusted cubic spline revealed increased offspring Systolic BP Percentile (SBP) and risk for elevated BP following $PM_{2.5}$ concentrations at $\geq 13 \mu g/m^3$ during third-trimester (4.85 percentile increase in child SBP) [42].

National scenario

Tamil Nadu Air Pollution and Health Effects (TAPHE) study was organized into five components from pregnant mother-child cohort and adult cohort (n=1200 each cohort). Continuous 24-48 h. exposures of $PM_{2.5}$ were measured in household microenvironments together with ambient measurements. The association between $PM_{2.5}$ and air toxicants exposure, maternal birth weight, child acute respiratory infections as well as adult chronic respiratory and lung impairments were examined while adjusting multiple covari-

ates. In addition, a bio-repository of peripheral and cord blood samples was created to investigate gene-environment interactions [6].

Direct serial measurements of 24-h household $PM_{2.5}$ exposures during pregnancy were analyzed for LBW in an integrated rural-urban, mother-child cohort study in Tamil Nadu, India. First-trimester pregnant women (n=1285) from primary health care centers and urban health posts were kept in observation for each trimester until birth to collect antenatal care data and birth weight. A detailed questionnaire was provided to each mother on covariates related to household, socio-economic, demographic and maternal health characteristics. $PM_{2.5}$ exposure ($10 \mu g/m^3$) increase in pregnancy was associated with 4 g (95% CI: 1.08, 6.76 g) decrease in birth weight while 2% increase in its prevalence (OR=1.02; 95% CI:1.005, 1.041) following adjustment for infant sex, gestational age, maternal BMI, maternal age, history of previous LBW, season of conception and birth order [7].

In the household survey to find the correlation between household cooking fuels and risk of stillbirth, 188917 women aged 15-49 were included. Robust standard errors were considered as literacy of parents, in home lighting fuel and cooking fuel used, gravida status, previous abortion history, antenatal checkup history of mother, age during last pregnancy (>35 years), fetus and other complications and premature delivery. The characteristic risk for stillbirths of firewood as cooking fuel in India was 11% and 1% for kerosene cooking. Women who cook with firewood, risk factors were significantly associated with the occurrence of stillbirth (1.24; 95% CI: 1.08-1.41) while kerosene fuel is associated with risk factors (1.36; 95% CI: 1.10-1.67) than those who cook with LPG/electricity. Kerosene lamp usage for home lighting was also associated with stillbirths compared to electric lighting (1.15; 95% CI:

1.06-1.25). By providing access to cleaner cooking fuel about 12% of stillbirths in India could be prevented [25].

Annual secondary health data of 1404337 live births of rural and urban areas from 284 districts of 09 Indian states were analyzed for neonatal mortality rate (NMR) following exposure to cow dung/crop residues/firewood as fuel. Socio-demographic, healthcare and behavioral aspects of the mother were considered as covariates during the study. The average rural NMR was recorded 42.4 out of 1000 live births and urban NMR was 33.1 out of 1000 live births. The proportion of household air pollution was 92.2% ($P < 0.001$) in rural areas as compared to 40.8% in urban areas, and the difference was statistically significant. HAP values were strongly associated with NMR ($\beta = 0.22$; 95% confidence interval [CI] = 0.09 to 0.35) for urban and rural areas combined. For rural areas separately, the association was significant ($\beta = 0.30$; 95% CI = 0.13 to 0.45) after adjustment which represents that HAP is associated with NMR in rural areas, but not in urban areas in India [28].

The association between exposure to indoor biomass fuel sources, second-hand tobacco smoke and adverse health outcomes in early infancy in a population in rural south India was observed by Tielsch et al. 2009. A population-based cohort of newborns from birth through 6 months was performed. Follow up visits for morbidity associated anthropometric measurements were performed bimonthly after delivery. 92.3% of 11728 live-born infants resided in households and used wood and/or dung as a primary fuel source, such exposure was associated with 49% increase in LBW risk, 34% increase in respiratory illness incidence and 21% increase in six-month infant mortality risk. Exposed neonatal also had 45% and 30% increase in risks of underweight and stunting.

Exposure was also linked with adverse health outcomes except for accomplished growth [35].

Conclusion

Pollutant mediated oxidative stress and ROS production following exposure to various air pollutants causes harmful health effects. The biological system though possesses the repair mechanism i.e. antioxidants to neutralize the resulting cellular toxicity, DNA damage and cell death, yet repeated and/or severe pollutant exposure defense mechanism provide insufficient protection. During pregnancy, such exposure severely affects both the maternal organs as well as the neonatal milieu. Maternal pollution exposure during pregnancy is attributed to increased chances of cesarean section, intrauterine growth retardation, preterm birth or small gestational age, stillbirth, increased fetal heart rate, neonatal asphyxia, cerebral palsy, low birth weight, low birth size, congenital malformations, decreased neurological development, chromosomal aberrations causing teratogenic effects, respiratory syndrome like-wise bronchiolitis/bronchitis, sudden infant death syndrome, elevated blood pressure and hypertension and increased risk to Kawasaki disease. Though studies are suggestive of adverse fetal and pregnancy outcomes following air pollutant exposure yet some research provides information about the gender-specific increase in leptin levels following NO_x exposure, no association between stillbirths and accomplished growth caused by maternal complications and pollutant exposure.

List of abbreviations

adjOR- Adjusted Odds Ratio; BMI- Body Mass Index; CAD- Caspase Activated DNase; CAT- Catalase; CI- Confidence Interval; CO- Carbon Monoxide; COS- Controlled Ovarian Stimulation; C-sections- Cesarean Sections; cSLE-Systemic

lupus erythematosus; DENPs- Diesel Exhaust Nanoparticles; EndoG- Endonuclease G; FHR- Fetal Heart Rate; GPx- Glutathione Peroxidase; GST-Glutathione-s-transferase; HAP- Household air pollution; HRs- Hazard Ratios; LBW- Low Birth Weight; MAPK- Mitogen Activated Protein Kinases; NF- κ B- Nuclear Factor Kappa-B; NMR- Neonatal Mortality Rate; NO₂-Nitrogen Dioxide; NO_x- Nitrogen Oxides; OR- Odd Ratio; O₃- Ozone; PAHs- Polycyclic Aromatic Hydrocarbons; PENPs- Petrol Exhaust Nanoparticles; PM- Particulate Matter; ROS- Reactive Oxygen Species; SBP- Systolic BP percentile; SOD- Superoxide Dismutase; SPM-Suspended particulate matter; TTN- Transient tachypnea of the newborn; VLBW- Very Low Birth Weight; VOCs- Volatile Organic Compounds; 8-OHdG- 8-hydroxydeoxyguanosine.

Financial supports

The author(s) received no specific funding for this work.

Competing interest

Authors report no competing interest.

Acknowledgements

Authors are thankful to the Director, ICMR- National Institute of Occupational Health (ICMR-NIOH) for providing necessary facilities.

Ethical considerations

Ethical issues (Including plagiarism, Informed Consent, misconduct, data fabrication and/or falsification, double publication and/or submission, redundancy, etc.) have been completely observed by the authors.

References

1. Durga M, Nathiya S, Rajasekar A, Devasena T. Effects of ultrafine petrol exhaust particles on cytotoxicity, oxidative stress generation, DNA damage and inflammation in human A549 lung cells and murine RAW 264.7 macrophages. *Environmental toxicology and pharmacology*. 2014;38(2):518-30.
2. EPA (Environmental Protection Agency). A decade of children's environmental health research: highlights from EPA's science to achieve results program. Office of research and development, U.S.; Washington, DC.
3. Møller P, Danielsen PH, Karotki DG, Jantzen K, Roursgaard M, Klingberg H, et al. Oxidative stress and inflammation generated DNA damage by exposure to air pollution particles. *Mutation research/Reviews in mutation research*. 2014; 762:133-66.
4. Cao Z, Meng L, Zhao Y, Liu C, Yang Y, Su X, et al. Maternal exposure to ambient fine particulate matter and fetal growth in Shanghai, China. *Environmental health*. 2019;18(1):49.
5. Alderete TL, Song AY, Bastain T, Habre R, Toledo-Corral CM, Salam MT, et al. Prenatal traffic-related air pollution exposures, cord blood adipokines and infant weight. *Pediatric obesity*. 2018;13(6):348-56.
6. Balakrishnan K, Sambandam S, Ramaswamy P, Ghosh S, Venkatesan V, Thangavel G, et al. Establishing integrated rural-urban cohorts to assess air pollution-related health effects in pregnant women, children and adults in Southern India: an overview of objectives, design and methods in the Tamil Nadu air pollution and health effects (TAPHE) study. *BMJ open*. 2015;5(6): e008090.
7. Balakrishnan K, Ghosh S, Thangavel G, Sambandam S, Mukhopadhyay K, Puttaswamy N, et al. Exposures to fine particulate matter (PM_{2.5}) and birthweight in a rural-urban, mother-child cohort in Tamil Nadu, India. *Environmental research*. 2018;161:524-31.
8. Basu R, Pearson D, Ebisu K, Malig B. Association between PM_{2.5} and PM_{2.5} constituents and preterm delivery in California, 2000–2006. *Paediatric and perinatal epidemiology*. 2017;31(5):424-34.
9. Brauer M, Lencar C, Tamburic L, Koehoorn M, Demers P, Karr C. A cohort study of traffic-related air pollution impacts on birth outcomes. *Environmental health perspectives*. 2008;116(5):680-6.
10. Bunkar N, Bhargava A, Chaudhury K, Sharma RS, Lohiya NK, Mishra PK. Fetal nucleic acids in maternal plasma: from biology to clinical translation. *Frontiers in Bioscience*. 2018; 23:397-431.
11. Choe SA, Jun YB, Lee WS, Yoon TK, Kim SY. Association between ambient air pollution and pregnancy rate in women who underwent IVF. *Human reproduction*. 2018;33(6):1071-8.
12. Conde PG, Farhat LC, Braga AL, Sallum AE, Farhat SC, Silva CA. Are prematurity and environmental factors determinants for developing childhood-onset systemic lupus erythematosus? *Modern rheumatology*. 2018;28(1):156-60.
13. Contreras ZA, Heck JE, Lee PC, Cui X, Hobel CJ, Janzen C, et al. Prenatal air pollution exposure, smoking,

- and uterine vascular resistance. *Environmental epidemiology* (Philadelphia, Pa.). 2018;2(3).
14. Dodd-Butera T, Quintana PJ, Ramirez-Zetina M, Batista-Castro AC, Sierra MM, Shaputnic C, et al. Placental biomarkers of PAH exposure and glutathione-S-transferase biotransformation enzymes in an obstetric population from Tijuana, Baja California, Mexico. *Environmental research*. 2017;152:360-8.
 15. Ebisu K, Malig B, Hasheminassab S, Sioutas C, Basu R. Cause-specific stillbirth and exposure to chemical constituents and sources of fine particulate matter. *Environmental research*. 2018;160: 358-64.
 16. Gehring U, Tamburic L, Sbihi H, Davies HW, Brauer M. Impact of noise and air pollution on pregnancy outcomes. *Epidemiology*. 2014;25(3):351-8.
 17. Stieb DM, Chen L, Eshoul M, Judek S. Ambient air pollution, birth weight and preterm birth: a systematic review and meta-analysis. *Environmental research*. 2012;117:100-11.
 18. Golan R, Kloog I, Almog R, Gesser-Edelsburg A, Negev M, Jolles M, et al. Environmental exposures and fetal growth: The Haifa pregnancy cohort study. *BMC public health*. 2018;18(1):132.
 19. Guo LQ, Zhang Q, Zhao DD, Wang LL, Chen Y, Mi BB, et al. Relationship between air pollution exposure during pregnancy and birth weight of term singleton live-birth newborns. *Zhonghua liu xing bing xue za zhi= Zhonghua liuxingbingxue zazhi*. 2017;38(10):1399-403.
 20. Guo T, Wang Y, Zhang H, Zhang Y, Zhao J, Wang Q, et al. The association between ambient PM2.5 exposure and the risk of preterm birth in China: A retrospective cohort study. *Science of the total environment*. 2018;633:1453-9.
 21. Ha S, Sundaram R, Louis GM, Nobles C, Seeni I, Sherman S, et al. Ambient air pollution and the risk of pregnancy loss: a prospective cohort study. *Fertility and sterility*. 2018;109(1):148-53.
 22. Johnson S, Bobb JF, Ito K, Savitz DA, Elston B, Shmool JL, et al. Ambient fine particulate matter, nitrogen dioxide, and preterm birth in New York city. *Environmental health perspectives*. 2016;124(8):1283-90.
 23. Kingsley SL, Eliot MN, Glazer K, Awad YA, Schwartz JD, Savitz DA, et al. Maternal ambient air pollution, preterm birth and markers of fetal growth in Rhode Island: results of a hospital-based linkage study. *Journal of epidemiology and community health*. 2017;71(12):1131-6.
 24. La Marca L, Gava G. Air Pollution Effects in Pregnancy. In: *Clinical handbook of air pollution-related diseases* 2018:479-494 Springer, Cham.
 25. Lakshmi PV, Virdi NK, Sharma A, Tripathy JP, Smith KR, Bates MN, et al. Household air pollution and stillbirths in India: analysis of the DLHS-II national survey. *Environmental research*. 2013;121:17-22.
 26. Lavigne E, Yasseen III AS, Stieb DM, Hystad P, Van Donkelaar A, Martin RV, et al. Ambient air pollution and adverse birth outcomes: differences by maternal co-morbidities. *Environmental research*. 2016;148:457-66.
 27. Morokuma S, Michikawa T, Yamazaki S, Nitta H, Kato K. Association between exposure to air pollution during pregnancy and false positives in fetal heart rate monitoring. *Scientific reports*. 2017;7(1):12421.
 28. Neogi SB, Pandey S, Sharma J, Chokshi M, Chauhan M, Zodpey S, et al. Association between household air pollution and neonatal mortality: an analysis of annual health survey results, India. *WHO South-East Asia journal of public health*. 2015;4(1):30.
 29. Rogers JF, Thompson SJ, Addy CL, McKeown RE, Cowen DJ, Decoufle P. Association of very low birth weight with exposures to environmental sulfur dioxide and total suspended particulates. *American journal of epidemiology*. 2000;151(6):602-13.
 30. Seeni I, Ha S, Nobles C, Liu D, Sherman S, Mendola P. Air pollution exposure during pregnancy: maternal asthma and neonatal respiratory outcomes. *Annals of epidemiology*. 2018;28(9):612-8.
 31. Soh SE, Goh A, Teoh OH, Godfrey KM, Gluckman PD, Shek LP, et al. Pregnancy trimester-specific exposure to ambient air pollution and child respiratory health outcomes in the first 2 years of life: Effect modification by maternal pre-pregnancy BMI. *International journal of environmental research and public health*. 2018;15(5):996.
 32. Son JY, Lee HJ, Koutrakis P, Bell ML. Pregnancy and lifetime exposure to fine particulate matter and infant mortality in Massachusetts, 2001-2007. *American journal of epidemiology*. 2017;186(11):1268-76.
 33. Song J, Chen Y, Wei L, Ma Y, Tian N, Huang SY, et al. Early-life exposure to air pollutants and adverse pregnancy outcomes: protocol for a prospective cohort study in Beijing. *BMJ open*. 2017;7(9):e015895.
 34. Tanwar V, Gorr MW, Velten M, Eichenseer CM, Long III VP, Bonilla IM, et al. In utero particulate matter exposure produces heart failure, electrical remodeling, and epigenetic changes at adulthood. *Journal of the American heart association*. 2017;6(4):e005796.
 35. Tielsch JM, Katz J, Thulasiraj RD, Coles CL, Sheeladevi S, Yanik EL, et al. Exposure to indoor biomass fuel and tobacco smoke and risk of adverse reproductive outcomes, mortality, respiratory morbidity and growth among newborn infants in south India. *International journal of epidemiology*. 2009;38(5):1351-63.
 36. Van den Eeden L, Lambrechts N, Verheyen V, Berth M, Schoeters G, Jacquemyn Y. Impact of particulate matter on mothers and babies in Antwerp (IPANEMA): a prospective cohort study on the impact of pollutants and particulate matter in pregnancy. *BMJ open*. 2018;8(3):e020028.
 37. Winkelmann E, Cox B, Martens E, Fierens F, Nemery B, Nawrot TS. Fetal growth and maternal exposure to particulate air pollution--More marked effects at lower exposure and modification by gestational duration. *Environmental research*. 2015;140:611-8.
 38. Wu J, Wilhelm M, Chung J, Ritz B. Comparing expo-

- sure assessment methods for traffic-related air pollution in an adverse pregnancy outcome study. *Environmental research*. 2011;111(5):685-92.
39. Xiao Q, Chen H, Strickland MJ, Kan H, Chang HH, Klein M, et al. Associations between birth outcomes and maternal PM_{2.5} exposure in Shanghai: A comparison of three exposure assessment approaches. *Environment international*. 2018;117:226-36.
40. Yorifuji T, Tsukahara H, Kashima S, Doi H. Intrauterine and early postnatal exposure to particulate air pollution and kawasaki disease: A nationwide longitudinal survey in Japan. *The Journal of pediatrics*. 2018;193:147-54.
41. Zhang Q, Xiang XM, Song H, Dong M, Zhang SP, Mi BB, et al. Relationship between exposure to air pollutants during pre-pregnancy or early pregnancy and birth defects in Xi'an, 2013-2015: a case crossover study. *Zhonghua liu xing bing xue za zhi=Zhonghua liuxingbingxue zazhi*. 2017;38(12):1677-82.
42. Zhang M, Mueller NT, Wang H, Hong X, Appel LJ, Wang X. Maternal exposure to ambient particulate matter $\leq 2.5 \mu\text{m}$ during pregnancy and the risk for high blood pressure in childhood. *Hypertension*. 2018;72(1):194-201.
43. Zhu Y, Zhang C, Liu D, Grantz KL, Wallace M, Mendola P. Maternal ambient air pollution exposure pre-conception and during early gestation and offspring congenital orofacial defects. *Environmental research*. 2015;140:714-20.
44. Jain KK. Carbon monoxide and other tissue poisons. In: *Textbook of hyperbaric medicine* 2017;131-154. Springer, Cham.
45. Queiroga CS, Almeida AS, Vieira HL. Carbon monoxide targeting mitochondria. *Biochemistry research international*. 2012;2012.
46. Rose JJ, Wang L, Xu Q, McTiernan CF, Shiva S, Tejero J, Gladwin MT. Carbon monoxide poisoning: pathogenesis, management, and future directions of therapy. *American journal of respiratory and critical care medicine*. 2017;195(5):596-606.
47. Manoj KM, Ramasamy S, Parashar A, Soman V, Pakshirajan K. Murburn concept explains the acutely lethal effect of cyanide. *bioRxiv*. 2019;1:555888.
48. Mansoor K, Parkins G, Wilson L, White J, Shiflett BS, Ajmeri A, Zeid MD. Carbon Monoxide: A rare cause of myocardial ischemia. *Marshall journal of medicine*. 2019;5(1):3.
49. Thom SR. Options for treatment of carbon monoxide poisoning, including hyperbaric oxygen therapy. *Carbon monoxide*. 2019;12:215.
50. Singer M. Mitochondrial Function. In: *Hemodynamic Monitoring* 2019;97-106. Springer, Cham.
51. McCarthy JT, Pelle E, Dong K, Brahmabhatt K, Yarosh D, Pernodet N. Effects of ozone in normal human epidermal keratinocytes. *Experimental dermatology*. 2013;22(5):360-1.
52. Jiang C, Stewart LT, Kuo HC, McGilberry W, Wall SB, Liang B, Van Groen T, Bailey SM, Kim YI, Tipple TE, Jones DP. Cyclic O₃ exposure synergizes with aging leading to memory impairment in male APOE $\epsilon 3$, but not APOE $\epsilon 4$, targeted replacement mice. *Neurobiology of aging*. 2019;81:9-21.
53. Stieb DM, Yao J, Henderson SB, Pinault L, Smith-Doiron MH, Robichaud A, van Donkelaar A, Martin RV, Ménard R, Brook JR. Variability in ambient ozone and fine particle concentrations and population susceptibility among Canadian health regions. *Canadian journal of public health*. 2019;110(2):149-58.
54. Duan C, Talbott E, Brooks M, Park SK, Broadwin R, Matthews K, et al. Five-year exposure to PM_{2.5} and ozone and subclinical atherosclerosis in late midlife women: The study of women's health across the nation. *International journal of hygiene and environmental health*. 2019;222(2):168-76.
55. Fuks KB, Woodby B, Valacchi G. Skin damage by tropospheric ozone. *Der Hautarzt*. 2019;1:1-5.
56. Meng Z, Qin G, Zhang B, Bai J. DNA damaging effects of sulfur dioxide derivatives in cells from various organs of mice. *Mutagenesis*. 2004;19(6):465-8.
57. Ji PY, Li ZY, Wang H, Dong JT, Li XJ, Yi HL. Arsenic and sulfur dioxide co-exposure induce renal injury via activation of the NF- κ B and caspase signaling pathway. *Chemosphere*. 2019; 224:280-8.
58. Rosser FJ, Forno E, Acosta-Perez E, Canino G, Celedon JC. Annual Sulfur Dioxide Exposure, Asthma, and Lung Function in Puerto Rican Children. In: *D96. Environmental asthma epidemiology*. American Thoracic Society. 2019;A7055.
59. Madhu BP, Singh KP, Saminathan M, Singh R, Shivasharanappa N, Sharma AK, et al. Role of nitric oxide in the regulation of immune responses during rabies virus infection in mice. *Virusdisease*. 2016;27(4):387-99.
60. Wlodkowic D, Telford W, Skommer J, Darzynkiewicz Z. Apoptosis and beyond: cytometry in studies of programmed cell death. In: *Methods in cell biology* 2011;103: 55-98. Academic Press.
61. Tekpli X, A Holme J, Sergent O, Lagadic-Gossmann D. Importance of plasma membrane dynamics in chemical-induced carcinogenesis. *Recent patents on anti-cancer drug discovery*. 2011;6(3):347-53.
62. Chen D, Stetler RA, Cao G, Pei W, O'Horo C, Yin XM, et al. Characterization of the rat DNA fragmentation factor 35/inhibitor of caspase-activated DNase (Short Form) the endogenous inhibitor of caspase-dependent DNA fragmentation in neuronal apoptosis. *Journal of biological chemistry*. 2000;275(49):38508-17.
63. Ishihara Y, Shimamoto N. Involvement of endonuclease G in nucleosomal DNA fragmentation under sustained endogenous oxidative stress. *Journal of biological chemistry*. 2006;281(10):6726-33.
64. Norberg E, Orrenius S, Zhivotovsky B. Mitochondrial regulation of cell death: processing of apoptosis-inducing factor (AIF). *Biochemical and biophysical research communications*. 2010;396(1):95-100.

65. Van Meteren N, Chevanne M, Gallais I, Gauffre F, Chevance S, Burel A, et al. Polycyclic aromatic hydrocarbons can trigger a hepatocyte release of cytotoxic extracellular vesicles. *Journal of Hepatology* 2019;70 E812. Elsevier science BV, Amsterdam, Netherlands.
66. Peixoto MS, de Oliveira Galvão MF, de Medeiros SR. Cell death pathways of particulate matter toxicity. *Chemosphere*. 2017;188:32-48.
67. Bhargava A, Tamrakar S, Aglawe A, Lad H, Srivastava RK, Mishra DK, et al. Ultrafine particulate matter impairs mitochondrial redox homeostasis and activates phosphatidylinositol 3-kinase mediated DNA damage responses in lymphocytes. *Environmental pollution*. 2018;234: 406-19.
68. Laing S, Wang G, Briazova T, Zhang C, Wang A, Zheng Z, et al. Airborne particulate matter selectively activates endoplasmic reticulum stress response in the lung and liver tissues. *American journal of physiology-cell physiology*. 2010;299(4):C236-49.
69. Andreau K, Leroux M, Bouharrour A. Health and cellular impacts of air pollutants: from cytoprotection to cytotoxicity. *Biochemistry research international*. 2012;2012.
70. Veronesi B, Oortgiesen M, Carter JD, Devlin RB. Particulate matter initiates inflammatory cytokine release by activation of capsaicin and acid receptors in a human bronchial epithelial cell line. *Toxicology and applied pharmacology*. 1999;154(1):106-15.
71. Devasagayam TP, Tilak JC, Boloor KK, Sane KS, Ghaskadbi SS, Lele RD. Free radicals and antioxidants in human health: current status and future prospects. *Journal of association of physicians of India*. 2004;52(794804):4.
72. Huff RD, Carlsten C, Hirota JA. An update on immunologic mechanisms in the respiratory mucosa in response to air pollutants. *Journal of allergy and clinical immunology*. 2019;1;143(6):1989-2001.
73. Hiemstra PS, Grootaers G, van der Does AM, Krul CAM, Kooter IM. Human lung epithelial cell cultures for analysis of inhaled toxicants: lessons learned and future directions. *Toxicology in vitro*. 2018;47:137-46.
74. Sies H, Berndt C, Jones DP. Oxidative stress. *Annual review of biochemistry*. 2017;86: 715-48.
75. Franklin P, Tan M, Hemy N, Hall GL. Maternal exposure to indoor air pollution and birth outcomes. *International journal of environmental research and public health*. 2019;16(8):1364.
76. Bauer RN, Diaz-Sanchez D, Jaspers I. Effects of air pollutants on innate immunity: the role of Toll-like receptors and nucleotide-binding oligomerization domain-like receptors. *Journal of allergy and clinical immunology*. 2012;129(1):14-24.
77. Smith RB, Fecht D, Gulliver J, Beevers SD, Dajnak D, Blangiardo M, et al. Impact of London's road traffic air and noise pollution on birth weight: Retrospective population based cohort study. *BMJ*. 2017;359:j5299.
78. Milanzi EB, Namacha NM. Maternal biomass smoke exposure and birth weight in Malawi: Analysis of data from the 2010 Malawi demographic and health survey. *Malawi medical journal*. 2017;29(2):160-5.
79. He JR, Wei DM, Chan FF, Luan YZ, Tu S, Lu JH, et al. Associations between maternal exposure to incense burning and blood pressure during pregnancy. *Science of the total environment*. 2018; 610:1421-7.