



# Association Between Sexual and Genital Hygiene Habits With the Urinary Tract Infection During Pregnancy: A Case-Control Study

Simin Haghdoost<sup>1</sup>, Farzaneh Pazandeh<sup>2\*</sup>, Mehdi Khabazkhoob<sup>3</sup>, Tahereh Behroozi Lak<sup>4</sup>

## Abstract

**Objectives:** Urinary tract infections (UTIs) are the most common type of infection during pregnancy, which cause serious complications for the mother and baby. The present study was conducted to investigate the relationship between sexual and genital hygiene habits and UTIs in pregnant women.

**Materials and Methods:** This case-control study was conducted on 187 pregnant women including 97 pregnant women with symptomatic bacteriuria (case) and 90 healthy pregnant women (control). Controls were frequency matched with the cases with respect to the number of pregnancy and trimesters of pregnancy. Both groups were compared for the presence of UTI. Finally, the binary multivariate unconditional logistic regression approach was used to evaluate the association between UTI and the confounding variables.

**Results:** The results of this study showed a significant association between UTI in pregnancy and sexual and genital hygiene habits ( $P < 0.05$ ). Based on the binary multivariate logistic regression analysis, attributable risks for UTI included having sexual intercourse  $> 3$  times/week (adjusted odds ratio [AOR] = 3.68, 95% CI = 2.09-5.41,  $P = 0.001$ ), changing the underwear per week (AOR = 1.39, 95% CI = 1-1.76,  $P = 0.012$ ), not voiding post-coitus (AOR = 2.01, 95% CI = 1.34-2.69,  $P = 0.01$ ), washing genitalia from back to the front (AOR = 1.96, 95% CI = 1.06-2.78,  $P = 0.01$ ), and not washing genitals pre-coitus by the husband (AOR = 1.20, 95% CI = 0.54-1.9,  $P = 0.024$ ) were among the sexual and hygiene habit variables associated with UTI in pregnant women in this model.

**Conclusions:** In general, sexual and genital hygiene habits are vital preventive actions for controlling the UTI in pregnant women. These actions may lead to UTI complications and a preterm birth reduction. Thus, informing spouses about these practices is highly suggested.

**Keywords:** Sexual habits, Hygienic habits, Urinary tract infection, Pregnancy

## Introduction

Urinary tract infections (UTIs) are considered as the most common type of bacterial infection during pregnancy. In addition, UTIs during pregnancy are classified into the infections of the lower (cystitis) or upper (pyelonephritis) urinary tracts. The lower UTIs are associated with a 40% increased risk of developing pyelonephritis in pregnancy, which occur in response to the physiological changes in the urinary tract during pregnancy, an increase in sex hormones, the increased bladder volume, and decreased bladder tone, along with the decreased urethral tone contributing to increased urinary stasis and vesicoureteral reflux (1,2). These changes, along with a short urethra-vagina distance increase the frequency of UTI during pregnancy (3). The prevalence of UTI is considerable among pregnant women in Asian countries such as Pakistan (12.4%), Turkey (8.5%), India (13.3%), and Iran (13%) (4-7). Further, UTI during pregnancy can have

serious complications affecting the health of both mother and fetus, including preterm labor, low birth weight, intrauterine growth retardation, fetal death, or maternal systemic infection (2,8).

The UTI prevalence is considerable among Iranian pregnant women and antibiotics resistance is also high in Iran (7). Importantly, preventing UTI is essential for maternal and infant health improvement, as well as the reduction of the preterm birth risk (9). There is inadequate evidence about the influence of sexual and genital hygiene habits on UTI during pregnancy. Thus, the current study aimed to determine the association between UTI and sexual and genital hygiene habits in pregnant women.

## Materials and Methods

### Study Design and Selection of Cases and Controls

In this case-control study, the association between sexual and genital hygiene habits and UTI was investigated in

Received 15 June 2019, Accepted 11 August 2019, Available online 9 September 2019

<sup>1</sup>Student Research Committee, Department of Midwifery and Reproductive Health, School of Nursing and Midwifery, Shahid Beheshti University of Medical Sciences, Tehran, Iran. <sup>2</sup>Department of Midwifery and Reproductive Health, Midwifery and Reproductive Health Research Center, Shahid Beheshti University of Medical Sciences Tehran, Iran. <sup>3</sup>Department of Noor Research Center for Ophthalmic Epidemiology, Noor Eye Hospital, Tehran, Iran. <sup>4</sup>Reproductive Health Research Center, Department of Infertility, Urmia University of Medical Sciences, Urmia, Iran.

\*Corresponding Author: Farzaneh Pazandeh, Email: pazandehf@gmail.com



pregnant women seeking care in Imam Reza Hospital in Urmia, the Northwest of Iran from February to July in 2018. In general, 187 pregnant women including 97 cases diagnosed with a symptomatic UTI were compared with 90 healthy controls who had neither symptomatic nor asymptomatic bacteriuria. The inclusion criteria for both groups were the age range between 18 and 35 years old. On the other hand, the exclusion criteria for both groups included a history of recurrent UTI, urinary tract anomalies, anemia, obesity (body mass index of  $30 \geq \text{kg/m}^2$ ), diabetes, urinary tract stent, urinary incontinence, neurogenic bladder, asymptomatic bacteriuria (10), kidney stones, or substance abuse, and the use of immunosuppressive drugs. The two groups were matched in terms of the number of pregnancies and trimesters of pregnancy. Furthermore, the group matching was performed to select the control group who met the inclusion criteria and was presented in the prenatal care clinic. Moreover, the case group included pregnant women within the age range of 18-35 years old who attended the prenatal care clinic complaining of UTI symptoms. Those who met the inclusion criteria were consecutively enrolled from the prenatal care clinic until attaining the calculated sample size.

#### Data Collection

The data were collected in the prenatal care clinic by using a self-administered questionnaire. The questionnaire included three sections. The first section was related to demographic and obstetric characteristics. The second section was about sexual habits including the number of sexual intercourses per week, pre-coitus genitals washing by the husband, and post-coitus voiding by the woman. Additionally, the third part included some questions about the genital hygiene habits including changing the underwear clothes per week, drying genitalia after washing, washing genitalia by the women, and considering the amount of the daily intake of water. The data about the history of diagnosis, treatment, and the results of laboratory analysis were extracted from medical charts in the clinics. In addition, clinical signs, urinalysis, and urine cultures were used for the diagnosis of UTI in the case group by obstetricians working in the prenatal care clinic. Further, the results of the urine cultures and urinalysis were used to confirm the diagnosis of UTI. The clinical signs of UTI were defined by dysuria, urgency, and the frequency of urination (acute cystitis), or the presence of fever ( $>38^\circ\text{C}$ ), chills, vomiting, nausea and/or loin pain in addition to the presence and absence of the symptoms of cystitis (acute pyelonephritis) (11). Furthermore, the abnormal urine analysis was characterized by pyuria (detecting more than 10 leukocytes per microscopic field), as well as the presence of red blood cells and positive nitrite test (2,11). Gestational age was estimated based on the last menstrual period. Body mass index (BMI) was calculated using height and weight measurements and stratified to

four categories including normal BMI  $\geq 18.5$ - $24.9 \text{ kg/m}^2$ , underweight BMI  $< 18.5 \text{ kg/m}^2$ , overweight BMI  $\geq 25.0$ - $29.9 \text{ kg/m}^2$ , and obese BMI  $\geq 30.0 \text{ kg/m}^2$ .

#### Diagnostic Assessment

UTI was diagnosed by midstream urine samples in the HiChrome UTI Blood Agar plate. Moreover, the samples were incubated under aerobic conditions at  $37^\circ\text{C}$  and then read after 48 hours. The growth of more than  $10^5$  colony-forming units of one organism per milliliter on the urine culture plate was considered positive (2).

#### Sample Size

The sample calculation was based on the formula used for a case-control study according to the study by Jorde et al (12). The sample size was determined considering a 95% confidence interval ( $Z_{\alpha/2} = 1.96$ ), 90% power ( $Z_{\beta} = 1.28$ ), with a case-to-control ratio of 1:1, as well as considering a 15% non-response rate. Therefore, the final sample size was calculated as 180 (90 cases and 90 controls).

#### Statistical Analysis

The Kolmogorov-Smirnov test was done to evaluate the normal distribution of data. Additionally, categorical variables were evaluated using the chi-square test or Fisher exact test, as appropriate. Similarly, the continuous variables were assessed using the *t* test or the Mann-Whitney U test, as appropriate, in order to determine the association between the variables in case and control groups. A *P* value of less than 0.05 was considered statistically significant. In addition, binary multivariate unconditional logistic regression analysis was used for adjusted genital hygiene habits and sexual variables, as well as demographic characteristics to determine the association between the risk factors and UTI. Eventually, the data were analyzed in SPSS software, version 24.

#### Results

Overall, 187 participants were recruited, including 97 pregnant women with UTI (case group) and 90 pregnant women without UTI (control group). The demographic and obstetrical characteristics of the participants are shown in Table 1. The mean gestational age of case and control groups was  $26 \pm 11.7$  and  $25.1 \pm 10.5$  weeks, respectively. Further, the mean BMI of women in case and control groups was equal to  $24.7 \pm 2.9$  and  $24.62 \pm 3 \text{ kg/m}^2$ , respectively. As subjects in both groups were matched with respect to the gravid and gestational age, no significant differences were observed in this regard. However, there was a significant difference regarding the monthly income of the participants between the two groups ( $P < 0.05$ , Table 1).

Table 2 summarizes the results of the comparison of sexual and genital hygiene habits between the participants. There were significant differences in the association

**Table 1.** Demographic and Obstetric Characteristics of the Participants

Variable		Case (n = 97) No. (%)	Control (n = 90) No. (%)	P Value
Maternal age (year, Mean ± SD)		26.4±4.49	27.5±4.69	0.106 <sup>a</sup>
Monthly income	Low	72 (74.%)	53 (60%)	0.034 <sup>b</sup>
	Moderate	20 (20.7%)	28 (29.7%)	
	High	5 (5.3%)	9 (10.3%)	
Education	University	16 (16.5%)	21 (22.2%)	0.255 <sup>b</sup>
	High school	31 (32.4%)	35 (40%)	
	Middle school	21 (21.6%)	17 (18.3%)	
	Primary school	25 (25.4%)	14 (15%)	
	Illiterate	4 (4.1%)	3 (4.5%)	
Husband education	University	21 (21.6%)	22 (23.2%)	0.285 <sup>b</sup>
	High school	33 (33.9%)	37 (41.3%)	
	Middle school	19 (19.6%)	20 (23.2%)	
	Primary school	22 (22.8%)	11 (12.3%)	
	Illiterate	2 (2.1%)	0	
Gravida	One	27 (28%)	23 (25.6%)	0.595 <sup>b</sup>
	Two	38 (39.6%)	39 (43.2%)	
	Three	24 (24.4%)	21 (22.7%)	
	Four	8 (8%)	7 (8.8%)	
BMI	Normal (18.5-24.9 kg/m <sup>2</sup> )	54 (56.7%)	50 (57.8%)	0.625 <sup>c</sup>
	Over weight (25-29.9 kg/m <sup>2</sup> )	43 (43.3%)	40 (42.2%)	
Gestational age	First trimester	22 (22.6%)	24 (26.7%)	0.570 <sup>b</sup>
	Second trimester	26 (26.6%)	26 (28.9%)	
	Third trimester	49 (50.8%)	40 (44.4%)	

Note. SD: Standard deviation; BMI: Body mass index.

<sup>a</sup> Student's *t* test.

<sup>b</sup> Mann-Whitney values are presented as mean ± SD.

<sup>c</sup> Chi-square values are presented as numbers (%).

**Table 2.** Comparison of Sexual and Genital Hygiene Habits Between the Participants

Variable		Case (n = 97)	Control (n = 90)	P Value
Sexual intercourse per week	<3 times	55 (56.7%)	72 (80%)	0.001 <sup>a</sup>
	≥3 times	47 (43.3%)	18 (20%)	
Changing the underwear per week	<3 times	42 (43.3%)	28 (31.1%)	0.031 <sup>a</sup>
	≥3 times	55 (56.7%)	62 (68.9%)	
Drying genitalia after washing	Yes	81 (83.5%)	76 (84.4%)	0.833 <sup>a</sup>
	No	16 (16.5%)	14 (15.6%)	
Washing genitals pre-coitus by husband	Yes	73 (75.4%)	81 (90%)	0.013 <sup>a</sup>
	No	24 (24.6%)	9 (10%)	
Voiding postcoitus by the woman	Yes	74 (76.4%)	85 (94.4%)	0.007 <sup>a</sup>
	No	23 (23.6%)	5 (5.6%)	
Washing genitals	From front to back	71 (73.3%)	80 (88.9%)	0.014 <sup>a</sup>
	From back to front	26 (26.7%)	10 (11.1%)	
Daily intake of water (Mean ± SD)		2±1.08 glasses	2.5±1.05 glasses	0.426 <sup>b</sup>

Note. SD: Standard deviation.

<sup>a</sup> Chi-square values are presented as numbers (%).

<sup>b</sup> Mann-Whitney U values are presented as mean ± SD.

between the UTI and some sexual and genital hygienic habits variables such as having sexual intercourse, changing the underwear per week, not voiding post-coitus by woman, not washing genitals pre-coitus by husband, washing genitals from back to front, and changing the underwear between the study groups ( $P < 0.05$ , Table 2). In this study, *Escherichia coli* was reported as the most common pathogen among pregnant women (70.3%),

followed by *Staphylococcus aureus* (11.1%) as the second most common pathogen (Table 3).

Table 4 presents the results of the multivariate unconditional logistic regression analysis. The association was evaluated between the confounding risk factors, namely, sexual and genital hygiene habits (i.e., Considering the frequency of intercourse per week, not voiding post-coitus, washing genitals from back to front,

**Table 3.** The Types of Bacteriologic Species Isolated From the Urine Culture of Symptomatic Pregnant Women With UTI

Germ	N	%
<i>Escherichia coli</i>	68	70.3
<i>Staphylococcus aureus</i>	11	11.1
<i>Citrobacter spp.</i>	1	1.1
<i>Staphylococcus epidermidis</i>	5	5.1
<i>Klebsiella spp.</i>	7	7.2
<i>Pseudomonas</i>	5	5.2
Total	97	100

Note. UTI: urinary tract infection.

not washing genitals pre-coitus by the husband, changing the underwear per week, drying genitalia per week,

and considering daily water intake) and demographic characteristics (i.e., BMI, monthly income, women's age, and the education level) and UTI. The presence of at least one of these factors was considered as a predictive of UTI: Intercourse >3 times per week (OR = 3.68, 95% CI = 2.09-5.41,  $P=0.001$ ), not voiding post-coitus (OR = 2.01, 95% CI = 1.34-2.69,  $P=0.001$ ), washing genitals from back to front (OR = 1.96, 95% CI = 1.06-2.78,  $P=0.01$ ), not washing genitals pre-coitus by the husband (OR = 1.20, 95% CI = 0.54-1.9,  $P=0.024$ ), and changing the underwear less than three times per week (OR = 1.39, 95% CI = 1-1.76,  $P=0.012$ ) were among the variables of sexual and genital hygiene habits that were associated with UTI in the binary multiple logistic regression model. Women's (OR = 1.29, 95% CI = 0.88-2.94,  $P=0.043$ ) and husbands' (OR = 1.15, 95% CI = 0.18-2.32,  $P=0.049$ ) primary level of education

**Table 4.** The Multiple Logistic Regression Analysis Adjusted for Sexual and Genital Hygiene Habits and Confounding Variables of UTI

Variable	Category	Adjusted OR (95% CI)	P Value
Sexual intercourse per week	Intercourse per week ≤3 times	1	
	Intercourse per week >3 times	3.68 (2.09-5.41)	<0.001
Voiding post-coitus by the woman	Voiding after coitus	1	
	Not voiding after coitus	2.01 (1.34-2.69)	<0.001
Washing genitals from back to front	Washing genitals from front to back	1	
	Washing genitals from back to front	1.96 (1.06-2.78)	0.01
Washing genitals pre-coitus by husband	Washing genitals pre-coitus	1	
	Not washing genitals pre-coitus	1.20 (0.54-1.9)	0.024
Changing underwear per week	Changing underwear more than three time per week	1	
	Changing underwear less than three time per week	1.38 (1-1.76)	0.012
Drying genitalia after washing	Yes	1	
	No	1.17 (0.51-2.64)	0.709
Daily water intake	1-2 glass	1	
	3-4 glass	1.49 (0.22-9.84)	0.67
	5-6 glass	0.62 (0.10-3.77)	0.60
	7-8 glass	1.32 (0.20-8.32)	0.76
	>8 glass	0.55 (0.8-3.63)	0.53
BMI	Normal	1	
	Over weight	1.21 (0.66-2.23)	0.528
Family income (monthly)	High	1	
	Moderate	1.41 (0.77-2.80)	0.32
	Low	5.03 (1-7.64)	0.052
Education	University	1	
	High school	0.37 (0.06-2.28)	0.28
	Middle school	0.77 (0.24-2.48)	0.66
	Primary school	1.29 (0.88-2.94)	0.043
	Illiterate	0.97 (0.35-2.76)	0.96
Age	18-23 year	1	
	24-29 year	0.57 (0.22-1.48)	0.25
	30-35 year	1.13 (0.53-2.41)	0.73
Husband's education	University	1	
	High school	0.41 (0.07-2.71)	0.33
	Middle school	0.42 (0.11-1.58)	0.20
	Primary school	1.15 (0.48-2.32)	0.049
	Illiterate	1.11 (0.38-3.18)	0.84

Note. UTI: Urinary tract infections; OR: odds ratio; CI: confidence interval; BMI: body mass index.

were also among the demographic characteristics that were significantly associated with the risk of UTI in pregnant women in this model.

### Discussion

The results of this study showed that sexual and genital hygiene habits were associated with UTI in pregnant women after adjusting for the confounders. In this study, most of the identified risk factors for UTI were similar to those found in previous studies on the urogenital infections of non-pregnant women in the menstruating age (3,13). In accordance with these results, two observational studies evaluated the association between UTI and sexual and genital hygiene habits during pregnancy. Amiri et al, in a case-control study on a total of 250 pregnant women with asymptomatic bacteriuria, found that sexual and genital hygiene habits such as sexual intercourse for more than or equal to 3 times per week, not washing genitals pre-coitus, not voiding after coitus, and washing genitals from back to front may be associated with UTI in pregnant women (14). Similarly, Abdel-Aziz Elzayat et al reported that genital hygiene habits such as washing genitals from back to front and having sexual intercourse for more than or equal to 2 times per week in pregnant women were associated with the incidence of UTIs (15).

In our study, sexual intercourse for more than 3 times per week was related to UTI, attributing to the fact that the mechanical manner of sexual intercourse may increase the transfer of uropathogens into the urethra and bladder. This relationship was also reported for recurrent cystitis (3,9,16). However, washing the genital area and voiding after intercourse may have a preventive effect against the UTI. The results of our study demonstrated that women who usually did not void post-coitus were most likely to develop the UTI compared to those who did urinate after intercourse, which is in line with the results of studies by Ahmed et al and Amiri et al (14,17). UTI was also found to be prevalent among pregnant women whose husbands did not wash genitals pre-coitus. These findings are in agreement with those of Amiri et al and Mohamed et al (14,18).

Based on the results of the present study, the direction of wiping or washing the genital area after voiding was considered to be important and women who washed their genital area from back to front showed a higher incidence of UTIs. This was reported in some other studies like Ahmed et al (17), Abdel-Aziz Elzayat (15), and Amiri et al (14). The hygiene habits related to toilet hygiene such as wiping or washing perineal area from back to front after urination or defecation are more likely to develop the anal or vaginal flora into the urethra (19). Thus, educating the women on the direction of washing the genital and suggestion for urination after sexual can decrease the prevalence of UTI in pregnant women (20).

The results of this study also revealed that women who changed the underwear less than three times per week were

more at the risk of UTI. Excessive vaginal discharge due to physiological and hormonal changes during pregnancy would lead to some consequences like an increase in the amount of moisture, microbial growth, and urogenital infection risk (21,22). Previous studies suggested that the frequent changing of underwear can decrease the risk of urogenital infection (14,19).

In our study, the mean of daily water intake was low in both groups, but there was no significant difference between the groups, which was not consistent with the results of other studies (14,23). This decrease in daily water intake may be due to the intake of other liquids instead of water.

Finally, it was found that spouses with low education levels were more at risk for UTI. In other words, the risk of UTI was 1.29 times higher in the group of women who had the primary level of education compared to women who had a higher level of education. In addition, the risk of UTI was 1.15 times higher in the group of men who had a primary level of education compared to those who had a higher level of education. Additionally, some Iranian studies reported that a low educational level was associated with an increase in the UTI risk in pregnant women (24-26). Sexual hygienic habits are influenced by the knowledge, beliefs, and habits considering health improvement as the main goal (27). Therefore, educating the spouses about the effect of behavioral risk factors on UTI is very important in planning strategies for preventing UTI during pregnancy. Furthermore, educating the spouses by health professionals with respect to their culture can increase their health knowledge about the sexual and genital hygiene and reduce the risk of urogenital infection among pregnant women (28,29).

The findings of the present study revealed that *E. coli* was the most frequent organism in positive cases, followed by *S. aureus*. In this study, staphylococcal infection was related to poor personal hygiene habits. These findings corroborate with those of previous studies conducted in Pakistan, Egypt, and Iran. Based on these studies, *S. aureus* was the second most frequent organism followed by *E. coli* (18,30,31). The increased levels of amino acids and lactose stimulate *E. coli* growth in the urinary tract, which could be due to the infection through faecal contamination as a result of poor hygiene habits (32).

Inaccurate responses about the sexual and genital hygiene habits of the participants could be considered as a limitation of this study. Therefore, randomized clinical trials are needed to better investigate the efficacy of sexual and genital hygiene habits in reducing the UTI prevalence.

### Conclusions

The results of this study showed that there was a relationship between sexual and genital hygiene habits and UTI in pregnant women. Our findings are very important in relation to planning the preventive strategies for UTIs during pregnancy. Therefore, it is recommended

to educate pregnant women about sexual and hygiene genital habits in prenatal sessions.

### Conflict of Interests

Authors declare that they have no conflict of interests.

### Ethical Issues

The study design was approved by the Ethics Committee (code 901.1396.PHNM.SBMU.IR) of Shahid Beheshti University of Medical Sciences in Tehran, Iran. All participants were informed about the purpose of the study and provided signed informed consent forms.

### Financial Support

This study was financially supported by Shahid Beheshti University of Medical Sciences.

### Acknowledgments

This paper was extracted from the MSc thesis submitted by Simin Haghdoost. We are very grateful to all of the participant women for giving their time, as well as the prenatal care unit and clinics of the Imam Reza Hospital in Urmia, Iran for their collaboration.

### References

1. Foxman B. Urinary tract infection syndromes: occurrence, recurrence, bacteriology, risk factors, and disease burden. *Infect Dis Clin North Am.* 2014;28(1):1-13. doi:10.1016/j.idc.2013.09.003
2. Kalinderi K, Delkos D, Kalinderis M, Athanasiadis A, Kalogiannidis I. Urinary tract infection during pregnancy: current concepts on a common multifaceted problem. *J Obstet Gynaecol.* 2018;38(4):448-453. doi:10.1080/01443615.2017.1370579
3. Gyftopoulos K, Matkaris M, Vourda A, Sakellaropoulos G. Clinical implications of the anatomical position of the urethra meatus in women with recurrent post-coital cystitis: a case-control study. *Int Urogynecol J.* 2019;30(8):1351-1357. doi:10.1007/s00192-018-3710-7
4. Shuja S, Pario S, Muneeb A. Asymptomatic Bacteriuria in Women Presenting to Antenatal Clinic at a Tertiary Care Hospital in Karachi, Pakistan. *Pak Armed Forces Med J.* 2018;68(5):1138-1142.
5. Rajaratnam A, Baby NM, Kuruville TS, Machado S. Diagnosis of asymptomatic bacteriuria and associated risk factors among pregnant women in mangalore, karnataka, India. *J Clin Diagn Res.* 2014;8(9):Oc23-25. doi:10.7860/jcdr/2014/8537.4842
6. Celen S, Oruç AS, Karayağın R, et al. Asymptomatic bacteriuria and antibacterial susceptibility patterns in an obstetric population. *ISRN Obstet Gynecol.* 2011;2011:721872. doi:10.5402/2011/721872
7. Ghafari M, Baigi V, Cheraghi Z, Doosti-Irani A. The prevalence of asymptomatic bacteriuria in Iranian pregnant women: a systematic review and meta-analysis. *PLoS One.* 2016;11(6):e0158031. doi:10.1371/journal.pone.0158031
8. Curtiss N, Meththananda I, Duckett J. Urinary tract infection in obstetrics and gynaecology. *Obstet Gynaecol Reprod Med.* 2017;27(9):261-265. doi:10.1016/j.ogrm.2017.06.006
9. Schneeberger C, Geerlings SE, Middleton P, Crowther CA. Interventions for preventing recurrent urinary tract infection during pregnancy. *Cochrane Database Syst Rev.* 2015(7):CD009279. doi:10.1002/14651858.CD009279.pub3
10. Ipe DS, Sundac L, Benjamin WH Jr, Moore KH, Ulett GC. Asymptomatic bacteriuria: prevalence rates of causal microorganisms, etiology of infection in different patient populations, and recent advances in molecular detection. *FEMS Microbiol Lett.* 2013;346(1):1-10. doi:10.1111/1574-6968.12204
11. Grabe M, Bartoletti R, Bjerkklund Johansen TE, et al. Guidelines on Urological Infections. *European Association of Urology (EAU);* 2015:182.
12. Jorde R, Sollid ST, Svartberg J, Joakimsen RM, Grimnes G, Hutchinson MY. Prevention of urinary tract infections with vitamin D supplementation 20,000 IU per week for five years. Results from an RCT including 511 subjects. *Infect Dis (Lond).* 2016;48(11-12):823-828. doi:10.1080/23744235.2016.1201853
13. Scholes D, Hooton TM, Roberts PL, Stapleton AE, Gupta K, Stamm WE. Risk factors for recurrent urinary tract infection in young women. *J Infect Dis.* 2000;182(4):1177-1182. doi:10.1086/315827
14. Amiri FN, Rooshan MH, Ahmady MH, Soliamani MJ. Hygiene practices and sexual activity associated with urinary tract infection in pregnant women. *East Mediterr Health J.* 2009;15(1):104-110.
15. Abdel-Aziz Elzayat M, Barnett-Vanes A, Dabour MF, Cheng F. Prevalence of undiagnosed asymptomatic bacteriuria and associated risk factors during pregnancy: a cross-sectional study at two tertiary centres in Cairo, Egypt. *BMJ Open.* 2017;7(3):e013198. doi:10.1136/bmjopen-2016-013198
16. Al-Badr A, Al-Shaikh G. Recurrent urinary tract infections management in women: a review. *Sultan Qaboos University Medical Journal.* 2013;13(3):359. doi: 10.12816/0003256
17. Ahmed AF, Solyman AA, Kamal SM. Potential host-related risk factors for recurrent urinary tract infection in Saudi women of childbearing age. *Int Urogynecol J.* 2016;27(8):1245-1253. doi:10.1007/s00192-016-2965-0
18. Mohamed NR, Omar H, Abd-Allah I. Prevalence and risk factors of urinary tract infection among pregnant women in Ismailia City, Egypt. *IOSR Journal of Nursing and Health Science.* 2017;6(3):62-72. doi:10.9790/1959-0603076272
19. Kisa S, Taskin L. Behavioral risk factors that predispose women to vaginal infections in Turkey. *Pak J Med Sci Q.* 2010;26(4):800-804.
20. Badran YA, El-Kashef TA, Abdelaziz AS, Ali MM. Impact of genital hygiene and sexual activity on urinary tract infection during pregnancy. *Urol Ann.* 2015;7(4):478. doi: 10.4103/0974-7796.157971
21. Sutaria P, Cholera M, Donga SB. A prevalence study of vaginal candidiasis among pregnant women. *Int J Adv Med.* 2019;6(3):922. doi:10.18203/2349-3933.ijam20192265
22. Farage MA, Miller KW, Sobel JD. Dynamics of the vaginal ecosystem--hormonal influences. *Infectious Diseases: Research and Treatment.* 2010;3:IDRT.S3903. doi:10.4137/idrt.s3903
23. Morgan KL. Management of UTIs during pregnancy. *MCN Am J Matern Child Nurs.* 2004;29(4):254-258.

doi:10.1097/00005721-200407000-00011

24. Enayat K, Fariba F, Bahram N. Asymptomatic bacteriuria among pregnant women referred to outpatient clinics in Sanandaj, Iran. *Int Braz J Urol.* 2008;34(6):699-704; discussion 704-697. doi:10.1590/s1677-55382008000600004
25. Ramzan M, Bakhsh S, Salam A, Khan GM, Mustafa G. Risk factors in urinary tract infection. *Gomal J Med Sci.* 2004;2(1):1-4.
26. Alijahan R, Pourfarzi F, Salimi S, Tahmasebi P, Moradi S, Hazrati S. Prevalence of urinary tract infection and associated risk factors in pregnant women. *Iran South Med J.* 2014;17(4):676-686. [Persian].
27. Fakhri M, Hamzehgardeshi Z, Hajikhani Golchin NA, Komili A. Promoting menstrual health among persian adolescent girls from low socioeconomic backgrounds: a quasi-experimental study. *BMC Public Health.* 2012;12:193. doi:10.1186/1471-2458-12-193
28. Najmabadi KM, Sharifi F. Sexual education and women empowerment in health: a review of the literature. *International Journal of Women's Health and Reproduction Sciences.* 2019;7(2):150-155. doi:10.15296/ijwhr.2019.25
29. Navidian A, Kykhaee A, Imani M, Taimoori B, Soltani P. The effect of group sexual counseling on the sexual response of pregnant women. *International Journal of Women's Health and Reproduction Sciences.* 2017;5(1):60-65. doi:10.15296/ijwhr.2017.11
30. Haider G, Zehra N, Munir AA, Haider A. Risk factors of urinary tract infection in pregnancy. *J Pak Med Assoc.* 2010;60(3):213-216.
31. Aseel MT, Al-Meer FM, Al-Kuwari MG, Ismail MFS. Prevalence and predictors of asymptomatic bacteriuria among pregnant women attending primary health care in Qatar. *Middle East Journal of Family Medicine.* 2009;7(4):10-13.
32. Obiobolu C, Okonko I, Anyamere C, et al. Incidence of urinary tract infections (UTIs) among pregnant women in Akwa metropolis, Southeastern Nigeria. *Sci Res Essays.* 2009;4(8):820-824.

© 2020 The Author (s); This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.