





## Evaluation of Anticancer and Neuroprotective Properties of Curcumin: a Network Analysis

Mohammad Rostami-Nejad<sup>1</sup> , Mohammadreza Razzaghi<sup>2</sup>, Somayeh Esmaeili<sup>3</sup>, Alireza Zali<sup>4</sup>, Majid Rezaei-Tavirani<sup>5</sup>, Mohammah Hossein Heidari<sup>6</sup>, Mostafa Rezaei-Tavirani<sup>6</sup>, Mona Zamanian-Azodi<sup>7\*</sup> , Nayebali Ahmadi<sup>6</sup>

<sup>1</sup>Gastroenterology and Liver Diseases Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

<sup>2</sup>Laser Application in Medical Sciences Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

<sup>3</sup>Traditional Medicine and Materia Medica Research Center and Department of Traditional Pharmacy, School of Traditional Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

<sup>4</sup>Functional Neurosurgery Research Center, Faculty of Medicine Shahid Beheshti University of Medical Sciences, Tehran, Iran.

<sup>5</sup>Firoozabadi Hospital, Faculty of Medicine, Iran University of Medical Sciences, Tehran, Iran.

<sup>6</sup>Proteomics Research Center, Faculty of Paramedical Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

<sup>7</sup>Proteomics Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

### Abstract

**Background and objectives:** Curcumin as a medicinal substance has shown effective in different kinds of diseases especially cancer. To understand its underlying mechanism, molecular complementary study of differentially expressed microRNAs (DEMs) could assist. In this view, regulatory network analysis of DEMs of melanoma cancer treated with curcumin versus the untreated male *Mus musculus* was investigated in this study. **Methods:** Data was obtained from the database of Gene Expression Omnibus (GEO), <https://www.ncbi.nlm.nih.gov/geo/>. At first, the log fold change (FC)  $\geq 2$  was assigned for predicting a cut off for DEMs in the following study. GEO2R detected a number of 250 top significantly changed microRNAs based on the priority of the most statistically significant ones. These miRNAs were then explored for regulatory network analysis via Cytoscape software v.3.7.2 and its plug-ins. **Results:** The findings indicated that a number of 21 miRNAs were statistically significant with differential expression amounts. Regulatory network also identified important microRNAs of mmu-miR-199a, mmu-miR-199b, mmu-miR-21, mmu-miR-142-3p, mmu-miR-148a, mmu-miR-214 and genes of Pkp3, Usp19, Ercc4, Ttc25, Atp13a2, Akr1b7, Umod, Nup188, Imp3, and Tmem74b. The highest ranked hub was mmu-miR-199a, which had nine connections. **Conclusion:** The present study offers new insights into the molecular mechanism of curcumin health benefits in melanoma cancer.

**Keywords:** curcumin; melanoma; microRNA

**Citation:** Rostami-Nejad M, Razzaghi M, Esmaeili S, Zali A, Rezaei-Tavirani M, Heidari MH, Rezaei-Tavirani M, Zamanian-Azodi M, Ahmadi N. Evaluation of anticancer and neuroprotective properties of curcumin: a network analysis. Res J Pharmacogn. 2020; 7(3): 85-92.

### Introduction

The biological activity of *Curcuma longa* L. (Zingiberaceae) known as turmeric has been

\* Corresponding author: mona.azodi@sbm.ac.ir

attributed to curcuminoids comprising of curcumin and two related compounds demethoxy curcumin and bisdemethoxycurcumin [1]. Curcumin as the mainly effective substance of turmeric has been famous for its many medical properties including anti-hepatotoxic, anti-microbial, anti-inflammatory, and anti-tumor advantages [2-4]. P53 upregulation and NF-kB, Akt/PI3K, and MAPK pathways inhibitions are the most prominent anticancer mechanisms which have been suggested for the functions of curcumin. On the other hand, microRNAs as potential regulatory targets in medicine are small non coding RNAs that have been reported playing role in different types of therapeutic approaches such as neural regeneration, heart health and cancer treatment [5-7]. In this sense, curcumin could also induce epigenetic changes in diseases such as cancer; for instance, down-regulation of miR-21, miR-17-5p, miR-20a, and miR-27a, and likewise up-regulation of tumor suppressor agents, miR-34 a/c and EMT (mesenchymal-epithelial transition) [8-10]. In addition, other studies have also showed that curcumin regulates miR-1, miR-7, miR-9, miR-34a, miR-181, miR-21, and miR-19 expressions in cancer [11]. This natural polyphenol substance has been also reported effective on melanoma as one of the hazardous type of cancers as well [12]. This type of skin cancer which effects on melanocytes has been accounted for yearly death of 55500 cases [13]. To identify the therapeutic molecular mechanism of curcumin influence, microarray analysis could assist. In a study by Indra N. Dahmke et al. microRNA expression profiling of curcumin treated mouse with melanoma was considered for the following bioinformatics research [14]. In silico analysis as a complementary study could provide more knowledge in this field by means of examining the regulatory network of differentially expressed microRNAs and their hub target genes. In this way, more information related to mechanisms by which curcumin provides health benefits could be reached. For this purpose, a regulatory network of curcumin influence on melanoma has been assessed.

## Materials and Methods

### Ethical considerations

This study was originated from the project with ethical code: IR.SBMU.REC.1398.154 approved by the ethical committee of Shahid Beheshti University of medical sciences.

## Data collection


The keywords, Curcumin + Melanoma were searched in GEO database and the corresponding data; the microRNA expression profiles which were included in GSE47211 were downloaded. The platform for this analysis is GPL16275 with 13 samples of controls and curcumin-treated individuals. The investigated organism is male *Mus musculus* in which the sampling was carried out from tumors for analyzing non-coding RNA profiling by array. Curcumin prescribed in this analysis was C3 complex that contains 77% curcumin, 17% demethoxycurcumin, and 3% bisdemethoxycurcumin. The samples were either dieting standard or with curcumin 4%= 160 mg daily. The comparison of the two samples was handled by GEO2R, the online analyzer (<http://www.ncbi.nlm.nih.gov/geo/geo2r/>).

Through this analysis, a list of 250 statistically significant expressed microRNAs were found and among them a cut off with  $\log_2 FC \geq 2$  and adjusted  $p\text{-value} \leq 0.05$  were designated to detect the DEMs. These candidates were then selected for further analysis known as “regulatory network analysis”. Cytoscape software 3.7.2 was used to construct and analyze the DEMs network and CluePedia presents a regulatory network of combined hub microRNAs and hub target genes [15,16]. There are certain criteria for statistical analysis of regulatory network, including number of hub genes that should be inserted in the regulatory network, which was 10; the miRanda score cut off that was assigned to miRanda-SCORE-v5  $\geq 0.6$ . The regulatory network analysis source was miRanda (miRNA-V5-2012-07-19.txt.gz). To get a better visualizations, the expression profiles of microRNA data, Heatmapper (<http://heatmapper.ca/expression/>) presentation was provided [17]. Furthermore, to get a better knowledge of corresponding hub genes, enrichment analysis was carried out by ClueGO [18]. The statistical criteria for biological process study were kappa score cut off  $\geq 0.5$  for grouping and gene number and percentage per term were assigned as 1 and 1, respectively. Bonferroni step down was the p-value correction method.

## Results and Discussion

To understand whether the groups are comparable in terms of microRNA expression profile, box plotting is carried out to get more insight. As it is shown in the figure 1, microRNA





**Figure 3.** The heat map presentation of the 18 microRNAs of regulatory network. The X axis refers to GSM series including GSM1146948-54 (7 rats as treated group) and GSM1146942-47 (6 rats as controls), right Y axis indicates microRNAs. Green and red colors code up-regulation and down-regulation respectively. Green to reds refers to the intermediate expression levels. Dendrogram is shown in the left Y axis which refers to the clustering of microRNAs.


**Table 1.** The list of hub microRNAs and the corresponded hub genes

R	mi-RNA ID	RC	K	FC	Hub genes
1	mmu-miR-199a	Up	9	4.98	Imp3, Tmem74b, Akr1b7, Umod, Ercc4, Pkp3, Usp19, Ttc25, Atp13a2
2	mmu-miR-199b	Up	6	4.42	Tmem74b, Akr1b7, Ercc4, Pkp3, Ttc25, Atp13a2
3	mmu-miR-21	Up	6	7.34	Umod, Nup188, Ercc4, Pkp3, Usp19, Atp13a2
4	mmu-miR-142	Up	5	4.89	Tmem74b, Akr1b7, Umod, Ttc25, Atp13a2
5	mmu-miR-148a	Up	5	4.37	Imp3, Umod, Nup188, Ercc4, Pkp3
6	mmu-miR-214	Up	5	4.37	Umod, Nup188, Ercc4, Pkp3, Usp19

The cut off assigned for hub microRNAs is  $\geq 5$ . R: Row; RC: regulation condition; K: degree; FC; fold change; Up refers to up-regulation

MicroRNAs are the important regulators of gene expression that could play a role in disease pathogenesis. Treatments with natural polyphenols such as curcumin could have some effects on culprit microRNAs and leading them to the normal function [14,19]. Melanoma cancer as one of the most serious kinds of skin cancers has been examined for health benefits of curcumin via microarray study [14]. DEMs were detected in the original research and in our study,

a regulatory network of these elements was investigated. GEO2R analysis implied that the samples of control and treated ones were comparable in terms of expression profile. In the next step, DEMs were assigned as 18 up-regulated and 3 down-regulated ones. Among 21 DEMs, 18 individuals were included in the main connected component of the constructed network which accepts one microRNA the other were upregulated.



**Figure 4.** Biological process analysis of 10 hub genes considering gene per term; 1, gene percentage; 1, and kappa score cut off  $\geq 0.5$ . The related genes are presented in boxes. Ttc25, Nup188, and Tmem74b were not related to the biological processes.

Heatmap analysis showed expression pattern clearly separates samples into two classes of control and treated groups. Among 18 microRNA, mmu-miR-199a, mmu-miR-199b, mmu-miR-21, mmu-miR-142, mmu-miR-148a, and mmu-miR-214 were introduced as hub microRNA. As it is depicted in the table 1 the 6 introduced hub microRNA are associated to all 10 genes and mmu-miR-199a has maximum connections.

The potent hub; mmu-miR-199a is connected to all gene target except Nup188. J Zhou et al. reported that mmu-mir-199a plays a role in the regulation of metastatic genes in the melanoma cells [20]. They showed that in the miR-199a-5p-silenced versus overexpressing cells of studied mice, Cd44, Cdh1, Cxcr4, Etv4, Fxyd5, Rpsa, Mmp3, Myc, Rb1, Tcf20, Hprt1, Actb1 were up-regulate while Ctsk, Itga7, and Tnfsf10 were down-regulated. FC>2 was considered to determine the deregulated genes. Involvement in different types of pathological conditions is attributed to the mmu-miR-199 microRNA family such as; progress of liver fibrosis,

impaired mitochondria fatty acid oxidation, and oncogenic activity [21-23].

The changes of microRNAs could have some impact on biological processes via changing the regulation of target genes [24]. As it is shown in figure 4, among 10 target genes; 7 individuals including Imp3, Akr1b7, Umod, Ercc4, Pkp3, Usp19, and Atp13a2 are connected to the associated biological processes. Ercc4 is connected to the cluster of BP which includes most biological terms. There are several reports that Ercc4 mutation is associated to DNA repair defect in various types of diseases such as Fanconi anemia, xeroderma pigmentosum, and cancers [25-28].

ATP13a2 is the other important target gene that is connected to the “polyamine transmembrane transport” cluster of biological processes. Investigations have revealed neuroprotective role of ATP13a2. It is reported that ATP13a2 deficiency leads to neurotoxicity [29]. Prominent role of ATP13a2 deregulation in Parkinson disease is reported by A Di Fonzo et al. [30].

Umod and pcp3 are the two target genes that are related to the five hub microRNA. Highlighted role of Umod deregulation in the kidney disease and hypertension has been investigated by researchers [31,32]. It is emphasized that deregulation of Pcp3 leads to various types of cancers such as breast and gastric cancers [33,34]. As it is discussed, the gene expression study and also gene ontology evaluation revealed the anticancer and also neuroprotective properties of curcumin. It can be concluded that curcumin consumption is associated to the protection of body against important spectrum of diseases such as kidney, skin, vascular and neurodegenerative diseases, and cancer.

### Acknowledgments

Shahid Beheshti University of Medical Sciences supported this research.

### Author contributions

All of the authors participated in the design, performing the project, and editing of the manuscript. Mona Zamanian-Azodi provided the draft of the manuscript. Mostafa Rezaei-Tavirani edited the final version of the manuscript.

### Declaration of interest

The authors declare that there is no conflict of interest. The authors alone are responsible for the accuracy and integrity of the paper content.

### References

- [1] Amalraj A, Pius A, Gopi S, Gopi S. Biological activities of curcuminoids, other biomolecules from turmeric and their derivatives - a review. *J Tradit Complement Med.* 2017; 7(2): 205-233.
- [2] Evance WC. Trease and Evance pharmacognosy. 16<sup>th</sup> ed. Edinburgh: Elsevier, 2009.
- [3] Hewlings SJ, Kalman DS. Curcumin: a review of its' effects on human health. *Foods.* 2017; 6(10): 1-11.
- [4] Xu XY, Meng X, Li S, Gan RY, Li Y, Li HB. Bioactivity, health benefits, and related molecular mechanisms of curcumin: current progress, challenges, and perspectives. *Nutrients.* 2018; 10(10): 1-33.
- [5] Chen Y, Cao S, Xu P, Han W, Shan T, Pan J, Lin W, Chen X, Wang X. Changes in the expression of miR-34a and its target genes following spinal cord injury in rats. *Med Sci Monit.* 2016; 22: 3981-3993.
- [6] Müller S, Janke F, Dietz S, Sülthmann H. Circulating MicroRNAs as potential biomarkers for lung cancer. *Recent Results Cancer Res.* 2020; 215: 299-318.
- [7] Pulido-Moran M, Moreno-Fernandez J, Ramirez-Tortosa C, Ramirez-Tortosa M. Curcumin and health. *Molecules.* 2016; 21(3): 1-22.
- [8] Slabáková E, Culig Z, Remšík J, Souček K. Alternative mechanisms of miR-34a regulation in cancer. *Cell Death Dis.* 2017; 8(10): 1-10.
- [9] Momtazi AA, Shahabipour F, Khatibi S, Johnston TP, Pirro M, Sahebkar A. Curcumin as a MicroRNA regulator in cancer: a review. *Rev Physiol Biochem Pharmacol.* 2016; 171: 1-38.
- [10] Sekhon K, Bucay N, Majid S, Dahiya R, Saini S. MicroRNAs and epithelial-mesenchymal transition in prostate cancer. *Oncotarget.* 2016; 7(41): 1-15.
- [11] Mirzaei H, Masoudifar A, Sahebkar A, Zare N, Sadri Nahand J, Rashidi B, Mehrabian E, Mohammadi M, Mirzaei HR, Jaafari MR. MicroRNA: a novel target of curcumin in cancer therapy. *J Cell Physiol.* 2018; 233(4): 3004-3015.
- [12] Mirzaei H, Naseri G, Rezaee R, Mohammadi M, Banikazemi Z, Mirzaei HR, Salehi H, Peyvandi M, Pawelek JM, Sahebkar A. Curcumin: a new candidate for melanoma therapy? *Int J Cancer.* 2016; 139(8): 1683-1695.
- [13] Schadendorf D, van Akkooi ACJ, Berking C, Griewank KG, Gutzmer R, Hauschild A, Stang A, Roasch A, Ugurel S. Melanoma. *Lancet.* 2018; 392(10151): 971-984.
- [14] Dahmke IN, Backes C, Rudzitis-Auth J, Laschke MW, Leidinger P, Menger MD, Meese E, Mahlknecht U. Curcumin intake affects miRNA signature in murine melanoma with mmu-miR-205-5p most significantly altered. *PLoS One.* 2013; 8(12): 1-10.
- [15] Smoot ME, Ono K, Ruscheinski J, Wang P-L, Ideker T. Cytoscape 2.8: new features for data integration and network visualization. *Bioinformatics.* 2011; 27(3): 431-442.
- [16] Bindea G, Galon J, Mlecnik B. CluePedia Cytoscape plugin: pathway insights using

- integrated experimental and in silico data. *Bioinformatics*. 2013; 29(5): 661-663.
- [17] Babicki S, Arndt D, Marcu A, Liang Y, Grant JR, Maciejewski A, Wishart DS. Heatmapper: web-enabled heat mapping for all. *Nucleic Acids Res*. 2016; 44: 147-153.
- [18] Bindea G, Mlecnik B, Hackl H, Charoentong P, Tosolini M, Kirilovsky A, Fridman WH, Pages F, Trajanoski Z, Galon J. ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation networks. *Bioinformatics*. 2009; 25(8): 1091-1093.
- [19] Norouzinia M, Azodi MZ, Seyfi DN, Kardan A, Naseh A, Akbari Z. Predication of hub target genes of differentially expressed microRNAs contributing in *Helicobacter Pylori* infection in gastric non-cancerous tissue. *Gastroenterol Hepatol Bed Bench*. 2019; 12(S1): 44-50.
- [20] Zhou J, Liu R, Wang Y, Tang J, Tang S, Chen X, Xia K, Xiong W, Xu D, Wang S, He Q, Cao K. Mir-199a-5p regulates the expression of metastasis-associated genes in B16F10 melanoma cells. *Int J Clin Exp Path*. 2014; 7(10): 7182-7190.
- [21] Murakami Y, Toyoda H, Tanaka M, Kuroda M, Harada Y, Matsuda F, Tajima A, Kosaka N, Ochiai T, Shimotohni K. The progression of liver fibrosis is related with overexpression of the miR-199 and 200 families. *PLoS One*. 2011; 6(1): e16081.
- [22] El Azzouzi H, Leptidis S, Dirx E, Hoeks J, van Bree B, Brand K, McClellan EA, Poels E, Sluimer JC, van den Hoogenhof MM, Armand AS, Yin X, Langley S, Bourajjaj M, Olieslagers S, Krishnan J, Vooijs M, Kurihara H, Stubbs A, Pinto YM, Krek W, Mayr M, da Costa Martins PA, Schrauwen P, De Windt LJ. The hypoxia-inducible microRNA cluster miR-199a~ 214 targets myocardial PPAR $\delta$  and impairs mitochondrial fatty acid oxidation. *Cell Metab*. 2013; 18(3): 341-354.
- [23] Kim S, Lee UJ, Kim MN, Lee E-J, Kim JY, Lee MY, Choung S, Kim YJ, Choi YC. MicroRNA miR-199a\* regulates the MET proto-oncogene and the downstream extracellular signal-regulated kinase 2 (ERK2). *J Biological Chem*. 2008; 283(26): 18158-18166.
- [24] Philippidou D, Schmitt M, Moser D, Margue C, Nazarov PV, Muller A, Vallar L, Nashan D, Behrmann I, Kreis S. Signatures of microRNAs and selected microRNA target genes in human melanoma. *Cancer Res*. 2010; 70(10): 4163-4173.
- [25] Bogliolo M, Schuster B, Stoepker C, Derkunt B, Su Y, Raams A, Trujillo JP, Minguillón J, Ramírez MJ, Pujol R, Casado JA, Baños R, Rio P, Knies K, Zúñiga S, Benítez J, Bueren JA, Jaspers NG, Schärer OD, de Winter JP, Schindler D, Surrallés J. Mutations in ERCC4, encoding the DNA-repair endonuclease XPF, cause Fanconi anemia. *Am J Hum Genet*. 2013; 92(5): 800-806.
- [26] Biggerstaff M, Szymkowski DE, Wood RD. Co-correction of the ERCC1, ERCC4 and xeroderma pigmentosum group F DNA repair defects in vitro. *EMBO J*. 1993; 12(9): 3685-3692.
- [27] Kornguth DG, Garden AS, Zheng Y, Dahlstrom KR, Wei Q, Sturgis EM. Gastrostomy in oropharyngeal cancer patients with ERCC4 (XPF) germline variants. *Int J Radiat Oncol Biol Phys*. 2005; 62(3): 665-671.
- [28] Milne RL, Ribas G, González-Neira A, Fagerholm R, Salas A, González E, Dopazo J, Nevanlinna H, Robledo M, Benitez J. ERCC4 associated with breast cancer risk: a two-stage case-control study using high-throughput genotyping. *Cancer Res*. 2006; 66(19): 9420-9427.
- [29] Usenovic M, Tresse E, Mazzulli JR, Taylor JP, Krainc D. Deficiency of ATP13A2 leads to lysosomal dysfunction,  $\alpha$ -synuclein accumulation, and neurotoxicity. *J Neurosci*. 2012; 32(12): 4240-4246.
- [30] Di Fonzo A, Chien H, Socal M, Giraudo S, Tassorelli C, Iliceto G, Fabbrini G, Marconi R, Fincati E, Abbruzzese G, Marini P, Squitieri F, Horstink MW, Montagna P, Libera AD, Stocchi F, Goldwurm S, Ferreira JJ, Meco G, Martignoni E, Lopiano L, Jardim LB, Oostra BA, Barbosa ER, Italian Parkinson Genetics Network, Bonifati V. ATP13A2 missense mutations in juvenile parkinsonism and young onset Parkinson disease. *Neurology*. 2007; 68(19): 1557-1562.
- [31] Hart T, Gorry M, Hart P, Woodard A, Shihabi Z, Sandhu J, Shirts B, Xu L, Zhu H, Barmada MM, Bleyer AJ. Mutations of the UMOD gene are responsible for

medullary cystic kidney disease 2 and familial juvenile hyperuricaemic nephropathy. *J Med Genet.* 2002; 39(12): 882-892.

- [32] Padmanabhan S, Melander O, Johnson T, Di Blasio AM, Lee WK, Gentilini D, Hastie CE, Menni C, Monti MC, Delles C, Laing S, Corso B, Navis G, Kwakernaak AJ, van der Harst P, Bochud M, Maillard M, Burnier M, Hedner T, Kjeldsen S, Wahlstrand B, Sjögren M, Fava C, Montagnana M, Danese E, Torffvit O, Hedblad B, Snieder H, Connell JM, Brown M, Samani NJ, Farrall M, Cesana G, Mancina G, Signorini S, Grassi G, Eyheramendy S, Wichmann HE, Laan M, Strachan DP, Sever P, Shields DC, Stanton A, Vollenweider P, Teumer A, Völzke H, Rettig R, Newton-Cheh C, Arora P, Zhang F, Soranzo N, Spector TD, Lucas G, Kathiresan S, Siscovick DS, Luan J, Loos RJ, Wareham NJ, Penninx BW, Nolte IM, McBride M, Miller WH, Nicklin SA, Baker AH, Graham

D, McDonald RA, Pell JP, Sattar N, Welsh P; Global BPgen Consortium, Munroe P, Caulfield MJ, Zanchetti A, Dominiczak AF. Genome-wide association study of blood pressure extremes identifies variant near UMOD associated with hypertension. *PLoS Genet.* 2010; 6(10): 1-11.

- [33] Demirag GG, Sullu Y, Yucel I. Expression of Plakophilins (PKP1, PKP2, and PKP3) in breast cancers. *Med Oncol.* 2012; 29(3): 1518-1522.

- [34] Demirag GG, Sullu Y, Gurgenyatagi D, Okumus NO, Yucel I. Expression of plakophilins (PKP1, PKP2, and PKP3) in gastric cancers. *Diagn Pathol.* 2011; 6(1): 1-5.

### Abbreviations

DEMs: differentially expressed microRNAs; FC: fold change; GEO: Gene Expression Omnibus; RC: regulation condition; K: degree