



Effect of Reminiscence Therapy on Death Anxiety in Retired Elderly Men: An Interventional Study

Abbasali Sheykhi¹, Farshid Saeedinezhad¹, Zahra Raiesi² and Nasrin Rezaee^{3,*}

¹Medical-Surgical Nursing Department, School of Nursing and Midwifery, Zahedan University of Medical Sciences, Zahedan, Iran

²Baharan Psychiatric Hospital, Zahedan University of Medical Science, Zahedan, Iran

³Community Nursing Research Center, Zahedan University of Medical Sciences, Zahedan, Iran

*Corresponding author: School of Nursing and Midwifery, Mashahir Sq, Postal Code: 9716913396, Zahedan, Iran. Tel: +98-5433442481, Email: nasrin_rezaee2005@yahoo.com

Received 2020 April 25; Accepted 2020 April 25.

Abstract

Background: Death anxiety is one of the concerns of the elderly population in the late stages of life. In this context, recalling memories is one of the favorite characteristics of these people that can affect different aspects of their mental and social health.

Objectives: This study aimed to determine the effect of reminiscence therapy on the death anxiety of men retired from the Islamic Republic of Iran Army living in Zahedan.

Methods: A quasi-experimental study was performed on 90 retired men over 60 years of age. The subjects were recruited through convenience sampling at the Army Retirement Center in Zahedan in winter 2019. The participants were randomly divided into the intervention (n = 45) and control (n = 45) groups. In the intervention group, reminiscence therapy was carried out based on major life events twice a week for six sessions. Data were gathered again two months after the intervention. The control group did not take part in any program. Data collection tools included a demographic form and the Death Anxiety scale (Templer, 1970). The results were analyzed by SPSS-21 using descriptive and inferential statistics. The P values < 0.05 were considered statistically significant.

Results: The mean scores of death anxiety were 25.37 ± 1.89 and 25.17 ± 1.61 in the intervention and control groups, respectively. After the experiment, this score was 9.09 ± 1.79 in the intervention group and 25.04 ± 1.62 in the control group. Thus, while reminiscence therapy significantly relieved death anxiety in the intervention group ($P < 0.0001$), the control group showed no significant difference in this respect at the end of the study ($P = 0.63$).

Conclusions: Group reminiscence therapy can reduce the death anxiety of older adults. Since the elderly are fond of expressing their memories, and such storytelling is an attractive, simple, and inexpensive intervention, it could be used to help these people mitigate their death anxiety.

Keywords: Death Anxiety, Older Adults, Reminiscence Therapy

1. Background

Aging is an inevitable biological process accompanied by both pleasant and unpleasant experiences (1). Social, economic, and scientific developments in recent years have led to higher life expectancy and lower mortality rates. The resulting population growth is far greater in developing countries than in developed countries. Iran, a developing country, is no exception to these demographic changes (2). As the number of older adults grows rapidly, the issue of their hygiene, health, and well-being in societies becomes ever more complicated and involves new areas. Aging is associated with both physical changes such as heart diseases and psychological changes like death anxiety (3).

Death anxiety is one of the main issues that older peo-

ple deal with. Death is an inevitable phenomenon, and thinking about it has played a decisive role in human life since the beginning of history. Some people see death as one stage of life, but some others regard it as the end of life (4). Death anxiety is a latent sentiment that makes one feel the taste of death in everything. Death anxiety is normal, but if it is too severe, it weakens one's performance (3) and negatively affects physical and mental health, especially in the elderly who feel it approaching after retirement (4). Death anxiety refers to the fear of dying, being cut off from the world, or dreading what happens afterward. While death anxiety is a single experience, people react to it differently. Thus, individual differences such as age, gender, occupation, and the experience of death, along with its confrontation, can affect the level of death anxiety one might have (5). In other words, people's attitude to death is

shaped both consciously and unconsciously by individual, cultural, social, and philosophical variables (6).

Retirement is a formative transition in an older person's life, and it entails important changes in lifestyle and social roles. There is no predetermined age for retirement, but it often comes after old age. The World Health Organization has proposed 60 - 65 years as a convenient yardstick to mark old age. Coupled with retirement, old age gives rise to several losses, including function, friends, and freedom, that might compromise one's mental health (5).

However, studies suggest that army retirees have experienced differently from other retired people, which can influence their physical and mental health. A study of more than 800 military retirees indicated that over half of these individuals had poor mental health (6). The findings from a cohort study addressing factors affecting the mental health of army retirees in 2001 - 2011 demonstrated that these older adults were more likely to experience psychological disorders such as depression and insomnia (7). In addition, the results of another study underscored that, despite the passage of many years since their experience in the Vietnam war, veterans still referred to medical centers because of psychological symptoms (8).

The lack of health can drive a person to social isolation (9). Mehri Nejad et al. (10) reported a significant relationship between social support and death anxiety in the elderly population. The results of another study in Turkey confirmed a significant relationship between death anxiety and depression in older adults living at a nursing home (11).

There is a variety of non-pharmacological therapies for old people with psychological problems such as anxiety, depression, and social isolation. Most notably, one may point to psychological interventions like cognitive therapy, narrative therapy, problem-solving therapy, bibliotherapy, movie therapy, and reminiscence therapy. In this context, reminiscence therapy can positively affect different aspects of social and mental health in the elderly. Musavi et al. (12) observed that reminiscence therapy could influence mental health, especially anxiety, of old people.

Reminiscence therapy is an intervention often used for older people. In the area of nursing interventions, it is used to evoke past events, feelings, and thoughts to promote mental and social health by establishing a happy and active atmosphere. Reminiscence therapy is an attractive, preventive, and therapeutic intervention for the elderly (12). People feel more valued by remembering memories. Indeed, those who successfully recreate their life story will acquire a sense of coherence (3). The results of a meta-analysis of clinical trials emphasized that reminiscence therapy should be integrated into routine care programs designed for older adults with depressive symptoms (13).

recalling memories reveals an individual's philosophy of life and the purpose he/she attaches to living. It helps represent the meaning of life and provides a framework for people's beliefs about this meaning, which also includes the concept of death. Older adults can release their daily anxieties by telling their past stories (3). Reminiscence therapy has patterns such as reviewing life based on different periods (2) or important life events (14).

Few studies have explored the impact of reminiscence therapy on depression and anxiety in older adults, and such studies have even been conducted on different populations (8, 12, 13, 15). The results of a systematic review, which proposed reminiscence therapy as a non-pharmacological intervention with many benefits, confirmed that group reminiscence therapy affected older people's depression, yet the authors also noted the need for further research into the effects of this therapeutic approach on anxiety and loneliness (16). Meanwhile, the results of a study conducted in a nursing home in Korea demonstrated that reminiscence therapy is beneficial for older adults with death anxiety (17). Iran's population is aging, and many old people are retired. Retirees have different experiences that can affect their psychological problems, such as death anxiety.

2. Objectives

The present study attempted to explore the impact of reminiscence therapy on the death anxiety of army retirees living in Zahedan in 2019.

3. Methods

We performed a quasi-experimental study in 2019 on 90 elderly people retired from the Islamic Republic of Iran Army. The study population consisted of all men over 60 years of age who referred to the Army Retirement Center in Zahedan. Convenience sampling was used to recruit eligible retirees, who were then randomized into the intervention and control groups. The grouping was done randomly by asking each person to choose a card, on which was written the name of each of the two groups. The number of cards equaled the sample size.

The inclusion criteria were men aged 60 - 75 years, relatively good physical and mental health to take part in the study (based on the statement of participant's family), acceptable hearing (based on participant's statement), a cognitive assessment score of higher than 7 (based on the Abbreviated Mental Test score (AMTS)), willingness to participate, and the passage of at least six months since retirement. On the other hand, the exclusion criteria were fail-

ure to attend more than one session, the occurrence of severe family crisis during the study (e.g., loss of a family member), and intolerance to the group.

The sample size was estimated at 14 for each group based on the mean total score of death anxiety reported by Jo and An (17), a confidence level of 95%, and statistical power of 95%, as well as the mean scores of $\bar{X}_1 = 56.06$ in the control group and $\bar{X}_2 = 52.67$ in the experimental group and standard deviations of $S_1 = 1.92$ and $S_2 = 2.95$ respectively. To ensure the adequacy of sample size and consider possible attrition, we finally allocated 45 individuals to each of the two groups (total = 90).

Data were collected using a self-report questionnaire consisting of two parts. The first section included personal and clinical information, age, education, retirement duration, number of children, post-retirement occupation, and AMTS. The AMTS scale has 10 questions designed to screen cognitive disorders in the elderly. The correct and incorrect answers to each question are scored 1 and 0, respectively, and the total score (ranging from 1 to 10) is the sum of correct answers. This instrument has been psychometrically tested in different countries. In Spain, it had a sensitivity of 100%, a specificity of 53%, and a cutoff point of 7 - 8. In Iran, the AMTS has been psychometrically evaluated, and its cutoff point has been reported at 7 (18). In the second part of data collection, we used the Death Anxiety scale (Templer, 1970) to assess participants' death anxiety. This questionnaire comprises 15 items that are scored based on a five-point Likert scale from 1 "strongly disagree" to 5 "strongly agree" (19). Questions 2, 3, 6, 7, and 15 give scores in the reverse order. The minimum and maximum scores of this instrument are 15 and 75, with higher scores indicating higher levels of death anxiety. This questionnaire has been psychometrically evaluated in Iran. The reliability (Cronbach's alpha) and internal consistency of the Death Anxiety scale have been appropriate (20). In our study, the reliability of this questionnaire was confirmed with Cronbach's alpha of 0.82.

We first examined the characteristics of those retirees who met some of the inclusion criteria (age and retirement duration) and found their phone numbers from their records. Then, we contacted these people and recruited those who had the rest of the inclusion criteria and were willing to participate in the study. In the next stage, they were randomly divided into the control and intervention groups. The subjects provided their written informed consent. The AMTS scale was completed for all of them before group allocation.

The subjects were then invited to the Army Retirement Center. The questionnaires were completed on the first day, and the dates for holding group sessions of 5 - 8 people

were determined. Overall, six reminiscence therapy sessions were held twice a week for three weeks. Each session lasted 45 to 60 minutes. The questionnaires were filled out again two months after the end of the intervention. At all sessions, the researcher, a psychiatric nurse, had the role of a leader. In other words, he would invite all individuals to participate in discussions and direct their attention to the main subject matter whenever a person diverted the discussion into current political or economic issues. The research assistant also attended all sessions to assess and care for the physical condition of elderly participants. The content presented in the sessions was based on the standards of reminiscence therapy, including attractiveness, and the characteristics of participants. More precisely, we addressed important life events like major (non-)occupational and post-retirement experiences (Table 1). To manage the time of the sessions, we gave each person an average of 5 - 7 minutes to talk in each session. The control group did not receive any intervention during this period, yet they underwent the post-test at the same time as the intervention group. At the end of the study, participants in the control group were contacted and invited to a reminiscence therapy session at the center. Some but not all of the control group members accepted the invitation.

The obtained data were analyzed in SPSS-21 using descriptive statistics (Shapiro-Wilk test) and inferential statistics (independent *t*-test, paired *t*-test, and chi-squared test) at the significance level of less than 0.05.

4. Results

Eventually, 90 participants completed the study. The mean age of the study subjects was 68 years in both intervention and control groups, and no significant difference was found in this regard ($P = 0.47$). Over 57% of the intervention group had a post-retirement job, whereas 42% of the control group reported such an occupation; nevertheless, the difference was not significant ($P = 0.14$) (Table 2).

The results showed that the mean score of death anxiety after reminiscence therapy decreased significantly in the intervention group, but this change was not significant in the control group. The results indicated no significant difference in the baseline scores of death anxiety between the two groups. After reminiscence therapy, however, the two groups exhibited a significant difference, with the intervention group reporting a lower score of death anxiety (Table 3).

5. Discussion

The findings of the present study highlight the significant impact of reminiscence therapy on death anxiety in

Table 1. Structure of Reminiscence Therapy Sessions

Session	Content
1. Influential life events	Creating a friendly atmosphere for the expression of memories
	recalling a decisive life event
	Illustration: A major life event is one that you have experienced and has significantly changed your life.
2. Family history	A brief review of memories narrated in the previous session
	Illustration: Who are your family members, or who have had a positive or negative impact on your life? who helped you most in your life and contributed to your progress?
3. Occupational memories	A brief review of memories narrated in the previous session
	Discussing occupational memories
	Illustration: How has your work influenced your roles in society and the family?
4. Stressful experiences	A brief review of memories narrated in the previous session
	Illustration: Love is an emotional attachment to a person, place, or something special. What are the major loves of your life? have you ever fallen in love or hated someone?
5. Meaning and purpose of life	A brief review of memories narrated in the previous session
	Illustration: Have you achieved any of your life goals? have you strived for the goals you have not reached?
6. Summing up	Evaluating memories
	Illustration: Pleasant and unpleasant memories

Table 2. Demographic Characteristics of Army Retirees in the Two Study Groups^a

Variable	Intervention Group	Control Group	Result
Age	68.02 ± 3.46	68.64 ± 4.60	P = 0.47 ^b
Number of children	2.35 ± 1.22	2.53 ± 1.32	P = 0.51 ^b
Education			P = 0.37 ^c
High school diploma	11 (24.4)	16 (35.6)	
Associate degree	13 (28.9)	14 (31.1)	
Bachelor's degree or higher	21 (46.7)	15 (33.3)	
Post-retirement job			P = 0.14 ^c
Yes	26 (57.8)	19 (42.2)	
No	19 (42.2)	26 (57.8)	

^aValues are expressed as mean ± SD or No. (%).

^bIndependent t-test.

^cChi-square test.

older adults. In the same vein, Ataie Moghanloo et al. (21) found that integrative reminiscence therapy focusing on cohesive memory could reduce anxiety in old widowers. Likewise, the results of a study conducted in Korea by Jo and An (17) supported the reminiscence therapy effect on the anxiety of older adults living at nursing homes.

In the present study, the opportunity was provided

to reduce the psychological burden of death anxiety in older adults by allowing them to interact with their peers through sharing their memories. The results of the study by Sadri Damirchi et al. (22), which examined the impact of reminiscence therapy on old women, displayed that older people who had a meaning and purpose in their life were less afraid of death and more likely to accept it. In addition, Moetamedi et al. (23) observed that life review therapy introduced consistency in the personality of the elderly. In our study, the elderly found the meaning of their life by telling their memories and reviewing their past. The final stage of Erik Erikson's theory (i.e., ego integrity versus despair) states that the elderly look at their past life and re-examine its positive and negative aspects (24), which was made possible in our study through reminiscence therapy. Indeed, by recollecting their memories, our study subjects developed a feeling of content and satisfaction, which, in turn, reduced their death anxiety.

Similarly, Moetamedi et al. (23) suggested that group therapy could reduce the death anxiety of older adults. The results of some studies support that group reminiscence therapy can also increase the life expectancy of the elderly (8, 16, 25). Thus, group therapies can help this population understand the meaning of their life. Such interventions seem to be more appropriate in terms of social support, coping with grief, losses of any kind, dealing with chronic illnesses, and ultimately acceptance of death (26). In line with our study, Garrow and Walker (26) concluded that group therapy created social cohesion in the elderly and

Table 3. Comparison of Participants' Death Anxiety on the Two Groups Before and After the Intervention^a

Stage Group	Before Intervention	After Intervention	Changes	Paired t-test
Intervention	25.37 ± 1.89	9.09 ± 1.79	16.28 ± 1.94	0.0001 ^b
Control	25.17 ± 1.61	25.04 ± 1.62	0.13 ± 1.87	0.63 ^b
Independent t-test	0.59 ^c	0.01 ^c	0.005 ^c	

^aValues are expressed as mean ± SD.^bPaired t-test.^cIndependent t-test.

provided conditions for them not to feel lonely; this homogeneity helps reduce anxiety.

Death anxiety is one of the components of mental health, yet older people experience high levels of this anxiety. The findings by Ghorbanalipour et al. (27) show that people with high death anxiety may especially benefit from health-promoting behaviors. Moreover, the results of the study by Soleimani et al. (28) emphasized that death anxiety is a predictor of poor quality of life. On the other hand, Siverova and Buzgova (29) studied older adults at a senior care center and noted that reminiscence therapy positively affected the quality of life and the attitude to aging and reduced the symptoms of depression in this population. These findings corroborate that reminiscence therapy is interesting to the elderly and it will be more durable and satisfying when conducted in a group setting.

While most related studies on older people have been undertaken in senior care centers, Chueh and Chang (30) investigated the effect of reminiscence therapy on veterans' depressive symptoms in Taiwan and observed that these symptoms declined three months and six months after the intervention. It could be concluded that reminiscence therapy, in the long run, can also have a positive impact on the mood of army retirees who have experienced differently from those of other elderly people.

One of the limitations of this study concerns the physical health status of participants, such that in some cases, their relatives were reluctant to leave them alone. Consequently, these companions were also allowed to attend the sessions, which was beyond our control.

5.1. Conclusions

The results of our study demonstrated that group reminiscence therapy could alleviate the death anxiety of older adults. Thanks to its great appeal, this method might be used to enable these people to reexamine different aspects of their lives. In fact, by encouraging older adults to look back on their life, this intervention can provide them with a sense of satisfaction, which, in turn, mitigates their death anxiety. It is suggested that this easy-to-use therapy be implemented in retirement centers and nursing homes.

Acknowledgments

This article is part of a Master's thesis in Psychiatric Nursing presented at Zahedan University of Medical Sciences. We would like to express our gratitude to all the seniors who participated in this research.

Footnotes

Authors' Contribution: Nasrin Rezaee did supervision, study design, and manuscript composition. Farshid Saeedinezhad did study design and data analysis. Abbasali Sheykhi did study design and data collection. Zahra Raiesi did study design and data collection.

Conflict of Interests: None of the authors has any conflict of interest to declare.

Ethical Approval: This study was approved by the Ethics Committee of Zahedan University of Medical Sciences (code: IR.ZAUMS.REC.1397.379).

Funding/Support: No financial support was received for this research.

Informed Consent: All participants gave their informed consent.

References

1. Fakari FR, Azimi Hashemi M, Rashidi Fakari F. A qualitative research: Postmenopausal women's experiences of abuse. *Proced Soc Behav Sci*. 2013;**82**:57-60. doi:10.1016/j.sbspro.2013.06.224.
2. Yazdanbakhsh K. Effects of life review on social adjustment of retirees' elderly persons. *J Aging Psychol*. 2015.
3. Majidi A, Moradi O. Effect of teaching the components of spiritual intelligence on death anxiety in the elderly. *Salmand*. 2018;**13**(1):110-23. doi:10.21859/sija.13.1.110.
4. Keshavarz-Afshar H, Jahan-Bakhshi Z, Anisi J, Azad-Marzabadi E, Ghahvehchi-Hosseini F. The determination of mental health among military retired forces based on organizational and humanistic variables. *J Mil Med*. 2016;**18**(2):191-6.
5. Rajabi S, Narimani M, Basari A. Evaluation and comparison of mental health statues a policemen and military personnel. *J Mil Psychol*. 2013;**4**(13):32-43.
6. Jahangiri MM. Prediction of death anxiety, according to religious orientation and loneliness in elderly in Arak City. *J Gerontol*. 2017;**1**(4):58-66. doi:10.18869/acadpub.joge.1.4.58.

7. Armenta RF, Rush T, LeardMann CA, Millegan J, Cooper A, Hoge CW, et al. Factors associated with persistent posttraumatic stress disorder among U.S. Military Service members and veterans. *BMC Psychiatry*. 2018;**18**(1):48. doi: [10.1186/s12888-018-1590-5](https://doi.org/10.1186/s12888-018-1590-5). [PubMed: 29452590]. [PubMed Central: PMC5816529].
8. Yousefi Z, Sharifi K, Tagharrobi Z, Akbari H. The effect of group reminiscence on happiness of the elderly. *Evidence Based Care*. 2014;**4**(3):33-46.
9. Ebrahimi L, Moradi F. Correlation of perceived social support and psychological hardness with perceived stress and aggression in retired veterans. *Iran J War Public Health*. 2018;**10**(3):157-64. doi: [10.29252/ijwph.10.3.157](https://doi.org/10.29252/ijwph.10.3.157).
10. Mehri Nejad SA, Ramezan Saatchi L, Paydar S. Death anxiety and its relationship with social support and adherence to religion in the elderly. *Salmand*. 2017;**11**(4):494-503. doi: [10.21859/sija-1104494](https://doi.org/10.21859/sija-1104494).
11. Bektas H, Korukcu O, Kabukcuoglu K. Undercover fear of elderly people in nursing homes: Death anxiety and depression. *J Hum Sci*. 2017;**14**(1). doi: [10.14687/jhs.v14i1.4268](https://doi.org/10.14687/jhs.v14i1.4268).
12. Musavi M, Mohammadian S, Mohammadinezhad B. The effect of group integrative reminiscence therapy on mental health among older women living in Iranian nursing homes. *Nurs Open*. 2017;**4**(4):303-9. doi: [10.1002/nop2.101](https://doi.org/10.1002/nop2.101). [PubMed: 29085656]. [PubMed Central: PMC5653382].
13. Song D, Shen Q, Xu TZ, Sun QH. Effects of group reminiscence on elderly depression: A meta-analysis. *Int J Nurs Sci*. 2014;**1**(4):416-22. doi: [10.1016/j.ijnss.2014.10.001](https://doi.org/10.1016/j.ijnss.2014.10.001).
14. Huang HC, Chen YT, Chen PY, Huey-Lan Hu S, Liu F, Kuo YL, et al. Reminiscence therapy improves cognitive functions and reduces depressive symptoms in elderly people with dementia: A meta-analysis of randomized controlled trials. *J Am Med Dir Assoc*. 2015;**16**(12):1087-94. doi: [10.1016/j.jamda.2015.07.010](https://doi.org/10.1016/j.jamda.2015.07.010). [PubMed: 26341034].
15. Abdollahzadeh H, Khabbazi M. Effectiveness of integrative reminiscence on health anxiety, death depression and perceived stress in elderly. *Psychol Health*. 2017;**6**(2):101-4.
16. Syed Elias SM, Neville C, Scott T. The effectiveness of group reminiscence therapy for loneliness, anxiety and depression in older adults in long-term care: A systematic review. *Geriatr Nurs*. 2015;**36**(5):372-80. doi: [10.1016/j.gerinurse.2015.05.004](https://doi.org/10.1016/j.gerinurse.2015.05.004). [PubMed: 26099638].
17. Jo KH, An GJ. Effects of a group reminiscence program on self-forgiveness, life satisfaction, and death anxiety among institutionalized older adults. *Korean J Adult Nurs*. 2018;**30**(5). doi: [10.7475/kjan.2018.30.5.546](https://doi.org/10.7475/kjan.2018.30.5.546).
18. Foroughan M, Wahlund LO, Jafari Z, Rahgozar M, Farahani IG, Rashedi V. Validity and reliability of Abbreviated Mental Test score (AMTS) among older Iranian. *Psychogeriatrics*. 2017;**17**(6):460-5. doi: [10.1111/psyg.12276](https://doi.org/10.1111/psyg.12276). [PubMed: 28589659].
19. Templer DI. The construction and validation of a Death Anxiety scale. *J Gen Psychol*. 1970;**82**(2d Half):165-77. doi: [10.1080/00221309.1970.9920634](https://doi.org/10.1080/00221309.1970.9920634). [PubMed: 4394812].
20. Sharif Nia H, Ebadi A, Lehto RH, Mousavi B, Peyrovi H, Chan YH. Reliability and validity of the persian version of templer death anxiety scale-extended in veterans of Iran-Iraq warfare. *Iran J Psychiatry Behav Sci*. 2014;**8**(4):29-37. [PubMed: 25798171]. [PubMed Central: PMC4364474].
21. Ataie Moghanloo V, Ataie Moghanloo R, Pishvaei M, Goudarzian M. Treatment efficacy of reminders of life with emphasis on integrative reminiscence on self-esteem and anxiety of old widowers. *J Sabzevar Univ Med Sci*. 2016;**23**(3):422-9. doi: [10.21859/sums-2303422](https://doi.org/10.21859/sums-2303422).
22. Sadri Damirchi E, Ghomi M, Esmaeli Ghazi Valoii F. Effectiveness of life review group therapy on psychological well-being and the life expectancy of elderly women. *Salmand*. 2017;**12**(3):312-25. doi: [10.21859/sija.12.3.312](https://doi.org/10.21859/sija.12.3.312).
23. Moetamedi A, Pajouhinia S, Fatemi Ardestani MH. The impact of spiritual wellbeing and resiliency in predicting death anxiety among elderly people in Tehran. *Neurosci J Shefaye Khatam*. 2015;**3**(2):19-26. doi: [10.18869/acadpub.shefa.3.2.19](https://doi.org/10.18869/acadpub.shefa.3.2.19).
24. Shamsaei F, Cheraghi F, Esmaeili R, Mohsenifard J. Explanation of loneliness in the elderly and comparison with psychosocial development theory: A quantitative study. *J Educ Community Health*. 2014;**1**(2):30-8. doi: [10.20286/jech-010230](https://doi.org/10.20286/jech-010230).
25. Nori pourlayavali R, Alikhani M, Hoseynian S, Soheylizadeh S, Mohammadiar M. The effect of group reminiscence on happiness and life expectancy of elderly journal of aging psychology. *J Aging Psychol*. 2016;**1**(3):147-15.
26. Garrow S, Walker JA. Existential group therapy and death anxiety. *Adultspan J*. 2001;**3**(2):77-87. doi: [10.1002/j.2161-0029.2001.tb00109.x](https://doi.org/10.1002/j.2161-0029.2001.tb00109.x).
27. Ghorbanalipour M, Falsafinejad MR, Sohrabi F, Borjali A. The effect of death anxiety and age on health promoting behaviors. *J Urmia Univ Med Sci*. 2010;**21**(2):286-92.
28. Soleimani MA, Lehto RH, Negarandeh R, Bahrami N, Nia HS. Relationships between death anxiety and quality of life in Iranian patients with cancer. *Asia Pac J Oncol Nurs*. 2016;**3**(2):183-91. doi: [10.4103/2347-5625.182935](https://doi.org/10.4103/2347-5625.182935). [PubMed: 27981157]. [PubMed Central: PMC5123493].
29. Siverova J, Buzgova R. The effect of reminiscence therapy on quality of life, attitudes to ageing, and depressive symptoms in institutionalized elderly adults with cognitive impairment: A quasi-experimental study. *Int J Ment Health Nurs*. 2018;**27**(5):1430-9. doi: [10.1111/inm.12442](https://doi.org/10.1111/inm.12442). [PubMed: 29427397].
30. Chueh KH, Chang TY. Effectiveness of group reminiscence therapy for depressive symptoms in male veterans: 6-month follow-up. *Int J Geriatr Psychiatry*. 2014;**29**(4):377-83. doi: [10.1002/gps.4013](https://doi.org/10.1002/gps.4013). [PubMed: 23946260].