

## Generation mean analysis of some physiological traits in the hybrid maize cv. SC704 under different water regimes

Mozhgan Shirinpour<sup>1</sup>, Ali Asghari<sup>1\*</sup>, Saeid Aharizad<sup>2</sup>, Ali Rasoulzadeh<sup>3</sup> and Saeid Khavari Khorasani<sup>4</sup>

Received: November 16, 2019 Accepted: June 7, 2020

<sup>1</sup>Department of Agronomy and Plant Breeding, University of Mohaghegh Ardabili, Ardabil, Iran.

<sup>2</sup>Department of Plant Breeding and Biotechnology, University of Tabriz, Tabriz, Iran.

<sup>3</sup>Department of Water Engineering, University of Mohaghegh Ardabili, Ardabil, Iran.

<sup>4</sup>Khorasan Razavi Agriculture and Natural Resources Research Center, Mashhad, Iran.

\*Corresponding author; Email: a\_asghari@uma.ac.ir

### Abstract

This research was carried out using seven generations of MO17 (the paternal line), B73 (the maternal line), F<sub>1</sub> (SC704), F<sub>2</sub>, F<sub>3</sub>, BC<sub>1</sub> and BC<sub>2</sub> to study the inheritance of chlorophyll fluorescence parameters and relative water content using generation mean analysis. The experiment was conducted in the research station of the University of Tabriz, Iran, with two replications for each generation under normal irrigation and two water deficit conditions. The existence of epistatic gene interaction was observed in controlling of F<sub>0</sub>, F<sub>m</sub>, F<sub>v</sub> (in the severe stress condition), F<sub>v</sub>/F<sub>m</sub> (in the normal condition), F<sub>v</sub>/F<sub>0</sub> (in the normal and severe stress conditions) and RWC (in the normal and moderate and severe stress conditions). Duplicate epistasis was observed for all studied traits under the normal and water deficit conditions except RWC in the severe stress condition. Therefore, exploiting non-additive gene action and delaying the selection to advanced generations for physiological traits in maize was suggested to adopt in the programs aimed at developing drought-tolerant lines and hybrids.

**Keywords:** Generation mean analysis; Heritability; Maize (*Zea mays* L.); Physiological traits; Water deficit stress

**Citation:** Shirinpour M, Asghari A, Aharizad S, Rasoulzadeh A and Khavari Khorasani S, 2020. Generation mean analysis of some physiological traits in hybrid maize cv. SC704 under different water regimes. Journal of Plant Physiology and Breeding 10(1): 141-148.

### Introduction

Various abiotic stresses such as drought, reduce the growth and yield of plants below the optimum level (Cramer *et al.* 2011). Maize (*Zea mays* L.) as one of the most important crops in the world, is drought sensitive, and its yield significantly decreases under water stress conditions (Shiferaw *et al.* 2011; Todaka *et al.* 2015). Drought causes degradation of photosynthetic pigments and thus decreases the light energy absorption on the photochemical apparatus and damages thylakoid membranes and centers of photosystem reaction (Batra *et al.* 2014). The chlorophylls and PSII absorb the light energy

which can be used to drive photosynthesis processes. When this energy is lost from PSII, it appears as heat or emitted chlorophyll fluorescence (Kalaji *et al.* 2012). Chlorophyll fluorescence has been extensively used to study the variation in the efficiency of light processing and stress tolerance in plants (Lichtenthaler *et al.* 2005; Stirbet *et al.* 2018). Relative leaf water content (RWC) has also been utilized for the evaluation of drought tolerance in crops (Dhanda and Sethi 2002). Decreasing this trait causes stomata closure and CO<sub>2</sub> assimilation declines under drought stress (Gindaba *et al.* 2004).

Studies about the inheritance of chlorophyll fluorescence components are very limited in maize. Hola *et al.* (2010) examined the response of several photosynthetic parameters of maize plants to drought stress in five inbred lines and their F1 hybrids. They reported that the nonadditive genetic effects were often more important than the additive effects for the photosynthetic characteristics. However, the genetic changed when the plants were exposed to drought conditions. Also, it was not possible to predict the response of F1 hybrids to drought based on the photosynthetic performance of their parents. Simic *et al.* (2014) conducted a quantitative trait loci (QTL) analysis on nine chlorophyll fluorescence parameters in maize at four environments that differed in weather conditions. The chlorophyll fluorescence parameters were analyzed among 205 recombinant inbred lines derived from the intermated B73 × Mo17 maize population. The results revealed 10 significant QTLs for seven chlorophyll fluorescence parameters, of which five were co-localized after combining over the four environments. These results indicated the role of polygenic inheritance in governing the chlorophyll fluorescence parameters under study. They also detected one pleiotropic locus on chromosome 7, coinciding with the gene *gst23* that possibly related to efficient photosynthesis in different field conditions.

Generation mean analysis is a biometrical method for estimating the genetic effects including additive, dominance and especially epistatic interactions by using means of different generations (Mather and Jinks 1982; Viana 2000).

Information about the nature and magnitude of gene actions involved in the tolerance to water deficit stress can be useful for breeding drought-resistant varieties in maize. Thus, a generation mean analysis was designed to study the inheritance of physiological characteristics in maize under different water regimes.

## Materials and Methods

### Plant material and experiment

Seven generations including B73 (the maternal line), MO17 (the paternal line), SC704 (F<sub>1</sub>), F<sub>2</sub>, BC<sub>1</sub>, BC<sub>2</sub> and F<sub>3</sub> were grown in PVC pipes with 20 cm diameter and the height of one meter, under normal irrigation and two water regimes (55% and 75% of available water depletion) in the research station of the University of Tabriz. With consideration of a proper distance between the seeds, two maize seeds were sown per pipe to secure at least one plant appearance in each pipe. Then, thinning was done at the tillering stage and only one plant was left in a pipe. Eventually, 20 plants for each generation were evaluated. Plants were grown in similar conditions in terms of soil type, light, temperature and pipes size. Also, the experimental site was protected from the rainfall by a plastic cover throughout the cropping period.

### Chlorophyll fluorescence parameters

Chlorophyll fluorescence was measured on a fully developed leaf using a hand-held chlorophyll fluorometer (Opti-Sciences, OS-30p). The leaves were adapted in dark for 15 min at first and then fluorescence was recorded at 650  $\mu\text{mol m}^{-2} \text{s}^{-1}$ . Fluorescence values recorded included: F<sub>0</sub>:

minimum fluorescence,  $F_m$ : maximum fluorescence,  $F_v$ : variable fluorescence =  $(F_m - F_0)$ ,  $F_v/F_m$ : maximum quantum yield of photosystem II (PS II) and  $F_v/F_0$ : maximum primary yield of the photochemistry of PS II.

### RWC

The measurement of RWC was accomplished by excising 1-cm disks from the fresh leaves. Five disks from each pipe were weighed immediately and the fresh weight (FW) was measured. Then, the leaf disks were placed in the distilled water for 24 h and the turgid weight (TW) was obtained. Finally, the samples were dried in an oven at 72 °C for 24 h and their dry weight (DW) was recorded. RWC was determined as follows (Salisbury and Ross 1998):

$$RWC = ((FW - DW) / (TW - DW)) \times 100$$

### Statistical and genetic analysis

Generation mean analysis was carried out using the weighted least squares method considering the inverse of the variance of the means within each generation as the weight (Mather and Jinks 1982). In this method, the overall average for each trait is shown as follows:  $Y = m + \alpha [d] + \beta [h] + \alpha^2 [i] + 2\alpha\beta [j] + \beta^2 [l]$  Where,  $Y$ = the generation mean,  $m$ =  $F_{\infty}$  metric,  $d$ = additive effects,  $h$ = dominance effects,  $i$ = additive  $\times$  additive interaction,  $j$ = additive  $\times$  dominance interaction,  $l$ = dominance  $\times$  dominance interaction and  $\alpha$ ,  $2\alpha\beta$  and  $\beta^2$  are coefficients of genetic parameters. To verify the goodness of fit of the model, a joint scaling test was carried out for each trait at each irrigation level (Cavalli 1952). All

genetic parameters were tested for significance using a t-test.

### Results and Discussion

The results showed that the scaling tests for the three-parameter model were not significant for  $F_v$  under the normal and mild stress conditions (55% AWD),  $F_v/F_m$  under both stress conditions and  $F_v/F_0$  under the mild stress condition, which indicates the lack of epistasis for these traits (Table 1). The scaling tests were significant for the other studied traits suggesting the inadequacy of the additive-dominance model or the presence of epistasis type of gene effects.

Estimates of genetic effects based on the three- or six-parameter models for the studied traits under different water regimes are given in Table 2. The generation mean analysis suggested the importance of both main effects (dominance and additive effects) and non-allelic interaction for most of the studied traits. However, for almost all characters the dominance effects were higher in magnitude as compared to the additive gene effects under different water regimes. The opposite signs of  $[h]$  and  $[l]$  revealed duplicate epistasis for all traits in the normal and water deficit conditions except RWC in the severe stress condition. Negative and positive signs of the  $[h]$  parameter showed that partial dominance was towards the decreasing  $F_0$ ,  $F_m$ ,  $F_v$  and RWC and increasing of  $F_v/F_m$  and  $F_v/F_0$  in the cross under consideration, respectively.

**F<sub>0</sub>**: The best-fit model for this trait was the six-parameter model  $[m-d-h-i-j-l]$ . Additive  $\times$  additive

Table 1. Scaling tests ( $\pm$ SE) for physiological traits in maize generations at different water regimes.

Traits	Water Regimes	A	B	C	D
$F_0$ ( $\mu\text{mol. m}^2\text{s}^{-1}$ )	Full irrigation	$-41.85 \pm 5.84^{**}$	$-18.1 \pm 7.66^{**}$	$-10.55 \pm 14.80^{\text{ns}}$	$8.55 \pm 14.23^{\text{ns}}$
	55% AWD	$-32.45 \pm 10.20^{**}$	$-11.55 \pm 9.81^{\text{ns}}$	$-30.20 \pm 21.23^{\text{ns}}$	$40.50 \pm 16.76^{**}$
	75% AWD	$-12.50 \pm 6.06^*$	$-17.25 \pm 7.09^{**}$	$12.95 \pm 14.76^{\text{ns}}$	$28.65 \pm 14.32^{**}$
$F_m$ ( $\mu\text{mol. m}^2\text{s}^{-1}$ )	Full irrigation	$-35.35 \pm 30.81^{\text{ns}}$	$-25.35 \pm 32.91^{\text{ns}}$	$40.60 \pm 55.44^{\text{ns}}$	$109.80 \pm 49.79^{**}$
	55% AWD	$-61.30 \pm 25.35^{**}$	$-64.85 \pm 25.51^{**}$	$13.05 \pm 44.61^{\text{ns}}$	$77.05 \pm 42.42^{\text{ns}}$
	75% AWD	$-50.70 \pm 29.04^{\text{ns}}$	$-72.60 \pm 34.95^{**}$	$-144.30 \pm 56.86^{**}$	$-8.90 \pm 64.72^{\text{ns}}$
$F_v$ ( $\mu\text{mol. m}^2\text{s}^{-1}$ )	Full irrigation	$5.40 \pm 31.60^{\text{ns}}$	$-7.30 \pm 29.20^{\text{ns}}$	$51.80 \pm 60.19^{\text{ns}}$	$95 \pm 54.48^{\text{ns}}$
	55% AWD	$-28.85 \pm 28.63^{\text{ns}}$	$-53.30 \pm 27.57^{\text{ns}}$	$43.25 \pm 45.70^{\text{ns}}$	$36.55 \pm 42.47^{\text{ns}}$
	75% AWD	$-40.10 \pm 31.75^{\text{ns}}$	$-66.35 \pm 35.99^{\text{ns}}$	$-119.65 \pm 54.81^{**}$	$-29.25 \pm 69.69^{\text{ns}}$
$F_v/F_m$	Full irrigation	$0.05 \pm 0.02^{**}$	$0.02 \pm 0.02^{\text{ns}}$	$0.03 \pm 0.04^{\text{ns}}$	$0.02 \pm 0.04^{\text{ns}}$
	55% AWD	$0.02 \pm 0.02^{\text{ns}}$	$-0.01 \pm 0.02^{\text{ns}}$	$0.05 \pm 0.03^{\text{ns}}$	$-0.03 \pm 0.02^{\text{ns}}$
	75% AWD	$-0.007 \pm 0.02^{\text{ns}}$	$-0.02 \pm 0.02^{\text{ns}}$	$-0.02 \pm 0.03^{\text{ns}}$	$-0.04 \pm 0.04^{\text{ns}}$
$F_v/F_0$	Full irrigation	$0.71 \pm 0.21^{**}$	$0.21 \pm 0.22^{\text{ns}}$	$0.46 \pm 0.42^{\text{ns}}$	$0.40 \pm 0.35^{\text{ns}}$
	55% AWD	$0.35 \pm 0.27^{\text{ns}}$	$-0.12 \pm 0.23^{\text{ns}}$	$0.70 \pm 0.46^{\text{ns}}$	$-0.42 \pm 0.35^{\text{ns}}$
	75% AWD	$-0.05 \pm 0.18^{\text{ns}}$	$-0.07 \pm 0.19^{\text{ns}}$	$-0.86 \pm 0.36^{**}$	$-0.47 \pm 0.39^{\text{ns}}$
RWC (%)	Full irrigation	$-24.22 \pm 3.65^{**}$	$-9.30 \pm 4.26^{**}$	$-38.42 \pm 7.59^{**}$	$-6.98 \pm 7.74^{\text{ns}}$
	55% AWD	$-20.11 \pm 2.72^{**}$	$-15.88 \pm 2.64^{**}$	$-43.66 \pm 4.73^{**}$	$-2.07 \pm 5.62^{**}$
	75% AWD	$-10.32 \pm 2.50^{**}$	$-3.75 \pm 3.30^{\text{ns}}$	$-26.83 \pm 5.39^{**}$	$-15.84 \pm 6.04^{**}$

ns, \* and \*\*: non-significant and significant at 0.05 and 0.01 probability levels, respectively; AWD: available water depletion;  $F_0$ ,  $F_m$ ,  $F_v$ ,  $F_v/F_m$  and  $F_v/F_0$ : minimum fluorescence, maximum fluorescence value, variable fluorescence, the maximum quantum yield of photosystem II (PSII) and maximum primary yield of the photochemistry of PSII, respectively; RWC: relative water content.

and dominance  $\times$  dominance interactions had a significant role in the genetic control of  $F_0$  in all three water regimes. Asadi *et al.* (2015) reported a high additive gene effect for  $F_0$  in wheat under normal conditions. Contrary, the non-additive gene effect was reported in sugarcane for  $F_0$  by Zhang *et al.* (2010).

**$F_m$ :** In the normal and mild stress conditions the six-parameter model [m-d-h-i-j-l] and in the severe stress conditions the five-parameter model [m-d-h-j-l] were the best-fit models for  $F_m$ . Dominance and dominance  $\times$  dominance effects were significant and governed the genetics of this character. Therefore, exploiting the non-additive gene action can be effective for improving  $F_m$ . The same findings were documented by Hola *et al.* (2010) in five inbred lines of maize. The authors revealed that epistatic and dominance gene effects were

important in controlling the chlorophyll fluorescence parameters.

**$F_v$ :** The findings revealed that the three-parameter model [m-d-h] was fitted well for  $F_v$  under normal and mild stress conditions. Both additive and dominance gene effects were significant for this trait under the two aforementioned conditions. Whereas, in the severe stress condition the five-parameter model [m-d-h-j-l] was determined as the best model for this character. The dominance and dominance  $\times$  dominance effects were also significant and had an important role in controlling  $F_v$ .

**$F_v/F_m$ :** The five-parameter model [m-d-h-j-l] was best fitted for this trait in the normal conditions. The dominance gene effect and dominance  $\times$  dominance interaction were significant and

Table 2. Estimates of genetic parameters ( $\pm$ SE),  $\chi^2$  for the joint scaling test for physiological traits in maize at different water regimes using generation mean analysis.

Traits	Water Regimes	m	d	h	i	j	l	$\chi^2$	df
$F_0$ ( $\mu\text{mol. m}^2\text{s}^{-1}$ )	Full irrigation	208.6 $\pm$ 5.85**	-2.54 $\pm$ 2.18 <sup>ns</sup>	-101.7 $\pm$ 17.27 <sup>ns</sup>	-23.83 $\pm$ 5.94**	-25.67 $\pm$ 8.54 <sup>ns</sup>	78.79 $\pm$ 13.38**	3.59 <sup>ns</sup>	1
	55% AWD	224.2 $\pm$ 7.08**	4.44 $\pm$ 2.51 <sup>ns</sup>	-99.4 $\pm$ 23.62**	-25.82 $\pm$ 7.22**	-21.05 $\pm$ 12.91 <sup>ns</sup>	72.35 $\pm$ 18.99**	0.32 <sup>ns</sup>	1
	75% AWD	244.8 $\pm$ 5.82**	8.39 $\pm$ 2.20**	-85.8 $\pm$ 16.89**	-27.12 $\pm$ 5.91**	4.06 $\pm$ 8.30 <sup>ns</sup>	54.04 $\pm$ 12.99**	1.36 <sup>ns</sup>	1
$F_m$ ( $\mu\text{mol. m}^2\text{s}^{-1}$ )	Full irrigation	738.1 $\pm$ 22.31**	13.87 $\pm$ 8.60 <sup>ns</sup>	-212.1 $\pm$ 74.47**	-78.19 $\pm$ 22.86**	-10.57 $\pm$ 42.48 <sup>ns</sup>	131.6 $\pm$ 58.06**	0.16 <sup>ns</sup>	1
	55% AWD	788.9 $\pm$ 18.94**	22.53 $\pm$ 6.85**	-272.3 $\pm$ 61.39**	-80.51 $\pm$ 19.33**	3.22 $\pm$ 34.12 <sup>ns</sup>	188.3 $\pm$ 46.91**	1.60 <sup>ns</sup>	1
	75% AWD	727.0 $\pm$ 9.54**	-2.65 $\pm$ 9.81 <sup>ns</sup>	-120.1 $\pm$ 43.11**	-	23.22 $\pm$ 42.11 <sup>ns</sup>	129.0 $\pm$ 41.29**	0.19 <sup>ns</sup>	2
$F_v$ ( $\mu\text{mol. m}^2\text{s}^{-1}$ )	Full irrigation	484.2 $\pm$ 7.38**	17.57 $\pm$ 8.21**	-9.10 $\pm$ 13.38 <sup>ns</sup>	-	-	-	6.35 <sup>ns</sup>	4
	55% AWD	513.5 $\pm$ 5.85**	20.22 $\pm$ 6.54**	-9.62 $\pm$ 11.55**	-	-	-	9.99 <sup>ns</sup>	4
	75% AWD	513.2 $\pm$ 10.20**	-9.52 $\pm$ 10.48 <sup>ns</sup>	-101.1 $\pm$ 44.79**	-	27.34 $\pm$ 45.08 <sup>ns</sup>	112.7 $\pm$ 42.36**	0.05 <sup>ns</sup>	2
$F_v/F_m$	Full irrigation	0.72 $\pm$ 0.006**	0.01 $\pm$ 0.007 <sup>ns</sup>	0.06 $\pm$ 0.03*	-	0.03 $\pm$ 0.02 <sup>ns</sup>	-0.06 $\pm$ 0.03**	0.50 <sup>ns</sup>	2
	55% AWD	0.72 $\pm$ 0.003**	0.004 $\pm$ 0.004 <sup>ns</sup>	0.003 $\pm$ 0.008 <sup>ns</sup>	-	-	-	4.40 <sup>ns</sup>	4
	75% AWD	0.70 $\pm$ 0.005**	-0.009 $\pm$ 0.005 <sup>ns</sup>	0.008 $\pm$ 0.008 <sup>ns</sup>	-	-	-	2.36 <sup>ns</sup>	4
$F_v/F_0$	Full irrigation	2.58 $\pm$ 0.06**	0.12 $\pm$ 0.06*	0.76 $\pm$ 0.28**	-	0.50 $\pm$ 0.29 <sup>ns</sup>	-0.78 $\pm$ 0.27**	0.99 <sup>ns</sup>	2
	55% AWD	2.60 $\pm$ 0.05**	0.05 $\pm$ 0.05 <sup>ns</sup>	0.05 $\pm$ 0.10 <sup>ns</sup>	-	-	-	4.74 <sup>ns</sup>	4
	75% AWD	2.03 $\pm$ 0.10**	-0.14 $\pm$ 0.06**	0.44 $\pm$ 0.13**	0.33 $\pm$ 0.11**	0.03 $\pm$ 0.02 <sup>ns</sup>	-	1.24 <sup>ns</sup>	2
RWC (%)	Full irrigation	71.3 $\pm$ 3.34**	1.10 $\pm$ 1.15 <sup>ns</sup>	-27.02 $\pm$ 10.20**	0.73 $\pm$ 3.39 <sup>ns</sup>	-14.72 $\pm$ 5.35**	33.92 $\pm$ 7.52**	0.29 <sup>ns</sup>	1
	55% AWD	64.9 $\pm$ 2.44**	-3.57 $\pm$ 0.75**	-37.09 $\pm$ 7.19**	-1.93 $\pm$ 2.46 <sup>ns</sup>	-4.40 $\pm$ 3.52 <sup>ns</sup>	40.88 $\pm$ 5.33**	4.58 <sup>ns</sup>	1
	75% AWD	41.2 $\pm$ 2.61**	-4.27 $\pm$ 0.85**	2.88 $\pm$ 7.75 <sup>ns</sup>	8.73 $\pm$ 2.64**	-6.15 $\pm$ 3.86 <sup>ns</sup>	6.49 $\pm$ 5.70 <sup>ns</sup>	0.59 <sup>ns</sup>	1

See Table 1 for abbreviations.

involved in governing the genetics of  $F_v/F_m$ . Under both water stress conditions, the three-parameter model [m-d-h] was fitted for this character. Similar findings were reported by Vijayalakshmi *et al.* (2010) in winter wheat which showed a low additive gene effect for  $F_v/F_m$ . The quantum efficiency of PS II ( $F_v/F_m$ ) has a major role in photosynthesis and it can be used as a screening tool to identify tolerant varieties to drought (Sharma *et al.* 2014). A low ratio of  $F_v/F_m$  shows low photosynthetic efficiency, thus, genotypes that have a higher ratio of  $F_v/F_m$  may be more tolerant to drought stress (Ristic *et al.* 2007; Kumar *et al.* 2012).

**$F_v/F_0$ :** In the normal and severe stress conditions the five-parameter models [m-d-h-j-l] and [m-d-h-i-j] were regarded as the best models explaining the

inheritance of this character, respectively. Thus, both main gene effects and epistatic interactions governed the control of this trait under these conditions. However, under the mild stress conditions, only the three-parameter model was best fitted for this trait.

**RWC:** Under normal and water deficit conditions the six-parameter model [m-d-h-i-j-l] was the best fitted this character. Both main effects were significant but the dominance effect was higher than the additive effect in the normal and mild stress conditions. Generation mean analysis in maize (Moharramnejad *et al.* 2018), as well as durum wheat (Salmi *et al.* 2019), under control and drought stress conditions showed the importance of dominance genes effects in operating the inheritance of RWC. In contrast to our results, the

additive gene effect for RWC in wheat was reported by Said (2014) under the normal condition and Asadi *et al.* (2015) under both normal and water deficit conditions. It seems that the genetic effects vary according to crop species, varieties and the environmental conditions in which the research was conducted.

### Conclusion

In this experiment, the importance of dominance and epistasis (duplicate) effect was recognized for most of the studied physiological traits under water stress conditions, which implies exploiting non-

additive gene action and delaying the selection to advanced generations for these traits in maize aiming at developing drought-tolerant lines and hybrids in maize. However, the additive effect was more important in the inheritance of  $F_0$  and  $F_V/F_m$  in the normal condition and early selection can be efficient for the improvement of these characters.

### Conflict of Interest

The authors declare that they have no conflict of interest with any organization concerning the subject of the manuscript.

### References

- Asadi AA, Valizadeh M, Mohammadi SA and Khodarahmi M, 2015. Genetic analysis of some physiological traits in wheat by generations mean analysis under normal and water deficit conditions. *Biological Forum* 7(2): 722-733.
- Batra NG, Sharma V and Kumari N, 2014. Drought-induced changes in chlorophyll fluorescence, photosynthetic pigments, and thylakoid membrane proteins of *Vigna radiata*. *Journal of Plant Interactions* 9(1): 712-721.
- Cavalli L, 1952. An analysis of linkage in quantitative inheritance. In: Reive ECR and Waddington CH (eds). *Quantitative Inheritance*. Pp. 135-144. HMSO, London.
- Cramer GR, Urano K, Delrot S, Pezzotti M and Shinozaki K, 2011. Effects of abiotic stress on plants: a systems biology perspective. *BMC Plant Biology* 11: 1-14.
- Dhanda S and Sethi G, 2002. Tolerance to drought stress among selected Indian wheat cultivars. *The Journal of Agricultural Science* 139(3): 319-326.
- Gindaba J, Rozanov A and Negash L, 2004. Response of seedlings of two Eucalyptus and three deciduous tree species from Ethiopia to severe water stress. *Forest Ecology* 201(1): 119-129.
- Hao D, Chao M, Yin Z and Yu D, 2012. Genome-wide association analysis detecting significant single nucleotide polymorphisms for chlorophyll and chlorophyll fluorescence parameters in soybean (*Glycine max*) landraces. *Euphytica* 186(3): 919-931.
- Hola D, Benesova M, Honnerova J, Hnilicka F, Rothova O, Kocova M and Hnilickova H, 2010. The evaluation of photosynthetic parameters in maize inbred lines subjected to water deficiency: can these parameters be used for the prediction of performance of hybrid progeny? *Photosynthetica* 48(4): 545-558.
- Kalaji HM, Goltsev V, Bosa K, Allakhverdiev SI and Strasser RJ, 2012. Experimental in vivo measurements of light emission in plants: a perspective dedicated to David Walker. *Photosynthesis Research* 114(2): 69-96.
- Kumar S, Sehgal SK, Kumar U, Prasad PV, Joshi AK and Gill BS, 2012. Genomic characterization of drought tolerance-related traits in spring wheat. *Euphytica* 186(1): 265-276.
- Lichtenthaler H, Buschmann C and Knapp M, 2005. How to correctly determine the different chlorophyll fluorescence parameters and the chlorophyll fluorescence decrease ratio  $R_{Fd}$  of leaves with the PAM fluorometer. *Photosynthetica* 43(3): 379-393.

- Mather K and Jinks JL, 1982. Biometrical Genetics: the Study of Continuous Variation. Third edition. Chapman and Hall, USA, 396 pp.
- Moharramnejad S, Valizadeh M and Emaratpardaz J, 2018. Generation mean analysis in maize (*Zea mays* L.) under drought stress. Fresenius Environmental Bulletin 27(4): 2518-2522.
- Ristic Z, Bukovnik U and Prasad PV, 2007. Correlation between heat stability of thylakoid membranes and loss of chlorophyll in winter wheat under heat stress. Crop Science 47(5): 2067-2073.
- Said AA, 2014. Generation mean analysis in wheat (*Triticum aestivum* L.) under drought stress conditions. Annals of Agricultural Sciences 59(2): 177-184.
- Salisbury F and Ross C, 1998. Plant Physiology. Fourth edition. Wads Worth Publishing Company, Belmont, California, USA.
- Salmi M, Benmahammed A, Benderradji L, Fellahi ZEA, Bouzerzour H, Oulmi A and Benbelkacem A, 2019. Generation means analysis of physiological and agronomical targeted traits in durum wheat (*Triticum durum* Desf.) cross. Revista Facultad Nacional de Agronomía Medellín 72(3): 8971-8981.
- Sharma DK, Fernandez JO, Rosenqvist E, Ottosen CO and Andersen SB, 2014. Genotypic response of detached leaves versus intact plants for chlorophyll fluorescence parameters under high temperature stress in wheat. Journal of Plant Physiology 171(8): 576-586.
- Shiferaw B, Prasanna BM, Hellin J and Banziger M, 2011. Crops that feed the world. 6. Past successes and future challenges to the role played by maize in global food security. Food Security 3(3): 307-327.
- Simic D, Lepedus H, Jurkovic V, Antunovic J and Cesar V, 2014. Quantitative genetic analysis of chlorophyll a fluorescence parameters in maize in the field environments. Journal of Integrative Plant Biology 56(7): 695-708.
- Stirbet A, Lazar D, Kromdijk J and Govindjee G, 2018. Chlorophyll a fluorescence induction: can just a one-second measurement be used to quantify abiotic stress responses? Photosynthetica 56: 86-104.
- Todaka D, Shinozaki K and Yamaguchi-Shinozaki K, 2015. Recent advances in the dissection of drought-stress regulatory networks and strategies for development of drought-tolerant transgenic rice plants. Frontiers in Plant Science 6: 84. doi: 10.3389/fpls.2015.00084.
- Viana JMS, 2000. Generation mean analysis in relation to polygenic systems with epistasis and fixed genes. Pesquisa Agropecuaria Brasileira 35(6): 1159-1167.
- Vijayalakshmi K, Fritz AK, Paulsen GM, Bai G, Pandravada S and Gill BS, 2010. Modeling and mapping QTL for senescence-related traits in winter wheat under high temperature. Molecular Breeding 26(2): 163-175.
- Zhang ZB, Xu P, Jia JZ and Zhou RH, 2010. Quantitative trait loci for leaf chlorophyll fluorescence traits in wheat. Australian Journal of Crop Science 4(8): 571-579.

## تجزیه میانگین نسل‌ها برای برخی صفات فیزیولوژیک در هیبرید سینگل کراس ۷۰۴ ذرت تحت رژیم‌های آبی مختلف

مژگان شیرین‌پور<sup>۱</sup>، علی اصغری<sup>۱\*</sup>، سعید اهری‌زاد<sup>۲</sup>، علی رسول‌زاده<sup>۳</sup> و سعید خاوری خراسانی<sup>۴</sup>

- ۱- گروه زراعت و اصلاح نباتات، دانشگاه محقق اردبیلی، اردبیل.
  - ۲- گروه به‌نژادی و بیوتکنولوژی گیاهی، دانشگاه تبریز، تبریز.
  - ۳- گروه مهندسی آب، دانشگاه محقق اردبیلی، اردبیل.
  - ۴- مرکز تحقیقات کشاورزی و منابع طبیعی خراسان رضوی، مشهد.
- \*مسئول مکاتبه؛ Email: a\_asghari@uma.ac.ir

### چکیده

تحقیق حاضر با استفاده از هفت نسل {MO17 به عنوان لاین پدری؛ B73 به عنوان لاین مادری؛ هیبرید (F1) SC704؛ F2؛ F3؛ BC1؛ BC2} به منظور بررسی وراثت پارامترهای فلورسانس کلروفیل و محتوای آب نسبی توسط تجزیه میانگین نسل‌ها انجام شد. این آزمایش در ایستگاه تحقیقاتی دانشگاه تبریز با استفاده از طرح بلوک‌های کامل تصادفی با دو تکرار در شرایط آبیاری نرمال و دو شرایط کمبود آب انجام شد. وجود اثرات متقابل اپیستازی در وراثت Fm، F0، FV (در شرایط تنش شدید)، FV/Fm (در شرایط عادی)، FV/F0 (در شرایط نرمال و تنش شدید) و RWC (در شرایط نرمال و تنش‌های ملایم و شدید) مشاهده شد. اپیستازی مضاعف در کلیه صفات مورد مطالعه به جز RWC در شرایط تنش شدید، مشاهده شد. این نتایج حاکی از لزوم بهره برداری از اثرات غیرافزایشی ژنتیکی و به تأخیر انداختن گزینش صفات فیزیولوژیکی مورد مطالعه به نسل‌های پیشرفته در برنامه‌های اصلاح ذرت برای تحمل خشکی می‌باشند.

**واژه‌های کلیدی:** تجزیه میانگین نسل‌ها؛ تنش کم‌آبی؛ ذرت؛ صفات فیزیولوژیکی؛ وراثت پذیری.