

## Nutritional Assessment in the Critically Ill Patients

Elham Bagheri<sup>1</sup>, Farzaneh Hematian<sup>2\*</sup>, Mandana Izadpanah<sup>2</sup>, Mahbobeh Rashidi<sup>3</sup>

<sup>1</sup>Student Research Committee, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.

<sup>2</sup>Department of Clinical Pharmacy, School of Pharmacy, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.

<sup>3</sup>Department of Anesthesiology, Faculty of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.

### ARTICLE INFO

#### Article history:

Received 05 May 2020

Revised 26 May 2020

Accepted 10 June 2020

#### Keywords:

Intensive care unit;

Malnutrition;

Nutritional support;

Modified-nutrition risk in critically ill (m-nutric) score

### ABSTRACT

**Background:** Malnutrition is a prevalent complication among critically ill patients. It has very detrimental effects on the patients' clinical course. This study aimed to investigate the impact of nutrition in the intensive care unit (ICU) patients.

**Methods:** In this epidemiologic-analytic study conducted in the surgical ICU of Imam Khomeini hospital, Ahvaz, Iran, 34 patients were selected and divided into two groups. The first group of patients received the appropriate nutrition. The second group received an inappropriate diet, and the nutritional risk was evaluated according to the modified- Nutrition Risk in Critically ill (m-NUTRIC) score. The energy was calculated by using 25 Kcal/kg, also the two groups were compared in terms of ICU mortality, ICU stays, Acute Physiologic Assessment and Chronic Health Evaluation (APACHE) II Scoring, and the Sequential Organ Failure Assessment (SOFA) Score.

**Results:** Baseline data, such as APACHE II score and mean age, except sex, were not significantly different between the two groups. In this study, results were toward shorter ICU stay, less mortality, and better SOFA score in the group receiving appropriate nutrition compared to the other group. However, due to the low number of patients, no significant differences were observed in the two groups.

**Conclusion:** Our data suggest that nutritional support should be considered as an essential part of the medication during critical illness.

© 2020 Tehran University of Medical Sciences. All rights reserved.

It is well known the role of nutritional support in critical illness is essential [1-3]. Malnutrition is prevalent complication among critically ill patients [4]. The large numbers of patients hospitalized in the intensive care unit (ICU) suffer from catabolic stress, which refers to be the body inflammatory response to stress or injury. Malnutrition significantly increases mortality and length of stay in critically ill patients [5-10].

The first step in the prevention and treatment of malnutrition is the assessment of nutritional status and correction of diet intake [9,11].

Enteral nutrition (EN) in these patients is superior to parenteral nutrition (PN) [12-13]. There is increasing recognition that aggressive early feeding, as well as prolonged underfeeding both should be avoided [1,3],

although specific amounts of EN required for improved outcomes continue to be questioned [14].

The guidelines suggest that a validated score, such as the Nutrition Risk in the critically ill (NUTRIC) score, be utilized to determine nutrition risk in ICU patients [15]. It was developed to link starvation, inflammation, and clinical outcomes [16]. The incorporation of IL-6 limits the original NUTRIC score, so IL-6 in the modified-NUTRIC (m-NUTRIC) score has been removed. It was shown to contribute very little to the overall prediction of the NUTRIC score [17].

One study reported that the prevalence of malnutrition increased significantly in ICU patients during hospitalization [18]. The amount of malnutrition prevalence and mortality of ICU patients in Iran, due to a few studies, is not clear. The few reviews have been reported in the range of 25-48% [19-20].

The authors declare no conflicts of interest.

\*Corresponding author.

E-mail address: Farzanehematian@yahoo.com

© 2020 Tehran University of Medical Sciences. All rights reserved.

The main aim of this study was to make a nutritional assessment of ICU patients referring to Imam Khomeini hospital in Ahvaz, Iran. Furthermore, we documented the length of stay in ICU, the mortality rate in the ICU, and the Sequential Organ Failure Assessment (SOFA) score.

### Methods

This study was performed in the surgical ICU of Imam Khomeini teaching hospital in Ahvaz, Iran. This was an epidemiologic-analytic study for Jul 23, 2019, to Oct 22, 2019, which included 50 ICU patients over the age of 18 with an ICU stay for a minimum duration of 3 days and m-NUTRIC score  $\geq 5$ . Patients were excluded if patients were <18 years, ICU stays less than three days, m-NUTRIC score <5, pregnancy, and lactation (Figure 1).


The ethical committee of the university approved the study (file number: IR.AJUMS.REC.1397.857)

Baseline data recorded were age, sex, Acute Physiologic Assessment and Chronic Health Evaluation (APACHE) II Scoring [21], and actual body weight. Also, diagnosis, type of feeding, SOFA score [22], ICU length of stay, and mortality rate in the ICU were documented. Nutritional support status for adult patients during the study period was enrolled for appropriateness evaluation according to the m-NUTRIC score [23].

We collected data on nutrition intake in the duration of ICU admission daily, including volume and type of EN and PN products administered. We divided patients with m-NUTRIC score  $\geq 5$ , into two groups. The first group of patients received the appropriate nutrition. The second group received an inappropriate diet. All the findings of the study were done in these two groups. According to the American Society for Parenteral and Enteral Nutrition (ASPEN) guideline, the energy target was defined as 25Kcal/Kg/d. Patients' requirements were compared to intake food. Findings were assessed and confirmed by a clinical pharmacist.

Baseline characteristics are presented as mean  $\pm$  standard deviation or median (minimum-maximum) for quantitative variables, and as the number (frequency percentage) for qualitative variables. Quantitative variables were compared using an independent sample t-test, and qualitative variables were compared using a chi-square test. A P-value  $\leq 0.05$  was considered significant in outcome analysis. Data analysis was performed using SPSS version 22.

**Figure 1- Frequency of patients included in (and excluded from) the study according to screening results**



### Results

Among the patients admitted to the ICU during in a 3-month study period, 50 patients were included. According to the m-NUTRIC score, 34 patients (68%) were classified as m-NUTRIC score  $\geq 5$ , and 16 (32%) were classified as m-NUTRIC score <5. In this study, only the patients, who had a mNUTRIC score  $\geq 5$ , were evaluated. Out of the 34 patients with a m-NUTRIC score  $\geq 5$ , 8 (23.5%) had appropriate nutrition (group I) and 26 (76.5%) had inappropriate nutrition (group II) (Figure1/Table2). There were 22 (64.7%) men and 12 (35.3%) women (Table1).

The baseline characteristics of the patients are summarized in table1. Fundamental patient characteristics recorded were age, sex, weight, and APACHE II scores. There was a significant difference between the two groups in sex ( $p=0.00$ ). There were no significant differences between the two groups in the average of the weight ( $p=0.26$ ), age ( $p=0.85$ ), and APACHE II scores ( $p=0.14$ ) (Table1).

The majority of the patients received enteral feeding, and 29.4% of them did not receive any form of feeding for the entire period of stay in the ICU (Table1).

Table2 shows nutritional support status according to the m-NUTRIC score in two groups. There was a significant positive difference between two groups ( $P \leq 0.05$ ) (Table2).

Table3 shows the secondary clinical outcomes in the patients over the study period. There were no significant differences between the two groups in ICU length of stay ( $P=0.96$ ) (Figure 2), SOFA score ( $P= 0.48$ ) (Figure 3) and ICU mortality ( $P=0.6$ ) (Table 3).

**Table 1- Characteristics of the patients and feeding parameters**

Variable N= 34	Mean ±SD, median (IQR) or N (%)			P-value*
	Total	Group I	Group II	
Age (y)	62.82±17.92	61.75±15.98	63.15±18.76	0.85
Sex	65.50 (30,87)	62 (33,85)	66 (30,87)	0.00
	Male 22 (64.7%)	1 (2.9%)	21 (61.8%)	
	Female 12 (35.3%)	7 (20.6%)	5 (14.7%)	
Weight	68.71±11.22	64.75±12.78	69.92±10.67	0.26
APACHE II	70 (42.90)	69 (43.80)	70 (42.90)	0.14
	26.97±7.11	23.75±6.56	27.96±70.09	
	27.50 (13.39)	23 (16.36)	29 (13.39)	
Diagnosis				
Postoperative care	31 (62%)			
Cardiovascular and respiratory arrest	6 (12%)			
Multiple trauma	3 (6%)			
Sepsis	3 (6%)			
Pemphigus Vulgaris	2 (4%)			
Others	5 (10%)			
Type of feeding				
EN: 16(47.1%)				
PN: 3 (8.8%)				
EN, PN: 5 (14.7%)				
Nothing: 10 (29.4%)				

Note: Group I: Patients with appropriate nutrition; Group II: patients with inappropriate nutrition

Abbreviation: N: number of patients; APACHE II score: Acute physiology and chronic health evaluation II score; IQR: Interquartile range (1st- 3rd quartile); EN: Enteral nutrition; PN: Parenteral Nutrition; SD: Standard deviation; \*calculated by chi-square test for qualitative variable and independent sample t-test for the quantitative variable.

**Table 2- Nutritional support status according to the m-NUTRIC score in two groups**

Total	Nutritional support status		P-value*
	Group I	Group II	
34 (100%)	8 (23.5%)	26 (76.5%)	0.00

Note: Group I: Patients with appropriate nutrition; Group II: patients with inappropriate nutrition

Abbreviation: N: number of patients; m-NUTRIC score: modified- Nutrition Risk in the critically ill score.

\* calculated by chi-square test for qualitative variable and independent sample t-test for the quantitative variable.

**Table3- The secondary clinical outcomes in the patients over the study period**


	Clinical outcomes in two groups			P-value*
	Total	Group I	Group II	
Length of ICU. stay	15.50±19/88	15.25±13/38	15.58±21.72	0.96
N(d)	8 (3,89)	13.50 (3,42)	6.50 (3,89)	
ICU mortality	23 (67.6%)	5 (14.7%)	18 (52.9%)	0.6
SOFA score	9.12±3.09	8.43±2.00	9.33±3.36	0.48
	n9.66 (0.33,16)	8.57 (5.67,11.33)	10.16(0.33,16)	

Note: Group I: Patients with appropriate nutrition; Group II: patients with inappropriate nutrition


Abbreviation: ICU: Intensive care unit, SOFA score: sequential organ failure assessment, IQR: Interquartile range (1st- 3rd quartile), SD: standard deviation

\* calculated by chi-square test for qualitative variable and independent sample t-test for the quantitative variable.

**Figure 2- The ICU length of stay in two groups**



**Figure 3- The mean number of SOFA score in two groups**



Note: SOFA= Sequential Organ Failure Assessment

**Discussion**

This epidemiologic-analytic study was conducted on 50 ICU inpatients for three months. Patients who met the inclusion criteria were categorized into two groups. We compared the SOFA score [22], ICU length of stay, and mortality rate in two groups.

In this study, there was a significant difference in appropriate nutrition between the two groups. The appropriate and inappropriate nutrition in two groups was 23.5% and 76.5%, respectively. It may be due to in ICU cure; the focus is more on the treatment of the disease than start proper nutrition. Unawareness that it is an important part of the patient's treatment. Nachvak et al. showed a significant increment in the prevalence of malnutrition among critically ill patients in the ICU discharge day (83.5%) compared to the ICU admission

day (42.6%) [18]. At the same time, Hejazi et al. reported a rate of 58.62% for malnutrition on the ICU discharge day compared to 28.8% on the admission day [24]. Based on the studies, malnutrition prevalence has been 13 to 78 percent between the years of 1996 to 2005 [25]. The different prevalence rates of malnutrition can be explained by the use of different nutritional support, disease severity, and performance of medical teams in study populations [25]. Malnutrition is associated with morbidity and mortality during ICU stay [26-27]. Leiva Badosa et al. showed that malnutrition was associated with longer ICU stay and higher ICU mortality [28].

Also, providing a type of nutrition (EN or PN) is another important issue that should be considered. EN nutrition in critically ill patients is superior to parenteral nutrition [12-13]. The benefits of EN in them cause maintenance of the functional and structural integrity of the gut, decreased infectious complications, shorter hospital stays, and lower mortality [15]. Also, high-risk patients should receive nutritional support, initiated within the first 24 to 48 hours after admission [29].

In this study, results were toward shorter hospital stay, less mortality, and better SOFA score in the group receiving appropriate nutrition compared to the other group. However, due to the low number of patients, no significant differences were observed in the two groups.

Similar and opposite results were observed in other studies. For example, the results of Rice et al.'s study showed that there were no significant differences in ventilator-free days, length of stay, and mortality rates among ICU patients [30]. However, Villet et al.'s study found a strong correlation between inadequate feeding and increased incidence of ICU mortality [31].

Raman et al. evaluated patient nutrition by m-NUTRIC score and showed adequate nutrition could decrease mortality in high-risk patients [23]. One possible explanation for this discrepancy is the small sample size in our study. Another limitation of the study was the lack of beds to measure body weight. Of course, to reduce errors, all estimated weights were conducted by one person.

**Recommendation**

We recommend further studies using a larger sample.

**Conclusion**

Our data suggest that nutritional support should be considered as an essential part of the medication during critical illness.

**References**

[1] Casaer MP, Ziegler TR. Nutritional support in critical illness and recovery. *Lancet Diabetes*

- Endocrinol. 2015; 3(9):734-45.
- [2] Hooper MH, Marik PE. Controversies and misconceptions in intensive care unit nutrition. *Clin Chest Med.* 2015; 36(3):409-18.
- [3] Preiser J-C, Taccone FS. Nutrition in critically ill patients: where do we stand? *Minerva Anestesiol.* 2016; 82(8):908-13.
- [4] Amiri Farahani L, Heidari T, Narenji F, Asghari Jafarabadi M, Shirazi V. Relationship between premenstrual syndrome with body mass index among university students. *J hayat.* 2012; 17(4):85-95.
- [5] Jensen GL, Compher C, Sullivan DH, Mullin GE. Recognizing malnutrition in adults: definitions and characteristics, screening, assessment, and team approach. *J Parenter Enteral Nutr.* 2013; 37(6):802-7.
- [6] Preiser J-C, van Zanten AR, Berger MM, Biolo G, Casaer MP, Doig GS, et al. Metabolic and nutritional support of critically ill patients: consensus and controversies. *Critical care.* 2015;19(1):35.
- [7] Tian F, Gao X, Wu C, Zhang L, Xia X, Wang X. Initial energy supplementation in critically ill patients receiving enteral nutrition: a systematic review and meta-analysis of randomized controlled trials. *Asia Pac J Clin Nutr.* 2017;26(1):11.
- [8] Weijts PJ, Looijaard WG, Beishuizen A, Girbes AR, Oudemans-van Straaten HM. Early high protein intake is associated with low mortality and energy overfeeding with high mortality in non-septic mechanically ventilated critically ill patients. *Critical care.* 2014;18(6):701.
- [9] Mondal MK, Jana TK, Giri S, Roy H. Height prediction from ulnar length in females: a study in Burdwan district of West Bengal (regression analysis). *J Clin Diagn Res: JCDR.* 2012;6(8):1401.
- [10] Harvey SE, Parrott F, Harrison DA, Bear DE, Segaran E, Beale R, et al. Trial of the route of early nutritional support in critically ill adults. *N Engl J Med.* 2014;371(18):1673-84.
- [11] Guerra R, Fonseca I, Pichel F, Restivo M, Amaral T. Hand length as an alternative measurement of height. *Eur J Clin Nutr.* 2014;68(2):229-33.
- [12] Taylor BE, McClave SA, Martindale RG, Warren MM, Johnson DR, Braunschweig C, et al. Guidelines for the provision and assessment of nutrition support therapy in the adult critically ill patient: Society of Critical Care Medicine (SCCM) and American Society for Parenteral and Enteral Nutrition (ASPEN). *Critical care medicine.* 2016;44(2):390-438.
- [13] Blaser AR, Starkopf J, Alhazzani W, Berger MM, Casaer MP, Deane AM, et al. Early enteral nutrition in critically ill patients: ESICM clinical practice guidelines. *Intensive Care Med.* 2017;43(3):380-98.
- [14] Mulherin DW, Cogle SV. Updates in Nutrition Support for Critically Ill Adult Patients. *Hosp Pharm.* 2017;52(1):17-26.
- [15] McClave S, Taylor B, Martindale R, Warren M, Johnson D, Braunschweig C. Guidelines for the Provision and Assessment of American Society for Parenteral and Enteral Nutrition (ASPEN). *Society of Critical Care Medicine. JPEN J Parenter Enteral Nutr.* 2016;40(2):159-211.
- [16] Heyland DK, Dhaliwal R, Jiang X, Day AG. Identifying critically ill patients who benefit the most from nutrition therapy: the development and initial validation of a novel risk assessment tool. *Crit Care.* 2011;15(6):R268.
- [17] Heyland DK, Dhaliwal R, Wang M, Day AG. The prevalence of iatrogenic underfeeding in the nutritionally 'at-risk' critically ill patient: Results of an international, multicenter, prospective study. *Clin Nutr.* 2015;34(4):659-66.
- [18] Nachvak M, Hedayati S, Hejazi N, Motamedi Motlagh A, Shafizade A, Shojaee M. Nutritional assessment in ICU patients with enteral feeding in Amol hospitals. *Razi J Med Sci.* 2018;24(10):92-104.
- [19] Ahsan B, Khaledi S. Knowledge and mortality of ICU patients in Tohid Hospital of Sanandaj in 2001. *Med Sci Kordestan Univ.* 2005;9:20-5.
- [20] Yoosef-zadeh C, Shab-Bidar S, Dehnadi A, Ahmadi D. Nutritional support in head damage patients in ICU. *Gorgan Univ Med Sci.* 2008;3:53-8.
- [21] Knaus WA, Draper EA, Wagner DP, Zimmerman JE. APACHE II: a severity of disease classification system. *Crit care med.* 1985;13(10):818-29.
- [22] Dewar DC, White A, Attia J, Tarrant SM, King KL, Balogh ZJ. Comparison of postinjury multiple-organ failure scoring systems: Denver versus Sequential Organ Failure Assessment. *J Trauma Acute Care Surg.* 2014;77(4):624-9.
- [23] Rahman A, Hasan RM, Agarwala R, Martin C, Day AG, Heyland DK. Identifying critically-ill patients who will benefit most from nutritional therapy: further validation of the "modified NUTRIC" nutritional risk assessment tool. *Clin Nutr.* 2016;35(1):158-62.
- [24] Hejazi N, Mazloom Z, Zand F, Rezaianzadeh A, Amini A. Nutritional assessment in critically ill patients. *Iran J Basic Med Sci.* 2016;41(3):171.
- [25] Kubrak C, Jensen L. Malnutrition in acute care patients: a narrative review. *Int J Nurs Stud.* 2007;44(6):1036-54.
- [26] Kyle ÚG, Kossovsky MP, Karsgaard VL, Pichard C. Comparison of tools for nutritional assessment and screening at hospital admission: a population study. *Clin Nutr.* 2006;25(3):409-17.
- [27] Tsai J-R, Chang W-T, Sheu C-C, Wu Y-J, Sheu Y-H, Liu P-L, et al. Inadequate energy delivery during early critical illness correlates with increased risk of mortality in patients who survive at least seven days: a retrospective study. *Clin Nutr.* 2011;30(2):209-14.
- [28] Leiva Badosam E, Badia Tahull M, Virgili Casas N, Elguezabal Sangrador G, Faz Méndez C, Herrero Meseguer I, et al. Cribado de la desnutrición hospitalaria en la admisión: la desnutrición aumenta la mortalidad y la duración de la estancia

- hospitalaria. *Nutr Hosp*. 2017;34(4):907-13.
- [29] Martindale R, McClave S, Vanek V, McCarthy M, Roberts P, Taylor B. Guidelines for the provision and assessment of nutrition in nutrition support therapy in the adult critically ill patient: Society of Critical Care Medicine (SCCM) and American Society for Parenteral and Enteral Nutrition (ASPEN). *JPEN J Parenter Enteral Nutr*. 2009;33(3):277-316.
- [30] Rice TW, Mogan S, Hays MA, Bernard GR, Jensen GL, Wheeler AP. A randomized trial of initial trophic versus full-energy enteral nutrition in mechanically ventilated patients with acute respiratory failure. *Crit Care Med*. 2011;39(5):967.
- [31] Villet S, Chiolero RL, Bollmann Md. Negative impact of hypocaloric feeding and energy balance on clinical outcome in ICU patients. *clin Nutr*. 2005; 24:502-9.