

Original Article

Sleep Symptoms and Polysomnographic Patterns of Obstructive Sleep Apnea in Obese Children

How to Cite This Article: Tavasoli A, Jalilolghadr Sh, Lotfi Sh. Sleep symptoms and polysomnographic patterns of obstructive sleep apnea in obese children. *Iran J Child Neurol.* Winter 2016; 10 (1):14-20.

Azita TAVASOLI MD¹,
Shabnam JALILOLGHADR MD²,
Shiva LOTFI MD³

1. Pediatric Neurology Department, Ali-Asghar Children's Hospital, Iran University of Medical Sciences, Tehran, Iran

2. Pediatrics and Sleep Disorder Department, Qods Hospital, Qazvin University of Medical Sciences, Qazvin, Iran

3. General Physicians, Qazvin University of Medical Sciences, Qazvin, Iran

Corresponding Author:
Tavasoli A. MD
N193, Division of Pediatric Neurology, Children's Ali Asghar Hospital, Shahid Dastgerdi St, 1919816766, Tehran, Iran
Tel: +98 21 22220041-5,
Fax: +98 21 22220063,
Email: azita_tavasoli@yahoo.com

Received: 7-Oct-2014
Last revised: 9-Aug-2015
Accepted: 5-Nov-2015

Abstract

Objective

This study was conducted to investigate the sleep symptoms and polysomnographic patterns of obstructive sleep apnea in overweight and obese children.

Materials & Methods

Overweight or obese children aging 6-18 yr old referred during 2010 to Endocrinology Clinic of Ghods Hospital in Ghazvin, central Iran were enrolled in the study. Polysomnography was done for the diagnosis of obstructive sleep apnea and the BEARS and Children's Sleep Habits questionnaires were used to survey sleep behaviors.

Results

We enrolled 30 children (14 males, 16 females). Twenty-one cases had body mass index (BMI) >95% and 9 had 85% <BMI<95%. Respiratory disturbance in polysomnography was seen in 90% of cases. Symptoms included snoring 18 (60%); frequent awakening 17 (56.6%); nocturnal sweating 15 (50%); daytime sleepiness 12 (40%); sleep talking 10 (33.3%); bedtime resistance 9 (30%); nightmares 8 (26.6%); waking up problems 6 (20%); sleep walking 6 (20%); difficult breathing 4 (13.3%); bedwetting 3 (10%) and sleep onset delay 2 (6.06%). Severe, moderate and mild apnea – hypopnea Index (AHI) were seen in 12, 9 and 6 subjects, respectively. A significant Pearson correlation was found between the BMI values and sleep latency.

Conclusion

Prevalence of obstructive sleep apnea is high among overweight and obese children. Physicians should be familiar with its manifestations and consider polysomnography as an invaluable diagnostic test. There was no relation between the degree of obesity and severity of obstructive sleep apnea.

Keywords: BEARS questionnaire; Polysomnography; Obstructive apnea; Obesity; Sleep apnea- hypopnea syndrome

Introduction

Sleep-disordered breathing (SDB) is a constellation of breathing disorders characterized by intermittent hypoxia, hypercapnia and sleep fragmentation. Obstructive sleep apnea hypopnea syndrome (OSAHS) is usually considered as an extreme of SDB (1). The underlying cause of SDB in children remains unclear and seems to be multifactorial. The most common cause is adenotonsillar hypertrophy but other factors such as obesity, anatomic or functional defect of upper or lower

airway and genetic factors are contributed (1). The prevalence of OSAHS in obese children has been reported 13% to 59% versus 2% in general pediatric population (2, 3). Obesity predisposes children to OSAHS through the adenotonsillar hypertrophy and local fat deposition within the airway luminal walls and upper airway narrowing. In addition, it may have direct effect on respiratory muscles and ventilatory control (1). On the other hand, for children, the prevalence of obesity is continuously increasing in both developed and developing countries, besides, it is considered as a significant public health problem (4).

Based on an Iranian national survey in 2007, the prevalence of overweight and obesity amongst children between 6 and 18 years was 10.1% and 4.79%, respectively (5). Children with OSAHS usually have a history of snoring, observed apneas, arousals, sweating, enuresis during sleep and restless sleep at nights as well as daytime sleepiness. Morning headaches, concentration difficulties, memory loss and educational failure are other daytime symptoms (6). Lack of early detection and treatment of OSAHS can have serious adverse effects on growth velocity, behavioral and cognitive status (7), quality of life (8), use of health care services (9) and resulting in more frequent cardiovascular and chronic illness comorbidities (10). Therefore, it is mandatory to monitor the symptoms and diagnosis of OSAHS in the suspicious children such as obese and overweight ones. Polysomnography (PSG) that consists of a concurrent recording of multiple physiologic parameters related to sleep and wakefulness is often considered as the gold standard test for the diagnosis of OSAHS that determine the severity of the disease. PSG can directly detect and quantify the number of respiratory events and the resultant hypoxemia as well as arousals related to the respiratory events.

Although obesity has been recognized as a risk factor for adult OSAHS, there are more conflicts about this correlation in children. In this study, we attempted to detect the clinical symptoms and PSG pattern of obstructive sleep apnea in overweight and obese children.

Materials & Methods

Overweight or obese children aging 6-18 years old

referred to Endocrinology Clinic of Ghods Hospital in Ghazvin, central Iran during 2010 were enrolled in the study. Children with upper airway disease, perennial allergy or asthma, genetic diseases with high susceptibility to sleep apnea such as neuromuscular disease, Pierre Robin or Down syndrome were excluded. Standing height was measured in millimeter using a wall-mounted stadiometer and weight through a Seca balance to the nearest 0.1 kg. The results expressed in the form of body mass index (BMI) reference charts and age-related prevalence for boys and girls. Children with 85% <BMI<95% and BMI>95% were considered overweight and obese, respectively.

PSG was conducted for all children under the observation of a trained nurse. One of the parents accompanied their child through the night. The bedroom had a proper temperature and lighting, adequate ventilation, minimal sources of noise and a comfortable bed. A 6-channel computerized PSG was developed with leads for an oronasal flow cannula, a thoracoabdominal strain gauge and electromyogram, pulse oximeter, body-position sensor, electroencephalogram and static charge-sensitive bed. After calibration of the device, along with turning off lights, computers began recording data. The measured parameters included sleep onset latency, sleep efficiency, sleep stages (N1, N2, N3, REM), arousal index (AI), apnea hypopnea index (AHI), mean saturation O₂ (SaO₂), total sleep time and waking after sleep onset. The BEARS questionnaire was used in children with PSG evidences of OSAHS to investigate their sleep symptoms. This questionnaire surveys five basic sleep domains; B: bedtime problems; E: excessive daytime sleepiness; A: awakenings during the night; D: duration and regularity of sleep/wake cycles; and S: snoring (12). The Persian version of this questionnaire is prepared and its validity and reliability is confirmed (13).

The second questionnaire we used for more details was the Children's Sleep Habits Questionnaire (CSHQ). Parents were asked to recall the sleep behavior of their child over a typical week. CSHQ consists of 33 items in eight subscales about sleep duration, bedtime resistance, sleep-onset delay, daytime sleepiness, night awakenings, parasomnias, sleep-disordered breathing, and sleep anxiety. This questionnaire is also translated

into Persian (14,15). Collected data were scored and analyzed by a pediatric sleep specialist using the American Academy of Sleep Medicine criteria for children (16). Apnea-hypopnea index (AHI) - which expresses the number of breathing abnormalities per hour during sleep - was used to assess the severity of sleep apnea. Apnea defined as the absence of both nasal and oral airflow, and hypopnea as a reduction of 50% or more in the airflow both lasting for more than two respiratory cycles. Respiratory events for at least 10 seconds accompanying by at least a 3% decrease in oxygen saturation and/or an associated arousal were considered significant. Obstruction was defined as the presence of apnea with continuous respiratory effort recorded by thoracoabdominal strain gauge and electromyogram. A central apnea was defined as the absence of both the airflow and breathing efforts. Mixed apneas begin with a central component and end with an obstructive one (17). The number of apneas/hypopneas event/h was used to evaluate the severity of OSAHS as divided into mild, moderate and severe if AHI number was 1-4, between 5-10 and >10, respectively (16). Data were presented as mean and standard deviation. The relationship between the SDB variables and obesity values was analyzed using Pearson correlation coefficients.

A written informed consent was taken from parents. The study was approved by the Ethics Committee of Tehran University of Medical Sciences, Tehran, Iran.

Results

Overall, 30 children (14 males, 16 females) were enrolled. The overall mean age was 11.18 ± 2.87 yr (6-17 yr). Twenty-one cases had BMI >95% and 9 had 85% <BMI <95%. The mean BMI was $26.18 \pm 10.50\%$. Based on questionnaires, children slept 9h, 25 min +/- 1h, in weekdays and 10 h, 27 min +/- 1 h, 51 min at the weekend. Sleep symptoms of OSAHS included snoring 18 (60%); frequent awakening 17 (56.6%); nocturnal sweating 15 (50%); daytime sleepiness 12 (40%); sleep talking 10 (33.3%); bedtime resistance 9 (30%); nightmares 8 (26.6%); waking up problems 6 (20%); sleep walking 6 (20%); difficult breathing 4 (13.3%); bedwetting 3 (10%) and sleep onset delay 2 (6.06%) subjects.

PSG monitoring: Total recording time was 457.56 ± 60.61 minutes and total sleep time 435.93 ± 55.51 min with sleep efficiency (the amount of efficient sleep) of $75.71 \pm 15.42\%$ and sleep latency of 13.57 ± 18.65 min. Sleep architecture consisted of $10.20 \pm 6.64\%$ stage N1, $49.80 \pm 9.34\%$ stage N2, $20.49 \pm 9.74\%$ stage N3 (three stages of non REM sleep) and $11.80 \pm 4.92\%$ rapid-eye-movement (REM) sleep. AI was $19.86 \pm 20.22/h$ (Short-term awakenings at night with 3-15 seconds duration that occurred spontaneously or due to respiratory events, snoring or body movements). AI was normal in 11 and more than 11/h in 19 subjects. Table 1 shows the sleep structure in obese children.

Severe, moderate and mild AHI were seen in 12, 9 and 6 subjects respectively. The patients' hypopnea index was $16.30 \pm 17.36/hour$. The specific obstructive AHI was measured $33.13 \pm 52.20/h$, central AHI $3.53 \pm 3.62/h$ and mixed AHI $1.96 \pm 3.85/h$. The patients' overall respiratory disturbances index (RDI) was $12.55 \pm 21.57/h$. The mean baseline arterial oxygen saturation was $96.77 \pm 1.29\%$ and $95.3 \pm 5.7\%$ during awaking and REM sleep stage, respectively. The lowest arterial oxygen saturation was $87.43 \pm 6.7\%$. Table 2 shows the breathing indices in obese children.

There was a significant Pearson correlation between the BMI values and sleep latency (0.421) ($P=0.02$) (Figure 1) but no correlation was seen between BMI values and total sleep time, sleep efficiency, AHI, RDI, arousal index and average O₂sat during REM.

Discussion

Cross-sectional studies around the world revealed short sleep duration might be associated with the development of obesity in children, which is the underlying factor for sleep apnea. The sufficient sleep duration recommended for children is varied regarding their age groups (18). In the current study, sleep duration based on questionnaires was 9 h and 25 min. Iranian children have shorter night sleep duration than expected to normal ranges for their age group (11, 19). This may be due to cultural sleeping habits and poor sleep hygiene (14). Short sleep duration results in daytime sleepiness and fatigue and decreased physical activity. Furthermore, it affects the body metabolism including immune and endocrine systems function, particularly glucose and insulin metabolism

that induces changes in the center of appetite followed by increased need to eat and calorie intake (moraba 20). Both of these mechanisms lead to increased BMI that predispose children to sleep apnea.

Regarding the major signs and symptoms, different prevalence of sleep problems has been reported (21, 22). Similar to the report of American Thoracic Society that dyspnea, restless sleep and night sweats were the other common symptoms, respectively, snoring was the most prevalent symptom of OSAHS in our study (23). Overweight and nocturnal enuresis in children were significantly associated with OSAHS especially when they display other accompanied symptoms (24). In addition, Brooks et al. showed higher risk for enuresis in children with RDI>1 and pertained it to the effects of obstructive sleep apnea on arousal response, bladder pressure, or urinary hormone secretion (25).

The prevalence of bedwetting in our study was much lower that may be partly due to the upper age group of children. In contrast, nocturnal sweating was found in 50% of our children that propound it as a diagnostic factor. Half of OSA patients had nocturnal sweating in the neck and upper body area (26). Arnardottir et al. showed threefold higher rate of nocturnal sweating in OSA patients comparing general population and stated it is related to low REM sleep percentage, lower sleep quality and more daytime sleepiness in patients. They propounded nocturnal sweating as a marker of increased risk of OSA in general population (27). Nocturnal sweating may be due to labored breathing (20), however, causes, evaluations and management of nocturnal sweating are still of matter of controverting (28). In a study, daytime sleepiness was observed in 13.5% of 37 obese children, which was much lower than our study (29). Excessive daytime sleepiness is a cardinal symptom in adolescents but hyperactivity or inattention is more obvious in preadolescent with OSAS (20).

Refaey et al. (30) reported the mean sleep efficacy and sleep latency in normal weight children as 83.46 and 39.06, while our results in obese children were much lower in both items. In addition, arousal index in our study was much higher than that of their normal group (19.86/h against 1.62/h). The lowest oxygen saturation in our study was 87.43, which was much lower

than reported earlier as 91.4% (30). However, sleep architecture reported there (30), as the mean REM and NREM as 8.6% and 92.6% in normal children were similar to our findings (11.80% and 80.40%).

Respiratory events in children more than 1/h, is abnormal and can lead to the clinical signs and symptoms (daily growth failure, attention deficit, behavioral disorders etc.) due to changes in sleep architecture and arterial oxygenation that affect prefrontal executive function (48 moraba). The mean RDI in our study was 12.55/h, which is much higher than 1.37/h reported in normal children (30). It seems that our results are similar to the report of Dayyat et al. in 105 obese children aged over 7 yr as the mean oxygen saturation was $92.95 \pm 10.5\%$, arousal index $15.89 \pm 2.8/h$ and RDI $9.1 \pm 1.1/h$ (31). In obese children, airway size is reduced and its resistance is increased in the favor of excessive fat deposition around the upper airway. In addition, lung volumes and central ventilatory drive both are reduced in obese children that lead to more decrement of the upper airway patency (20). Obesity is the most important risk factor for OSAS in 2 – 18 yr old children, with an odds ratio of 4.5 (32).

In the current study, 90% of children had OSAHS in PSG. Compared to similar studies, this rate is much higher, which could be due to our methodological limitations. Although our subjects were enrolled from an obesity clinic and not from an otolaryngology or sleep clinic, however, presumably only children with suspected OSAHS were referred.

The great limitation of this study is the lack of a control normal weight population for comparing with obese children. In addition, despite PSG is the standard test for the diagnosis of OSAHS, the high cost and lack of parental consent resulted in limited number of our samples

In conclusion, prevalence of OSAHS is high among overweight and obese Iranian children. Physicians should be familiar with the related manifestations and consider PSG as an invaluable diagnostic test. We did not find any relation between the degree of obesity and severity of OSAHS but A significant Pearson correlation was found between BMI values and sleep latency (0.421). Monitoring the symptoms of sleep apnea in obese children is recommended and if possible, PSG

should be done for confirmation. Therefore, proceeding programs to increase the knowledge of parents and physicians is suggested.

Acknowledgments

This manuscript is prepared according to the thesis of Shiva Lotfi with code number of 5000 documented at Tehran University of Medical Sciences in 2011 for acquisition of general physician degree.

Authors' Contributions:

Azita Tavasoli drafted, and prepared the manuscript. Shabnam Jalilolghadr prepared design of the manuscript, data analysis, and interpretation.

Shiva Lotfi performed the study, acquired the data and analyzed them.

Conflict of interest: Non declared

Table 1. Sleep Structure of Obese Children

	Minimum	Maximum	Mean	Std. Deviation	Variance
Sleep Efficiency (%)	41.10	97.50	75.7100	15.42596	237.960
Sleep Latency (minutes)	.00	84.70	13.5700	18.65962	348.181
N1 stage (%)	.90	27.70	10.2000	6.64379	44.140
N2 stage (%)	28.10	72.00	49.8067	9.34499	87.329
N3 stage (%)	4.00	42.60	20.4933	9.74807	95.025
REM stage (%)	.50	20.60	11.8000	4.92635	24.269
Arousal Index (time/hour)	.00	64.10	19.8667	20.22444	409.028

N1 stage: The first stage of non- REM sleep

N3 stage: The third stage of non- REM sleep

N2 stage: The second stage of non- REM sleep

REM stage: rapid-eye-movement stage of sleep

Table 2. Breathing indices of obese children

	Minimum	Maximum	Mean	Std. Deviation	Variance
Apneas Total central (times/h)	0.00	00.14	3.5333	3.62685	13.154
Apneas Total mixed (times/h)	0.00	00.18	1.9667	3.85499	14.861
Apneas Total obstructive (times/h)	0.00	00.270	33.1333	52.20763	2725.637
Hypopneas Total (times/h)	0.00	53.00	16.3000	17.36852	301.666
Total Events RDI (times/h)	0.20	93.90	12.5567	18.70581	349.907
AHI Total Back (times/h)	0.00	240.00	24.2267	44.92667	2018.405
Average o2 While Awake (%)	92	98.80	96.7733	1.29640	1.681
Average o2 While in REM (%)	65.90	99.20	95.0367	5.76078	33.187
Approximate Minimum o2 value (%)	73	96	87.4333	6.72455	45.220

RDI : Respiratory Disturbances Index

REM: Rapid eye movement sleep

AHI: Apnea Hypopnea Index




Fig 1. Correlation between the BMI values and sleep latency (0.421) ($P=0.02$)

References

- Marcus CL. Pathophysiology of childhood obstructive sleep apnea: current concepts. *Respiration Physiol* 2000;119 (2):143-54.
- Verhulst SL, Van Gaal L, De Backer W, Desager K. The prevalence, anatomical correlates and treatment of sleep-disordered breathing in obese children and adolescents. *Sleep Med Rev* 2008;12 (5):339-46.
- Marcus CL. Sleep-disordered breathing in children. *Am J Respir Crit Care Med* 2001;164 (1):16-30.
- Lau DC.W., Doukeits JD, Morrison KM, Hraminak IM, et al. 2006 canadian clinical practical guidelines on the management and prevention of obesity in adults and children. *Canadian Med Ass J* 2007;176 (8),suppl 2: 1-14.
- Kelishadi R, Ardalan G, Gheiratmand R, Majdzadeh R, Hosseini M, Gouya M, et al. Thinness, overweight and obesity in a national sample of Iranian children and adolescents: Caspian Study. *Child Care Health Dev* 2008;34 (1):44-54.
- Azagra-Calero E, Espinar-Escalona E, Barrera-Mora JM, Llamas-Carreras JM, Solano-Reina E. Obstructive sleep apnea syndrome (OSAS). Review of the literature. *Med Oral, Patología Oral y Cirugía Bucal* 2012;17 (6):e925.
- Kohler MJ, Lushington K, Kennedy JD. Neurocognitive performance and behavior before and after treatment for sleep-disordered breathing in children. *Nature Sci Sleep* 2010;2:159.
- Jackman AR, Biggs SN, Walter LM, Embuldeniya US, Davey MJ, Nixon GM, et al. Sleep disordered breathing in early childhood: quality of life for children and families. *Sleep* 2013;36 (11):1639-46.
- Reuveni H, Simon T, Tal A, Elhayany A, Tarasiuk A. Health care services utilization in children with obstructive sleep apnea syndrome. *Pediatrics* 2002;110 (1):68-72.
- Amin RS, Kimball TR, Kalra M, Jeffries JL, Carroll JL, Bean JA, et al. Left ventricular function in children with sleep-disordered breathing. *Am J Cardiol* 2005;95 (6):801-4.
- Jalilolghadr.SH. *Pediatrics Sleep Disorders*. 1 ed: Qazvin Medical University; 2011: 12.
- Owens JA, Dalzell V. Use of the 'BEARS' sleep screening

- tool in a pediatric residents' continuity clinic: a pilot study. *Sleep Med* 2005;6 (1):63-9.
13. Mohammadi M, Amintehran E, Farhad M, Ghalehbandi MRA, Shoaee S, Ghalehbaghi B. Reliability and validity of persian version of "BEARS" pediatric sleep questionnaire. *Indian J Sleep Med* 2008;3 (1): 14-19.
 14. Shoghy M, Khangary S, Farmany F, Hosseini F. Sleep habits of school age children. *Iran J Nurs* 2005;18 (41):131-8.
 15. Rafii F, Oskouie F, Shoghi M. The association between sleep and injury among school-aged children in Iran. *Sleep Disord* 2013;2013.
 16. Grigg-Demberger M, Respiratory rules for children. In: Iber C, Medicine AAoS. *The AASM manual for the scoring of sleep and associated events: rules, terminology and technical specifications*. 1st ed : Westchester, Illinois : American Academy of Sleep Medicine; 2007, p:48-49.
 17. Somers VK, Dyken ME, Clary MP, Abboud FM. Sympathetic neural mechanisms in obstructive sleep apnea. *J Clin Investig* 1995;96 (4):1897.
 18. Chen X, Beydoun MA, Wang Y. Is Sleep Duration Associated With Childhood Obesity? A Systematic Review and Meta - analysis. *Obesity* 2008;16 (2):265-74.
 19. Javadzadeh M, Hashemi Z, Roudbari M, Mahvelati F, Jalilolghadr S. Sleep patterns and sleep disorders in primary school children in Qazvin, Iran. *Iran J Child Neurol* 2008;2 (4):15-9.
 20. Jung Chang S, Young Chae K. Obstructive sleep apnea syndrome in children: Epidemiology, pathophysiology, diagnosis and sequelae. *Korean J Pediatr* 2010;53 (10):863-871.
 21. Ali N, Pitson D, Stradling J. Natural history of snoring and related behaviour problems between the ages of 4 and 7 years. *Arch Dis Childhood* 1994;71 (1):74-6.
 22. Blunden S, Chervin RD. Sleep, performance and behaviour in Australian indigenous and non-indigenous children: An exploratory comparison. *J Paediatr Child Health* 2010;46 (1-2):10-6.
 23. Loughlin G, Brouillette R, Brooke L, Carroll J, Chipps B, England S, et al. Standards and indications for cardiopulmonary sleep studies in children. *Am J Respirato Critical Care Med* 1996;153 (2):866-78.
 24. Barone JG, Hanson C, DaJusta DG, Gioia K, England SJ, Schneider D. Nocturnal enuresis and overweight are associated with obstructive sleep apnea. *Pediatrics* 2009;124 (1):e53-e9.
 25. Brooks LJ, Topol HI. Enuresis in children with sleep apnea. *J Pediatr* 2003 May;142 (5):515-8.
 26. Guilleminaault C, Bassiri A. Clinical features and evaluation of obstructive sleep apnea-hypopnea syndrome and upper airway resistance syndrome. In: Kryger MH, Roth T, Dement WC, eds. *Principles and practice of sleep medicine*. 4th edn. Philadelphia, PA: Elsevier Sanders, 2005. 1043-52.
 27. Arnardottir E, Janson Ch, Bjornsdottir E, Benediktsdottir B, Juliusson S, Kuna S, et al. Nocturnal sweating - a common symptom of obstructive sleep apnoea: the Icelandic sleep apnoea cohort. *BMJ Open* 2013;3:e002795. doi:10.1136/bmjopen-2013-002795.
 28. Mold JW, Holtzclaw BJ, McCarthy L. Night sweats: a systematic review of the literature. *J Am Board Fam Med* 2012;25:878-93
 29. Xu Z, Jiaqing A, Yuchuan L, Shen K. A case-control study of obstructive sleep apnea-hypopnea syndrome in obese and nonobese chinese children. *CHEST J* 2008;133 (3):684-9.
 30. El-Refaey A, ElSayed R, Sarhan A, Bakr A, Hammad A, Elmougy A, et al. Sleep quality assessment using polysomnography in children on regular hemodialysis. *Saudi J Kidney Dis Transplant* 2013;24 (4):714.
 31. Dayyat E, Kheirandish-Gozal L, Capdevila OS, Maarafeya MM, Gozal D. Obstructive Sleep Apnea in Children Relative Contributions of Body Mass Index and Adenotonsillar Hypertrophy. *CHEST J* 2009;136 (1):137-44.
 32. Redline S, Tishler PV, Schluchter M, Aylor J, Clark K, Graham G. Risk factors for sleep-disordered breathing in children. Association with obesity, race, and respiratory problems. *Am J Respir Crit Care Med* 1999;159:1527-32.