



Evaluation and Comparison of Body Mass Index and Albumin Level in Patients with Active Tuberculosis and Latent Tuberculosis Infection

Fatemeh Heidarneshad¹, Roghayeh Ghezelsofla¹, Aghigh Ziaeemehr², Amir Asnaashari³, Houshang Rafatpanah⁴, Abdolrahim Rezaee⁴, Kiarash Ghazvini⁵, Reza Basiri^{3*}

¹ Master of Immunology, Inflammation and Inflammatory Diseases Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

² Bachelors of Biology, Inflammation and Inflammatory Diseases Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

³ Pulmonologist, Lung Diseases Research Center, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

⁴ Immunologist, Inflammation and Inflammatory Diseases Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

⁵ Microbiologist, Antimicrobial Resistance Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

ARTICLE INFO

Article type:
Original article

Article history:
Received: 15 Aug 2015
Revised: 11 Apr 2016
Accepted: 1 May 2016

Keywords:
Albumin
Body Mass Index
Tuberculosis

ABSTRACT

Introduction: Limited data are available on the relationship between nutritional status and tuberculosis. The aim of this study was to evaluate and compare the body mass index (BMI) and serum albumin level in patients with active tuberculosis (ATB) and latent tuberculosis (LTB).

Materials and Methods: A cross-sectional study was conducted on 17 patients newly diagnosed with pulmonary TB who were referred in Iran, during September 2011 to March 2012 and 17 latent tuberculosis infection individuals. Standard method was performed to collect an early morning fasting blood sample for albumin (by the bromocresolgreen method). Also (BMI) was calculated as body weight divided by height squared (kg/m²).

Results: One-sample Kolmogorov-Smirnov test was used to check normal distribution data. The mean \pm Standard deviation (SD) for albumin in the patients and controls were 3.62 ± 0.56 and 4.68 ± 0.25 , respectively. BMI in the patients and controls were 19.46 ± 2.79 and 25.4 ± 3.46 , respectively. The serum albumin level was significantly lower in the patient group, compared to the control group ($P < 0.001$). BMI was significantly lower in the patient group, compared to the control group ($P < 0.001$).

Conclusion: Our findings demonstrated that BMI and serum Albumin were significantly lower in the active tuberculosis patients than latent tuberculosis groups.

► Please cite this paper as:

Heidarneshad F, Ghezelsofla R, Ziaeemehr A, Asnaashari A, Rafatpanah H, Rezaee A, Ghazvini K, Basiri R. Evaluation and Comparison of Body Mass Index and Albumin Level in Patients with Active Tuberculosis and Latent Tuberculosis Infection. J Cardiothorac Med. 2016; 4(2):437- 439

Introduction

Tuberculosis is one of the most important infectious diseases with a high global mortality rate, particularly in developing countries. According to World Health Organization, one-third of the world population is infected with *Mycobacterium tuberculosis*, while only 5-10% of infected individuals develop active tuberculosis

(ATB) with clinical symptoms (1-3).

Malnutrition is frequently reported in patients with pulmonary tuberculosis (4). Albumin facilitates the movement of various small molecules through the blood including bilirubin, calcium, progesterone and medicines. This protein plays an important role in restraining the blood fluid from

*Corresponding author: Reza Basiri, Lung Diseases Research Center, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran. Tel/Fax: +985138431252; Email: basirir@mums.ac.ir

© 2016 mums.ac.ir All rights reserved.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

leaking into tissues (5, 6). Considering its oncotic properties, albumin has been used in patients with cirrhosis and liver disease for the maintenance of vascular volume (7).

Considering the limited data on the nutritional status and serum albumin level in patients with ATB, we aimed to Evaluate BMI and serum Albumin serum albumin level in subjects with active tuberculosis (ATB) and latent tuberculosis (LTB).

Materials and Methods

The present cross-sectional study was conducted at Ghaem Hospital, Mashhad University of Medical Sciences, Mashhad, Iran, during September 2011 and March 2012. The study population consisted of 17 patients with recently diagnosed pulmonary tuberculosis and 17 control subjects with LTB (with positive tuberculin skin test). The subjects were matched in terms of age and sex.

The diagnosis of tuberculosis was confirmed, based on positive microscopic findings for acid-fast bacillus in at least two early morning sputum specimens (8, 9) and positive *M. tuberculosis* culture on Löwenstein-Jensen medium after four weeks. The patient and control groups were interviewed, using a structured questionnaire including information related to the inclusion and exclusion criteria.

The exclusion criteria were as follows: 1) pregnancy, 2) acute or chronic liver diseases, 3) renal diseases, and 4) other active inflammatory conditions. Five ml blood samples were obtained from the subjects and serum samples were stored at -20 °C. Bromocresol green assay kit was applied to measure serum albumin level. Body mass index (BMI) was also calculated as body weight in kilograms divided by height in meters squared (kg/m^2).

Statistical Analysis

Normal distribution of the data was assessed by Kolmogorov-Smirnov test. Mean and standard deviation (SD) were calculated for reporting normally distributed data. Independent sample's *t*-test was used to assess the differences between patient and control groups in terms of normally distributed parameters. Moreover, Pearson's correlation test was applied to calculate the correlation coefficients. SPSS version 20 (Chicago, USA) was used for all statistical analyses. *P*-value less than 0.05 were considered statistically significant.

The present study was approved by the ethic committee of Mashhad University of Medical Sciences, Mashhad, Iran. Informed consent forms were obtained from subjects before the start of the study.

Results

The mean age of the patients and control groups was 49.65 ± 18.27 and 44.12 ± 10.20 years (range: 20-70 years), respectively. There was no statistically significant (*p* value = 0.697) difference between the two groups in terms of age or sex. The mean BMI in the patients and control groups was 19.46 ± 2.79 and 25.4 ± 3.46 kg/m^2 , respectively. BMI was significantly lower in the patient group, compared to the control group ($P < 0.001$).

The mean albumin level in the patients and control groups was 3.62 ± 0.56 and 4.68 ± 0.25 , respectively. The serum albumin level was significantly lower in the patient group, compared to the control group ($P < 0.001$). According to Pearson correlation test a positive correlation observed between BMI and albumin. ($P < 0.01$).

Discussion

Infection leads to a decline in serum albumin level in humans. Albumin is also required for the growth of *M. tuberculosis*. In the present study, subjects with the following conditions were excluded from the study: pregnancy, acute and chronic liver disease, renal diseases and other active inflammatory conditions. In our findings serum albumin level and BMI was significantly reduced in patients with ATB, compared to control subjects with LTB.

According to previous research, serum albumin, zinc and BMI were lower in patients with ATB, compared to healthy control subjects (4). The reported findings were consistent with previous research in England (10), India (11) and Japan (12). In ATB, the production of cytokines (e.g., interleukin-6 and tumor necrosis factor- α) and acute-phase proteins inhibits serum albumin production (13-15).

In the acute-phase response, leakage of transthyretin (prealbumin) and albumin through the vascular endothelium occurs and production of transthyretin by the liver is reduced (16). In a previous study, lipid, when bound to albumin, can be used as a source of carbon and promote the growth of *M. tuberculosis* (17).

In a recent study in India, BMI was significantly lower in patients with tuberculosis, compared to healthy control subjects (18). In fact, low BMI is a known risk factor for mortality (19). According to several studies, low BMI of patients with pulmonary tuberculosis may be due to anorexia, impaired absorption of nutrients or increased catabolism (4, 9, 19, 20). In recent studies, patients with pulmonary tuberculosis and human immunodeficiency virus (HIV) infection had significantly low zinc level, albumin

and BMI. Micronutrient supplementation for this group of patients improves their health, increases their weight and improves the efficacy of drug treatments (9, 21-23).

In conclusion, based on the obtained findings, BMI and serum albumin concentration in patients with ATB were lower, compared to control subjects with LTB.

Acknowledgments

The study was financially supported by the Vice Chancellor for Research at Mashhad University of Medical Sciences, Mashhad, Iran (grant No: 86276).

Conflicts of interest

The authors declare no conflicts of interest.

References

1. Pai-Dhungat JV, Parikh F. Centenary of discovery of Tubercle bacillus. *J Assoc Physicians India*. 2003; 51:668.
2. Deveci F, Ilhan N. Plasma malondialdehyde and serum trace element concentrations in patients with active pulmonary tuberculosis. *Biol Trace Elem Res*. 2003; 95:29-38.
3. Sapozhnikova NV, Skvortsova LA, Pavlova MV, Vishnevskii BI, Otten TF, Vasil'eva SN, et al. Pulmonary tuberculosis caused by *Mycobacterium tuberculosis* of different genotypes. *Probl Tuberk Bolezn Legk*. 2003; 10:13-5.
4. Karyadi E, Schultink W, Nelwan RH, Gross R, Amin Z, Dolmans WM, et al. Poor micronutrient status of active pulmonary tuberculosis patients in Indonesia. *J Nutr*. 2000; 130:2953-8.
5. Cantin AM, Paquette B, Richter M, Larivée P. Albumin-mediated regulation of cellular glutathione and nuclear factor kappa B activation. *Am J Respir Crit Care Med*. 2000; 162:1539-46.
6. Mendez DL, Jensen RA, McElroy LA, Pena JM, Esquerra RM. The effect of non-enzymatic glycation on the unfolding of human serum albumin. *Arch Biochem Biophys*. 2005; 444:92-9.
7. Arroyo V. Review article: albumin in the treatment of liver diseases--new features of a classical treatment. *Aliment Pharmacol Ther*. 2002; 16:1-5.
8. Middleton AM, Chadwick MV, Nicholson AG, Wilson R, Thornton DJ, Kirkham S, et al. Interaction between mycobacteria and mucus on a human respiratory tissue organ culture model with an air interface. *Exp Lung Res*. 2004; 30:17-29.
9. Ramakrishnan K, Shenbagarathai R, Kavitha K, Uma A, Balasubramanian R, Thirumalaikolundusubramanian P. Serum zinc and albumin levels in pulmonary tuberculosis patients with and without HIV. *Jpn J Infect Dis*. 2008; 61:202-4.
10. Onwubalili JK. Malnutrition among tuberculosis patients in Harrow, England. *Eur J Clin Nutr*. 1988; 42:363-6.
11. Saha K, Rao KN. Undernutrition in lepromatous leprosy. V. Severe nutritional deficit in lepromatous patients co-infected with pulmonary tuberculosis. *Eur J Clin Nutr*. 1989; 43:117-28.
12. Tsukaguchi K, Yoneda T, Yoshikawa M, Narita N, Enoki Y, Miyazaki R, et al. Interaction between interleukin-1 and tumor necrosis factor productions by peripheral blood monocytes and nutritional disturbance in active pulmonary tuberculosis. *Kekkaku*. 1991; 66:477-84.
13. Beisel WR. Infection-induced depression of serum retinol--a component of the acute phase response or a consequence? *Am J Clin Nutr*. 1998; 68:993-4.
14. Xing Z, Gauldie J, Cox G, Baumann H, Jordana M, Lei XF, et al. IL-6 is an antiinflammatory cytokine required for controlling local or systemic acute inflammatory responses. *J Clin Invest*. 1998; 101:311-20.
15. Gabay C, Kushner I. Acute-phase proteins and other systemic responses to inflammation. *N Engl J Med*. 1999; 340:448-54.
16. Fleck A. Clinical and nutritional aspects of changes in acute-phase proteins during inflammation. *Proc Nutr Soc*. 1989; 48:347-54.
17. Lynn M, Wilson AR, Solotorovsky M. Role of bovine serum albumin in the nutrition of *Mycobacterium tuberculosis*. *Appl Environ Microbiol*. 1979; 38:806-10.
18. Swaminathan S, Padmapriyadarsini C, Sukumar B, Iliayas S, Kumar SR, Triveni C, et al. Nutritional status of persons with HIV infection, persons with HIV infection and tuberculosis, and HIV-negative individuals from southern India. *Clin Infect Dis*. 2008; 46:946-9.
19. van Lettow M, Fawzi WW, Semba RD. Triple trouble: the role of malnutrition in tuberculosis and human immunodeficiency virus co-infection. *Nutr Rev*. 2003; 61:81-90.
20. van Lettow M, Kumwenda JJ, Harries AD, Whalen CC, Taha TE, Kumwenda N, et al. Malnutrition and the severity of lung disease in adults with pulmonary tuberculosis in Malawi. *Int J Tuberc Lung Dis*. 2004; 8:211-7.
21. Kaiser JD, Campa AM, Ondercin JP, Leoung GS, Pless RF, Baum MK. Micronutrient supplementation increases CD4 count in HIV-infected individuals on highly active antiretroviral therapy: a prospective, double-blinded, placebo-controlled trial. *J Acquir Immune Defic Syndr*. 2006; 42:523-8.
22. McGrath N, Bellinger D, Robins J, Msamanga GI, Tronick E, Fawzi WW. Effect of maternal multivitamin supplementation on the mental and psychomotor development of children who are born to HIV-1-infected mothers in Tanzania. *Pediatrics*. 2006; 117:e216-25.
23. Karyadi E, West CE, Schultink W, Nelwan RH, Gross R, Amin Z, et al. A double-blind, placebo-controlled study of vitamin A and zinc supplementation in persons with tuberculosis in Indonesia: effects on clinical response and nutritional status. *Am J Clin Nutr*. 2002; 75:720-7.