



Quality of Life in Elderly Iranian Population Using the QOL-brief Questionnaire: A Systematic Review

Zahra CHERAGHI¹, Amin DOOSTI-IRANI¹, Sima NEDJAT², Parvin CHERAGHI³,
*Saharnaz NEDJAT⁴

1. Dept. of Epidemiology and Biostatistics, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran
2. Knowledge Utilization Research Center, University of Social Welfare and Rehabilitation Sciences, Tehran, Iran
3. Dept. of Health Education and Promotion, School of Public Health, Hamadan University of Medical Sciences, Hamadan, Iran
4. Dept. of Epidemiology and Biostatistics, School of Public Health, Knowledge Utilization Research Center, Tehran University of Medical Sciences, Tehran, Iran

*Corresponding Author: Email: nejatsan@tums.ac.ir

(Received 17 Nov 2015; accepted 21 Apr 2016)

Abstract

Background: Due to the increase in the elderly population in different societies, their primary needs, physical and mental health, and quality of life, is important. This study aimed to estimate the overall mean score of the QOL based on the Qol-Brief questionnaire in the elderly population of Iran.

Methods: The international and national databases, including; Medline, Scopus, Science Direct, MagIran, and SID were searched up to Feb 2015. All studies, addressed the quality of life among the healthy Iranian elderly population using WHO-QOL-BRIEF questionnaire, were included. The random effect model was used for data analysis and the results were reported with a 95% confidence.

Results: Out of 2150 studies, seven studies involved 1366 elderly participants were included in the meta-analysis. The pooled mean score of total QOL was 60.1 ± 4.6 . The pooled mean score of quality of life for physical health was 55.13 [51.03, 59.22], for environmental was 51.80 [45.50, 58.10], for psychological was 56.68 [53.29, 60.08] and for the social relationship was 57.82 [55.79, 59.86]. The men had a better status and in two health domains, including; physical and psychological health.

Conclusion: The results of this study stress the necessity of attention to the quality of life in the domains of physical health (especially in women), and environmental health in the elderly population.

Keywords: Quality of life, Systematic review, Elderly

Introduction

Population ageing is a common phenomenon that occurs due to increased life expectancy, improvement of the health care services, and decreased mortality rate. This phenomenon is observed in nearly all countries throughout the world (1, 2). According to a report by WHO in 2013, the global elderly population (aged 60 yr and over) increased from 9.2% in 1990 to 11.7% in 2013 and will reach 21.1% in 2050 (3). According to the results of the National Population and Housing Census in 2011, the proportion

of the elderly population of Iran has been 8.19% (4). Due to the increase in the elderly population in different societies, it is important to consider their primary needs, physical and mental health, and improvement of the quality of life (QOL). According to the WHO definition, the QOL is the understanding people from their position in life in terms of culture, appraisal, goals, hope, standards, and preferences (5). Several different tools are available to measure the QOL such as the SF-36 questionnaire, WHO-QOL-Brief ques-

tionnaire, Lipad questionnaire, etc. (6-8). The WHO-QOL-Brief can assess the quality of life in a variety of societies and population groups (6). This questionnaire has been standardized in many countries, so we could be able to compare the quality of life in different populations. On the other hand, the age structure is rapidly changing in Iran, so we need to compare the QOL status between the Iranian elderly population and other developing countries.

So far, limited and sporadic studies have been conducted to investigate the QOL in the Iranian elderly population (9-16) and their results are inconsistent. This meta-analysis aimed to estimate the overall mean score of the QOL based on the Qol-Brief questionnaire in the elderly population of Iran.

Methods

Instrument

The WHOQOL-BREF is a standard tool for assessing the quality of life based on four domains of physical health, psychological health, social relationships and environmental health. The physical health domain includes energy and fatigue, pain and discomfort, sleep and rest. The psychological domain includes bodily image and appearance, negative feelings, positive feelings, self-esteem, thinking, learning, memory and concentration, religion /spirituality/personal beliefs. The social domain involves personal relationships, social support and sexual activity, and finally the environmental domain includes the financial resources, freedom, physical safety and security, health and social care, accessibility and quality of home environment, opportunities for acquiring new information and skills participation and opportunities for recreation/leisure, environment (pollution/noise/climate) and transport (6).

Searching

International and national databases were searched using following key words: "quality of life", "aging", "aged", "elderly" and "Iran". International databases including Medline, Scopus and Science Direct were searched up to Feb

2015. National databases including Science Information Database, MagIran, IranMedex and Irandoc were searched up to Feb 2015. Moreover, to obtain additional studies, the reference lists of all selected studies were scanned manually, and the authors of included studies were contacted.

Criteria for including studies

All cross-sectional studies addressed the QOL in the healthy Iranian elderly population using the WHO-QOL-BRIEF questionnaire were included irrespective of the time of the study and the language of the publication. The Iranian elderly population was considered the study population regardless of sex and age. The main outcome of interest was the mean scores of the domains of the quality of life.

Data extraction and management

Two authors (PCh. and ADI) screened the title and abstract of the retrieved citations independently; in the next stage, they reviewed the full text of the selected studies to extract the studies that met the inclusion criteria for this meta-analysis. Any disagreement between authors in the selection of the studies was resolved by discussion and adjudication of a third author. In the case of missing data, we contacted the corresponding author of the study. Two authors (PCh, ADI) extracted the following variables for data analysis: the year of publication, location of the study (city), mean or median of age, gender, residence of participants, sample size, mean score of the QOL based on domains (physical health, psychological, social and environmental) and their standard deviation.

Assessment risk of bias

Seven selected items from the STROBE checklist (17) were used for evaluating the quality of the included studies. These items included the following; 1) presence of the key elements of the study design; 2) the inclusion and exclusion criteria; 3) the outcome, i.e., quality of life; 4) calculation of the sample size; 5) the setting of the study (location and date); 6) the precision of the estimates, i.e., standard deviation or confidence

interval and 7) the statistical methods for data analysis. Studies that satisfied all mentioned criteria were classified as high quality, studies that did not meet one item were classified as moderated quality, and studies that did not meet more than one item were classified as low quality.

Assessment of heterogeneity

Statistical heterogeneity was explored using the chi-squared test at a significance level of 10%. I^2 was used for quantifying the heterogeneity across the included studies (18). The variance between the studies was estimated using tau-squared statistics (19).

Estimation of summary measures

A meta-analysis was performed to estimate the pooled mean score of the QOL among the elderly population. The pooled estimation of the QOL was reported for each domain of the WHO-QOL-BREF questionnaire separately. The inverse variance (IV) method was used for calculating the pooled estimations. Sub-group analysis

was accomplished according to the age group, sex, residence, and the quality of the included studies. Both Review Manager 5 (20) and Stata 11 (StataCorp, College Station, TX, USA) were employed for data analysis. The random effect model (21) was used for data analysis and the results were reported with a 95% confidence interval.

Results

Description of studies

We retrieved 2150 records; 470 references were excluded because of duplication, 1000 reference were excluded because they were not related to the objective of the review, and 653 references were excluded because they were not eligible to be included in the meta-analysis after checking the full text. Finally, seven articles (10, 11, 13, 15, 22-24) remained for meta-analysis (Fig. 1 and Table 1), including 1366 Iranian elderly participants with a mean age of 70.8 ± 2.3 yr.




Fig. 1: Flow chart of study identification process

Table 1: Characteristic of included studies in meta-analysis

Author	Year	Province	Sex	Resident	Sample Size	Habitat	Age
Ahangari	2007	Tehran	Male/Female	Nursing Home	523	City	68.08
Panaghi	2010	Tehran	Male/Female	Nursing Home/ Home	243	City	71.41
Hasani	2011	Tehran	Male/Female	Nursing Home	130	City/Rural	73.19
Khoosheh Mehri	2005	Alborz	Male/Female	Nursing Home	62	City	Not reported
Amirhosseini	2015	Tehran	Male/Female	Nursing Home/ Home	408	City/Rural	Not reported
Payahoo	2013	Tabriz	Male/Female	Home	184	City	69.4
Zaeri	2014	Tehran	Male/Female	Home	287	City	67.3

Quality assessment

Eighty percent of the studies (five studies) referred to all (seven) STROBE items while others (two studies; 20%) referred to six STROBE items. No study referred to five items or less.

Heterogeneity assessment

The heterogeneity of the studies was assessed using the Chi² test and the I² statistics. The results of Chi² test and I² for all domains of the WHO QOL-Brief were as follows: physical health (I² =99% and Chi²=1044.10), psychological (I² =96% and Chi² =224.16), environmental (I² =99% and Chi² =1044.10), social (I² =88% and Chi² =76.92), and overall (I²=100% and Chi² =1436.1) (Fig. 2).

Publication Bias: We assessed publication bias using the funnel plot as well as Begg's and Egger's tests. The results of Begg's and Egger's tests for physical health ($P=0.325$), psychological health ($P=0.106$) and social relationship ($P=0.295$) confirmed the absence of publication bias and for environmental ($P=0.024$) confirmed presence of publication bias.

The Mean Score of the Quality of life

The pooled mean score of the total QOL was 60.1 ± 4.6 . The pooled mean score of the quality of life was 55.13 [51.03, 59.22] for physical health, 51.80 [45.50, 58.10] for environmental health, 56.68 [53.29, 60.08] for psychological

health, and 57.82 [55.79, 59.86] for social relationships. Therefore, the maximum and minimum-pooled mean scores were obtained in social relationship and environmental domains, respectively.

Subgroup Analysis

We developed a sub-group analysis according to available variables (gender and the quality of reporting the studies). Men had a better status and a higher mean score in two health domains including physical health (62.6 vs. 49.7, $P<0.001$) and psychological health (62.4 vs. 55.4, $P<0.001$) as well as the total mean score of the quality of life (61.4 vs. 58.4, $P<0.001$). In addition, these mean differences were statistically significant. Women had a slightly better status than men in the two remaining health domains including social relationships (59.1 vs. 58.5, $P=0.82$) and the environmental domain (52.2 vs. 51.0, $P=0.85$). However, the mean differences were not statistically significant.

Studies with more risk of bias had reported a higher mean scores in comparison to studies with low risk of bias (including; in the psychological (61.55 vs. 55.29, $P<0.001$), in the environmental health (60.70 vs. 49.45, $P<0.001$), in social relationship (60.65 vs. 56.94, $P=0.77$)). Although both types of studies (low risk and high risk of bias) had equal status (58.33 vs. 58.34, $P=0.99$) (Table 2).




Fig. 2: Forest plot of pooled mean score of quality of life in four domains

Table 2: Subgroup analysis of mean score of quality of life and four domains according some factor affecting on quality of life

Domains	Variables	Categories	N* (%)	Pooled M	95% CI	I ²	P-value**
Physical Health	Gender	Male	4 (0.50)	62.59	[61.13, 64.04]	0.98	<0.001
		Female	4 (0.50)	49.71	[48.36, 51.07]	0.88	
	Quality of Study	Low Risk Bias	4 (0.40)	58.34	[58.18, 58.50]	0.95	0.99
Psychological	Gender	Moderate Risk of Bias	6 (0.60)	58.33	[56.94, 59.72]	0.100	
		Male	4 (0.50)	62.45	[60.92, 63.98]	0.93	<0.001
	Female	4 (0.50)	55.90	[54.54, 57.25]	0.98		
Social Relationship	Quality of Study	Low Risk Bias	8 (0.80)	55.29	[52.28, 58.30]	0.89	<0.001
		Moderate Risk of Bias	2 (0.20)	61.55	[56.28, 66.82]	0.94	
	Gender	Male	4 (0.50)	58.55	[56.97, 60.12]	0.92	0.82
Environmental	Quality of Study	Female	4 (0.50)	59.07	[57.67, 60.48]	0.69	
		Low Risk Bias	8 (0.80)	56.97	[53.94, 60.01]	0.90	0.77
	Moderate Risk of Bias	2 (0.20)	60.65	[59.12, 62.19]	0.29		
Total Mean of Quality of Life	Gender	Male	4 (0.50)	50.99	[40.81, 61.17]	0.98	0.85
		Female	4 (0.50)	52.20	[44.79, 59.61]	0.97	
	Quality of Study	Low Risk Bias	8 (0.80)	49.45	[44.59, 54.32]	0.97	<0.001
Total Mean of Quality of Life	Quality of Study	Moderate Risk of Bias	2 (0.20)	60.70	[59.49, 61.92]	0.00	
		Male	4 (0.50)	61.37	[45.61, 77.13]	0.97	0.30
	Female	4 (0.50)	58.92	[47.93, 69.90]	0.99		
Total Mean of Quality of Life	Quality of Study	Low Risk Bias	4 (0.50)	51.37	[50.03, 52.71]	0.83	<0.001
		Moderate Risk of Bias	4 (0.50)	71.32	[70.64, 72.01]	0.100	

*Number of Studies/ ** P-Value of Subgroup Difference

Discussion

The results of this study as the first meta-analysis on WHO-QOL-Brief revealed that the total mean score of the QOL among the Iranian elderly people was 60.1 ± 4.6 . In addition, the maximum and minimum-pooled mean score was obtained in the domains of social relationships and environmental health, respectively. The overall mean score of the QOL in the elderly population was less than the mean score of the QOL (76.95) in the general population of Iran (25). Moreover in our study, all mean scores were less than in comparison to Skevingto study (that included the information of 23 countries) (26), including; in physical health (54.6 vs. 63.7), in psychological health (57.3 vs. 63.1), in social relationship (57.9 vs. 63.7) and in environmental health (51.6 vs. 61.2).

Furthermore, another important result was that the elderly men had a markedly better QOL in the physical health domain (62.6 vs. 49.7). The difference was about 13 scores. The elderly men had a better status in the psychological domain

(62.4 vs. 55.9). The total pooled mean score of the QOL in men was significantly more than women (61.4 vs. 58.9). This finding was concordant with the results of some studies that evaluated the QOL using the WHO-QOL-Brief questionnaire (10, 11, 15) and other tools (12, 14, 16, 27). The reason for the better QOL in men rather than women may be due to the better socioeconomic status and having the role of guardianship in the families in developing countries, and more communication and interaction in the community. In addition, the latter result was concordant with another study in Iran (9) showed female gender significantly associated with a lower quality of life. However, the male gender is associated with a higher level of education, higher socioeconomic status, and better job position in the Iranian elderly population, and these factors are associated with the better QOL among men. In addition, the pooled mean score of the QOL in the physical health domain was more in men when compared with women. This finding was acceptable according to health statistics at the national level; for example, the prevalence of ob-

esity was 14.9% and 19% in men and women respectively, according to the results of the second round of Urban Heart Tehran Experience (28). The lower mean score of the QOL in women may be associated with less physical activity and higher prevalence of obesity in them.

Both genders had an equal status in the environmental domain of the QOL. This finding is similar to the results of a study (9) (in the adjusted and unadjusted analysis) as well as other studies (11, 15, 29) that suggest environmental factors have a negligible relationship with individual wellbeing.

We could not assess the association of demographic variables such as marital status, level of education, employment status, income, and living status (having a personal house or living in the nursing home) with the overall QOL because the included primary studies in this meta-analysis did not evaluate the association of these factors with the QOL. Therefore, we recommend that further primary studies should be conducted to assess the potential predictors of the QOL in the Iranian elderly population.

In this meta-analysis, the included studies were limited to three provinces of Iran (Tehran, Alborz and Tabriz). Eighty and twenty percent of the studies were conducted in Tehran and Alborz provinces, respectively. We did not find the similar studies in the elderly population in other provinces of Iran. The measurement and assessment of the QOL in the elderly population in different societies are necessary for decision making for the promotion of their health status and wellbeing. There appears to be a major gap in the quality of life research in other provinces of Iran. We performed sub-group analysis based on the potential source of heterogeneity, but heterogeneity remained in sub-groups. The considerable heterogeneity in results of included studies may be related to different settings of the studies (such as different geographic regions, with different cultures and lifestyles) consequently, the QOL may be affected by these factors. However, we pooled the results of included studies using the random effect model in order to estimate the overall QOL, because of public health impor-

tance of QOL for the elderly population and health policy makers. If the results of meta-analysis were a guide for health decision-making, it is possible to pool the results of heterogeneous studies (30).

Conclusion

The results of this study stress the necessity of attention to the quality of life in the domains of physical health (especially in women), and environmental health in the elderly population.

Ethical considerations

Ethical issues (Including plagiarism, informed consent, misconduct, data fabrication and/or falsification, double publication and/or submission, redundancy, etc.) have been completely observed by the authors.

Acknowledgement

We would like to thank the Vic-chancellor of Research and Technology of Tehran University of Medical Sciences (TUMS) for financial support of this study (Code: 94-01-27-28359). The authors declare that there is no conflict of interests.

References

1. Watkins K. Human Development Report 2005. New York: 2005 ISBN 0-19-530511-6.
2. World Health Organization. Ageing [04.08.2015]. <http://www.who.int/topics/ageing/en/>.
3. United Nations. Department of Economic and Social Affairs PD. World Population Ageing 2013. New York: 2013 ST/ESA/SER.A/348.
4. Statistical Centre of Iran. 2015 [30.07.2015]. Available from: <http://www.amar.org.ir/>.
5. World Health Organization. Measuring quality of life 1997. Available from: http://www.who.int/mental_health/media/68.pdf
6. Organization WH (1996). WHOQOL BREF: Introduction, administration, scoring. Geneva: WHO.
7. De Leo D, Diekstra RF, Lonqvist J, Lonqvist J, Cleiren MH, Frisoni GB, et al. (1998). LEIPAD,

- an internationally applicable instrument to assess quality of life in the elderly. *Behav Med*, 24(1):17-27.
8. Lyons RA, PERRY IM, LITTLEPAGE BN (1994). Evidence for the validity of the Short-form 36 Questionnaire (SF-36) in an elderly population. *Age Ageing*, 23(3):182-4.
 9. Ardalan A, Mazaheri M, Vanrooyen M, Mowafi H, Nedjat S, Naieni KH, et al. (2011). Post-disaster quality of life among older survivors five years after the Bam earthquake: implications for recovery policy. *Ageing soc*, 31(2):179-96.
 10. Ahangari M, Kamali M, Arjmand H M. (2007). quality of elderly's life in tehran senile culture house clubs. *Iran J Ageing*, 2(3):182-9.
 11. Hasani F, Kamali M, Akbarfahimi M, Davatgaran K (2011). Factors affecting quality of life of the elderly in the residential homes of Tehran (2009). *J Bijand Univ Med Sci*, 18(4):320-8.
 12. Hesamzadeh A, Maddah S, Mohammadi F, Fallahi Khoshknab M, M R (2010). Comparison of elderly's "quality of life" living at homes and in private or public nursing homes. *Iran J Ageing*, 4(14):66-74.
 13. Khooshemehri G, Kaviani M, Asmari M, Rajabian H, Naserkhaki V (2013). Comparison of quality of life for elderly men and women elderly nursing home resident by Alborz province in 2012. *Alborz University Med J*, 2(1):19-24.
 14. Nabavi S H, Shoja M, Mohammadi S, Rashedi V (2014). Health-related quality of life in community-dwelling older adults of Bojnourd, 2014. *J North Khorasan Univ Med Sci*, 6(2):433-39.
 15. Panaghi L, Abarashi Z, Mansoori N, Dehghani M (2010). Quality of Life and Related demographic factor of the elderly in Tehran. *Iran J Ageing*, 4(2):77-87.
 16. Saber M, NOSRATABADI M (2014). Social support and health-related quality of life in elderly people covered by the Welfare organization of Kerman city. *J Health Dev*, 3(3):189-99
 17. Von Elm E, Altman DG, Egger M, Pocock SJ, Gotsche PC, Vandenbroucke JP, et al. (2007). The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *BMJ*, 335(7624):806-8.
 18. Higgins JP, Thompson SG, Deeks JJ, Altman DG (2003). Measuring inconsistency in meta-analyses. *BMJ*, 327(7414):557-60.
 19. Higgins JP, Green S. *Cochrane handbook for systematic reviews of interventions*: Wiley Online Library; 2008.
 20. Dezfulimanesh M, Tehranian N (2005). Endocervical gonorrhoea in pregnant and non pregnant women and follow up of the infected cases in Kermanshah, Iran 2004. *Pak J Med Sci*, 21(3):313-7.
 21. DerSimonian R, Laird N (1986). Meta-analysis in clinical trials. *Control Clin Trials*, 7(3):177-88.
 22. Amir Hosseini K, Nedjat S. Comparison of quality of life for the elderly residents of nursing homes and elderly resident in personal home in Tehran in 2013. Tehran: Tehran University of Medical Sciences; 2013.
 23. Zaeri S, Asgharzadeh S, Holakouie Naeini K, Rahimi Foroushani A (2014). The Quality of Life and Its Effective Factors in the Elderly Living Population of Azerbaijan District, Tehran, Iran. *Iran J Epidemiol*, 9(4):66-74.
 24. Payahoo L, Khaje-bishak Y, Pourghasem B, Kabiralavi MB (2014). The survey of the relationship between quality of life of elderly with depression and physical activity in Tabriz, Iran. *J Rehab Med*, 2(2):39-46.
 25. Kufi F, Nedjat S, Yaseri M, cheraghi Z. Assessment of quality of life of the general population in different countries, according to the human development index.: Tehran University of Medical Sciences; 2016.
 26. Skevington SM, Lotfy M, O'Connell KA; WHOQOL Group (2004). The World Health Organization's WHOQOL-BREF quality of life assessment: psychometric properties and results of the international field trial. A report from the WHOQOL group. *Qual Life Res*, 13(2):299-310.
 27. Bazrafshan M, Hosseini MA (2008). Quality of elderly's life in shiraz ,jahandidegan club. *Iran J Ageing*, 3(1):33-41.
 28. Mahdavi MM, Asadi Lari M, Malek Afzali H. *Urban Heart Tehran Experience, Round II Result*. Tehran: 2010.
 29. Schroder-Butterfill E, Mariani R (2006). A framework for understanding old-age vulnerabilities. *Ageing Soc*, 26(1):9-35.
 30. Poole C, Greenland S (1999). Random-effects meta-analyses are not always conservative. *Am J Epidemiol*, 150(5):469-75.