



Prevalence of the Skipping Breakfast among the Iranian Students: A Review Article

*Mahin GHAFARI¹, Amin DOOSTI-IRANI², Masoud AMIRI³, *Zahra CHERAGHI²*

1. Dept. of Public Health, School of Health, Shahrekord University of Medical Sciences, Shahrekord, Iran
2. Dept. of Epidemiology, School of Public Health, Hamadan University of Medical Sciences, Hamadan, Iran
3. Social Health Determinants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

*Corresponding Author: Email: cheraghiz@ymail.com

(Received 04 Jan 2017; accepted 21 May 2017)

Abstract

Background: Adolescence is an important period since the establishment of dietary pattern can also affect the adulthood. This study aimed to estimate the overall prevalence of skipping breakfast among Iranian students.

Methods: The international and national databases, including Medline, Scopus, Science Direct, Embase, Web of sciences, Google Scholar, MagIran, and SID were searched 1945-2016 as per case. All studies addressing the prevalence of skipping breakfast among Iranian students were extracted.

Results: Out of 322 records, 24 articles remained for meta-analysis. The total pooled prevalence of skipping the breakfast was 0.216 (95% CI: 0.213-0.22), the girls had a higher percentage for skipping breakfast compared with boys (26% vs. 18%).

Conclusion: Skipping breakfast is more prevalent in girls. Interventions are required to promote breakfast consumption in the targeted Iranian students, especially the girls.

Keywords: Skipping breakfast, Dietary pattern, Systematic review, Iran

Introduction

In most countries, adolescents make up a large proportion of the population. According to the World Health Organization, in 2003, 19% of the world's total populations were adolescents aged 10 to 19 yr, and 84% of them were living in the developing countries (1). In Iran, based on the results of the 2011 Population and Housing Census, the proportion of adolescents aged 10 to 19 was 16.34% (2).

Adolescence is an important period since the establishment of nutritional habits during this time can also affect adulthood (2). In addition, breakfast is known as the most important daily meal (2), and consumption of it is considered as an important indicator of a healthy lifestyle (2). Since the time between dinner and breakfast is usually the longest period without absorption of

energy and nutrients, skipping breakfast may lead to metabolic changes and have a negative effect on cognitive performance (3). Eating breakfast has beneficial effects on the quality of the diet and prevents chronic disease; in contrast, skipping this meal increases risk of the metabolic syndrome and cardiovascular diseases (4, 5). So far, breakfast plays an important role in maintaining the health of children and adolescents (5).

Skipping breakfast has been reported in several countries, and skipping the breakfast meal was more common in children and adolescents (6). Some studies indicate a relation between skipping breakfast and the occurrence of obesity (7, 8). Others indicate the effect of basic characteristics (such as parent's education and income) on skipping breakfast (9, 10).

Therefore, limited and sporadic studies have been conducted to investigate the prevalence of skipping breakfast in the Iranian student population, and the results are inconsistent. This study aimed to estimate the overall prevalence of skipping breakfast among Iranian students.

Methods

Searching

Using the PRISMA guideline (11), extensive electronic and manual sources were investigated to identify relevant reviews. The international and national databases were searched using following keywords: “prevalence,” “skipping breakfast,” and “Iran.” International databases including Medline, Scopus, Embase, Web of sciences, and Google Scholar were searched up to Jun 2016. In addition, national databases including MagIran, Irandoc, Medlib, Iranmedex, and SID were searched from 1945-2016 as per case (Fig. 1).




Fig. 1: Flow chart depicting the stages of retrieving articles, checking eligibility criteria, and including the articles into the meta-analysis

Inclusion criteria

All cross-sectional studies that addressed the prevalence of skipping breakfast among Iranian pu-

pils were included irrespective of the time of the study and the language of the publication. The main outcome of interest was the prevalence of skipping breakfast.

Data extraction and management

Two authors (M Gh. and Z Ch.) screened the titles and abstracts of the retrieved citations independently; in the next stage, they reviewed the full text of the selected studies to select the studies that met the inclusion criteria for this review. In case of missing data, we contacted the corresponding authors of the studies. The same reviewers extracted the following variables for data analysis: the year of publication, location of the study (city), mean of age, gender, education, mean of body mass index, habitat of participants, sample size, and percent of skipping breakfast.

Assessment risk of bias

Five selected items from the Newcastle-Ottawa Scale checklist (12) were used for evaluating the quality of the included studies. The items included the following: 1) the statistical test used to analyze the data is clearly described and appropriate, and the measurement of the association is presented, including confidence intervals and the probability level (P -value), 2) assessment of the outcome, 3) The subjects in different outcome groups are comparable, based on the study design or analysis. Confounding factors are controlled, 4) sample size Justified and satisfactory, 5) Representativeness of the sample; truly representative of the average in the target population.

Studies that satisfied all mentioned criteria were classified as high quality. Studies that did not satisfy one item were classified as moderated quality, and studies that did not satisfy more than one item were classified as low quality.

Assessment of heterogeneity

Statistical heterogeneity was explored using the chi-squared test at a significance level of 10%. I-square test was used to quantify the heterogeneity across the included studies (13). The variance between the studies was estimated using tau-squared statistics (14).

Assessment of the publication bias

We used funnel plot to investigate publication bias visually (14), as well as (15)(16) tests to assess publication bias statistically.

Estimation of summary measures

A meta-analysis was performed to estimate the percent of skipping breakfast among pupils. The inverse variance (IV) method was used for calculating the pooled estimations. Subgroup analysis was performed according to gender and the quality of the included studies. The Stata 11 (StataCorp, College Station, TX, USA) was employed for data analysis.

In order to deal with the bias caused by the size of the different populations studied (17), we used the sample size as the weight variable in the mean command in STATA.

Moreover, the meta-regression has been used to evaluate the potential factors on the heterogeneity (18). The random effect model (19) was used for data analysis and the results were reported with a 95% confidence interval.

Results

We retrieved 322 records; 104 references were excluded because of duplication, 166 references were excluded because they were not related to the objective of the review, and 36 references were excluded because they were not eligible to be included in the meta-analysis after checking the full text. Finally, 24 articles (20-43) remained for the meta-analysis (Fig. 1 and Table 1), which included 59292 Iranian pupils, aged 7–16 yr with a mean age of 11.66 ± 0.73 yr.

54.05% of the studies (20 studies) referred to all five STROBE items (low risk of bias) and 18.92% (seven studies) referred to four STROBE items (moderate risk of bias). Finally, 27.3% (10 studies) referred to maximum three items (high risk of bias). The results of chi-squared and I-square tests indicate high heterogeneity for the overall prevalence of skipping breakfast. The mentioned heterogeneity test results were as follows: I-squared=99.4% and Chi squared=5953.44, $P<0.001$. Therefore, we did not report the pooled summary for the overall prevalence of skipping the breakfast.

Table 1: Characteristic of included studies in the systematic review

Author	Publication Year	City	Gender	Education	Number of cases	Sample Size	Prevalence (%)
Lotfi	2012	Zahedan	Boy	Primary	13	223	0.06
Cakirglu	2007	Tabriz	Both	Primary	12	160	0.08
Jagari	2013	Tehran	Boy	High School	6	300	0.02
Amini	2007	Tehran	Both	High School	37	389	0.10
Alimoradi	2014	Sannadaj	Both	High School	41	553	0.07
Moghadam	2011	Ghazvin	Both	Primary	136	1300	0.10
Gholami	2014	Tehran	Girl	Primary	30	164	0.18
Mortazavi	2010	Zahedan	Boy	All level	67	1278	0.05
Rashidi	2007	Tehran	Both	High School	179	2321	0.08
Djalalinia	2013	Tehran	Girl	Middle School	492	1823	0.27
Bagherniya	2014	Shahinshahr	Girl	Middle School	53	172	0.31
Shahbazi	2013	Yazd	Both	High School	36	320	0.11
Rahimi	2011	Qom	Girl	Middle School	21	100	0.21
Baygi	2015	27 province	Both	Middle & High School	227	1092	0.21
Maddah	2009	Rasht	Boy	Primary	838	3551	0.24
Hajghanai	2015	Kerman	Both	Primary	56	320	0.18
Kelishadia	2015	30 province	Both	Middle School	2627	7320	0.36
Neamati	2003	Ardabil	Girl	Middle & Primary	103	611	0.17
Ahadi	2015	31 province	Both	All Level	2548	13486	0.19
Maddah	2009	Rasht	Both	Primary & High School	2784	8937	0.31
Maddah	2009	Guilan	Girl	High School	523	2090	0.25
Veghari	2012	Golestan	Both	Primary	254	3786	0.07
Karami	2015	Omideiyeh	Male	Primary	31	155	0.20
Karimi	2008	Semnan	Both	All Level	52	1193	0.04

* Prevalence of skipping breakfast

We used the meta-regression analyses via the random effect model to investigate potential sources of heterogeneity between the studies. According to the meta-regression analysis, some factors identified as possible factors that modified the results including gender, education, and the period of conducting the study.

We assessed the probability of publication bias using the funnel plot as well as Begg's and Egger's tests (Fig. 2). The studies scattered nearly symmetrically on both sides of the vertical line reflecting absence of publication bias. The results of Begg's and Egger's tests confirmed the absence of publication bias ($P=0.981$).




Fig. 2: Funnel plot of included studies

Subgroup Analysis

We developed a subgroup analysis according to identified variables in the meta-regression analysis (gender and the quality of reporting the studies). For the variables (education and the period of the studies), the high heterogeneity remained even after subgroup analysis, and then we limited a subgroup analysis just for gender and quality of reporting the studies.

Among the high-quality studies, the girls had a higher percent of skipping the breakfast compared with the boys, but this difference was not statistically significant (23% vs. 19%, $P=0.29$).

Among the moderate-quality studies, this difference was considerable and statistically significant (22% vs. 0.05%, $P=0.02$). Finally, in low-quality studies, the higher prevalence of skipping the breakfast among girls rather than boys was not statistically significant (12% vs. 0.07%, $P=0.25$) (Table 2). Finally, the overall skipping the breakfast was 0.216 (95% CI: 0.213-0.22) (Fig. 3).

Discussion

In this systematic review, we retrieved 24 studies that addressed the prevalence of skipping breakfast in Iranian pupils aged 7–16 yr. Because of the high heterogeneous results, we conducted a subgroup analysis based on the meta-regression analysis via random effect model. In the meta-regression analysis, some factors had potential effects on the heterogeneity such as gender, education, period of conducting the studies, and quality of studies.

Based on the subgroup analysis, in all levels of quality (high, moderate, and low), the prevalence of skipping breakfast among girls was more than boys were. However, the mentioned difference was statistically significant only in the moderate quality group. In addition, some primary studies that conducted in other countries also indicated greater prevalence of skipping the breakfast among girls in comparison to boys (11, 44, 45). According to the result of the researchers at Edinburg University, because of eating habits (e.g., Consumption of daily sugary drinks and snack type foods, such as sweets and crisps), girls skipped breakfast more often than boys (46).

Poor socioeconomic status was strongly associated to skipping breakfast in the Iranian population (4, 47, 48). In addition, eating breakfast was more common in pupils more educated parents and acceptable economic status. In this systematic review, we could not assess the prevalence of skipping breakfast by socio-economic status since the majority of the studies did not assess the socio-economic status of pupils.




Fig. 3: Forrest Plot of prevalence of skipping breakfast by gender

Table 2: Subgroup analysis of prevalence of skipping breakfast according the quality of studies and gender

Quality of studies	Categories	n (%)	Pooled Prevalence	95% CI	I ²	P-value**
Low Risk Bias	Boy	7 (0.38)	0.19	[0.11, 0.26]	0.100	0.29
	Girl	11 (0.62)	0.23	[0.15, 0.30]	0.100	
Moderate Risk of Bias	Boy	4 (0.57)	0.05	[0.04, 0.06]	0.00	0.02
	Girl	3 (0.43)	0.15	[0.08, 0.36]	0.99	
High Risk of Bias	Boy	5 (0.62)	0.07	[0.03, 0.11]	0.88	0.25
	Girl	3 (0.38)	0.12	[0.06, 0.18]	0.77	

** P-value of Subgroup Difference

One of the most important limitations of this study was that we did not assess the other important factors that affect breakfast skipping, such as socio-economic, behavioral (hours of sleeping at night, time of waking up in the morning, and level of physical activity), environmental (influence of friends and parents), and cognitive factors (perceived barriers, self-efficacy, and attitudes). Additionally, most studies did not report the mentioned information, while several observational studies showed some evidence of mentioned factors on skipping breakfast in different populations (3, 44, 49).

Conclusion

Skipping breakfast is more prevalent in girls than boys are. This study again highlighted the interventions required to promote breakfast consumption in the targeted Iranian students, especially in girls.

Ethical considerations

Ethical issues (Including plagiarism, informed consent, misconduct, data fabrication and/or falsification, double publication and/or submission, redundancy, etc.) have been completely observed by the authors.

Acknowledgements

The authors received no specific funding for this work.

Conflict of Interests

The authors declare that there is no conflict of interests.

References

1. United Nations Department of Economic and Social Affairs (2013). World Population Ageing 2013. <http://www.un.org/en/development/desa/population/publications/pdf/ageing/WorldPopulationAgeing2013.pdf>
2. Statistics Center of Iran (2011) Selection Results General Census of Population and Housing. <https://www.amar.org.ir/Portals/1/Iran/census-2.pdf>
3. Rampersaud GC, Pereira MA, Girard BL, et al (2005). Breakfast habits, nutritional status, body weight, and academic performance in children and adolescents. *J Am Diet Assoc*, 105:743-760.
4. Shafiee G, Kelishadi R, Qorbani M, et al (2013). Association of breakfast intake with cardiometabolic risk factors. *J Pediatr (Rio J)*, 89:575-82.
5. Mekary RA, Giovannucci E, Willett WC, et al (2012). Eating patterns and type 2 diabetes risk in men: breakfast omission, eating frequency and snacking. *Am J Clin Nutr*, 95:1182-9.
6. Howden J, Chong Y, Leung S, et al (1993). Breakfast practices in the Asian region. *Asia Pac J Clin Nutr*, 2:77-84.
7. Horikawa C, Kodama S, Yachi Y, et al (2011). Skipping breakfast and prevalence of overweight and obesity in Asian and Pacific regions: a meta-analysis. *Prev Med*, 53:260-7.
8. Dubois L, Girard M, Kent MP, et al (2009). Breakfast skipping is associated with differences in meal patterns, macronutrient intakes and overweight among pre-school children. *Public Health Nutr*, 12:19-28.
9. Vereecken C, Dupuy M, Rasmussen M, et al (2009). Breakfast consumption and its socio-demographic and lifestyle correlates in schoolchildren in 41 counties participating in the HBSC study. *Int J Public Health*, 54 Suppl 2:180-90.
10. Szajewska H (2010). Systematic review demonstrating that breakfast consumption influences body weight outcomes in children and adolescents in Europe. *Crit Rev Food Sci Nutr*, 50:113-119.
11. Abalkhail B, Shawky S (2002). Prevalence of daily breakfast intake, iron deficiency anaemia and awareness of being anaemic among Saudi school students. *Int J Food Sci Nutr*, 53:519-528.
12. Wells G, Shea B, O'connell D, et al (2000) The Newcastle-Ottawa Scale (NOS) for assessing

- the quality of nonrandomised studies in meta-analyses.
http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp
13. Higgins JP, Thompson SG, Deeks JJ, Altman DG (2003). Measuring inconsistency in meta-analyses. *BMJ*, 327:557-60.
 14. Higgins JP, Green S (2008). Cochrane handbook for systematic reviews of interventions. ed. Wiley Online Library.
<http://training.cochrane.org/handbook>
 15. Begg CB, Mazumdar M (1994). Operating characteristics of a rank correlation test for publication bias. *Biometrics*, 50:1088-101.
 16. Egger M, Smith GD, Schneider M, Minder C (1997). Bias in meta-analysis detected by a simple, graphical test. *BMJ*, 315:629-634.
 17. Pitblado J (2009) Survey data analysis in Stata. Canadian Stata Users Group Meeting. Stata-Corp LP Toronto, Canada.
http://www.stata.com/meeting/canada09/ca09_pitblado_handout.pdf
 18. Thompson SG, Higgins J (2002). How should meta-regression analyses be undertaken and interpreted? *Stat Med*, 21:1559-73.
 19. DerSimonian R, Laird N (1986). Meta-analysis in clinical trials. *Control Clin Trials*, 7:177-88.
 20. Ahadi Z, Qorbani M, Kelishadi R, et al (2015). Association between breakfast intake with anthropometric measurements, blood pressure and food consumption behaviors among Iranian children and adolescents: the CASPIAN-IV study. *Public Health*, 129:740-7.
 21. Alimoradi F, Barikani A, Mohammadpoor Asl A, Javadi M (2014). Evaluation of association of the breakfast and demographic factors among the adolescents 14 to 18 years old in Sanandaj in 1392. *J Neyshabur Univ Med Sci*, 2:64-57.
 22. Amini M, Omidvar N, Kimiagar M (2007). Prevalence of overweight and obesity among junior high school students in a district of Tehran. *J Res Med Sci*, 12:315-319.
 23. Bagherniya M, Keshavarz SA, Mostafavi F, et al (2014). Using social cognitive theory in predicting meal frequency in overweight and obese Iranian adolescents. *Bull Env Pharmacol Life Sci*, 3:197-203.
 24. Baygi F, Heshmat R, Kelishadi R, et al (2015). Regional Disparities in Sedentary Behaviors and Meal Frequency in Iranian Adolescents: The CASPIAN-III Study. *Iran J Pediatr*, 25:e182.
 25. Cakiroglu FP, Malek M (2007). Eating habits of 7-12 year-old children in Tabriz, Iran. *Pakistan J Nutr*, 6:430-435.
 26. Djalalinia S, Ramezani-Tehrani F, Malekafzali H, et al (2013). Development and evaluation of a nutritional health program for adolescents. *Iran J Nurs Midwifery Res*, 18: 425-427.
 27. Gholami M, Mohammadi-Asl L (2015). The relations between breakfast consumption, body composition and cardiorespiratory fitness in school children. *J Food Technol Nutr*, 12:63-70.
 28. Hajghani A, Olumi F, Dustan A (2015) The study of breakfast habits and adverse factors affecting learning mathematics in primary school students in the academic year 93-92 Kerman. National Conference on Literacy and Health Promotion, Mashhad-Iran.
 29. Jafari F, Rezaeipor N, Gerayeli Malek V, et al (2013). Association Between Breakfast Consumption and Math Scores of High School Students in Tehran. *Biomed Pharmacol J*, 6:285-291.
 30. Karami K, Ghaleh S (2015). Breakfast and snack patterns of primary school girls in Omidieh, Khuzestan Province, Iran. *J Prev Med*, 2:67-72.
 31. Karimi B, Hashemi MS, Habibian H (2008). Study of the breakfast habits and its relationship with some factors in Semnan (Iran) pupils. *Koomesh*, 9:285-291+41.
 32. Kelishadi R, Qorbani M, Motlagh ME, et al (2016). Association of eating frequency with anthropometric indices and blood pressure in children and adolescents: the CASPIAN-IV Study. *J Pediatr (Rio J)*, 92:156-67.
 33. Lotfi B, Rakhshani F (2014). Knowledge and perceived threat of students in relationship with their behavior in context of consumption of breakfast and snack in primary boy schools in zahedan. *Payesh*, 13:61-71.
 34. Maddah M, Nikooyeh B (2010). Factors associated with overweight in children in Rasht, Iran: gender, maternal education, skipping breakfast and parental obesity. *Public Health Nutr*, 13:196-200.
 35. Maddah M, Rashidi A, Mohammadpour B, et al (2009). In-school snacking, breakfast consumption, and sleeping patterns of normal and overweight Iranian high school girls: a

- study in urban and rural areas in Guilan, Iran. *J Nutr Educ Behav*, 41:27-31.
36. Maddah M (2009). Risk factors for overweight in urban and rural schoolgirls in Iran: Skipping breakfast and early menarche. *Int J Cardiol*, 136(2):235-8.
 37. Mortazavi Z, Roudbari M (2010). Breakfast consumption and body mass index in primary, secondary and high school boys in Zahedan 2005-2006. *Iran J Endocrinol Metab*, 12:345-351.
 38. Nemati A, Sagha M, Nouzad Charvadeh H, Dehghan M (2003). Evaluation of eating breakfast among adolescent girl students in Ardabil, 1999-2000. *J Ardabil Univ Med Sci*, 3:39-46.
 39. Rahimi T, Dehdari T, Ariacian N, Gohari M (2012). Survey of breakfast consumption status and its predictors among Qom students based on the Penders health promotion model constructs. *Iran J Nutr Sci Food Technol*, 7:75-84.
 40. Rashidi A, Mohammadpour-Ahramjani B, Karandish M, et al (2007). Obese and female adolescents skip breakfast more than their non-obese and male peers. *Cent Eur J Med*, 2:481-487.
 41. Reza khani H, Soheili Azad A, Razaghi M, Nemati A (2012). Pattern of Breakfast and Snack Consumption and Their Effective Factors among Primary School Students, Qazvin. *J Health Hyg*, 2:57-63.
 42. Shahbazi H, Baghianimoghadamm MH, Khjeh Z, et al (2014). Survey of health and nutritional behaviors among high school students. *Iran J Health Educ Health Promot*, 1:69-80.
 43. Veghari G, Mansourian AR (2012). Breakfast consumption amongst school children in northern Iran. *J Nepal Paediatr Soc*, 32:193-200.
 44. Videon TM, Manning CK (2003). Influences on adolescent eating patterns: the importance of family meals. *J Adolesc Health*, 32:365-73.
 45. O'Dea JA, Caputi P (2001). Association between socioeconomic status, weight, age and gender, and the body image and weight control practices of 6-to 19-year-old children and adolescents. *Health Educ Res*, 16:521-32.
 46. Edinburge TUo (2016) Girls skip breakfast 'more often'. <http://www.ed.ac.uk/news/all-news/breakfast-150210>
 47. Sharifirad G, Yarmohammadi P, Azadbakht L, et al (2013). Determinants of fast food consumption among Iranian high school students based on planned behavior theory. *J Obes*, 2013: 147589.
 48. Keski-Rahkonen A, Kaprio J, Rissanen A, et al (2003). Breakfast skipping and health-compromising behaviors in adolescents and adults. *Eur J Clin Nutr*, 57:842-53.
 49. Pearson N, Atkin AJ, Biddle SJ, et al (2009). Patterns of adolescent physical activity and dietary behaviours. *Int J Behav Nutr Phys Act*, 6:45.